

Socialization of children with autism spectrum disorder complicated by concomitant disorders

Rymkhanova Ainagul Rymbekovna¹, Turebayeva Klara Zhamanbaevna², Rakhmetova Anar Muratovna³, Olexyuk Zaryana Yaroslavovna¹, Yakovenko Irina Valeryevna¹, Zhussupbekova Zamzagul Dostiarovna^{1*}

¹Department of Special and Inclusive Education, Karaganda University named after Academician E.A. Buketov, Karaganda, Republic of Kazakhstan. ²Department of Pedagogical and Special Education, K.Zhubanov Aktobe Regional University, Aktobe, Republic of Kazakhstan. ³Republican Research Institute for Labor Protection of the Ministry of Labor and Social Protection of the Population of the Republic of Kazakhstan, Astana, Republic of Kazakhstan.

Correspondence: Zhussupbekova Zamzagul Dostiarovna, Department of Special and Inclusive Education, Karaganda University named after Academician E.A. Buketov, Karaganda, Republic of Kazakhstan. Dostiarovna@mail.ru

Received: 06 March 2025; **Revised:** 20 May 2025; **Accepted:** 10 June 2025

ABSTRACT

The article describes the difficulty of socialization and adaptation of children with autism spectrum disorders (ASD) complicated by concomitant disorders to the conditions of modern society. Considering that the integration process of children with ASD into the social environment is a multifaceted process of personality development and adaptation in a social atmosphere, the authors conducted an experimental approbation of the socialization model for children with autism spectrum disorders (ASD) complicated by concomitant disorders. The socialization model contains modules that reveal issues such as organizational and substantive. The main directions of the socialization model are technologies, principles of support and guidance, competence of the specialists, indicating the necessary criteria for assessing socialization, taking into account the characteristics of children with ASD, complicated by concomitant disorders, in conditions of special and inclusive education. According to the experimental study, established the socialization model aimed to increase the independence of children with ASD, the ability to organize social relationships, and participate in educational and extracurricular activities. The implementation (application) of the model makes it possible to normalize and improve quality of life of children with ASD, complicated by concomitant disorders, and their integration into the educational environment.

Keywords: Socialization model, Autism spectrum disorders, Mental retardation, Concomitant disorders, Education and upbringing of children with ASD, Impaired communication and social interaction

Introduction

The difficulties of socialization for children with autism spectrum disorders (ASD) is one of the urgent problems of childhood that requires the close attention of pedagogical, psychological, and

medical sciences. Across the world, the increasing number of children with autism spectrum disorder is frightening. According to international statistics, ASD exists in one child out of 100. The data of psychological, medical, and pedagogical consultations (PMPC), in the Karaganda region confirm the trend of stable growth in the number of children with autistic spectrum. For example, in 2017 107 children, however in 2023, it grows up to 967 [1-8].

At the 10th revision of the International Classification of Diseases (ICD), ASD was introduced by a medical diagnosis of F84.0 Childhood autism. At the ICD-11th revision, six types of ASD are presented under the code 6A02 (6A02.0-6A02.5) indicating the presence/absence of intellectual disability, functional language, and functional speech. Generally, the ICD-11 revision

Access this article online

Website: www.japer.in

E-ISSN: 2249-3379

How to cite this article: Rymbekovna RA, Zhamanbaevna TK, Muratovna RA, Yaroslavovna OZ, Valeryevna YI, Dostiarovna ZZ. Socialization of children with autism spectrum disorder complicated by concomitant disorders. *J Adv Pharm Educ Res.* 2025;15(3):61-8. <https://doi.org/10.51847/8cQnTPYjYg>

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

refers to ASD complicated by concomitant disorders. The latest data from the World Health Organization (WHO) indicates that autism exists in about one child out of 100. However, according to the results of early studies, the real numbers are much higher. The prevalence of autism spectrum in third-world and developing countries is unknown [9-19].

Moreover, from the latest data from the US Center for Disease Control and Prevention (CDC), autism spectrum disorders have been registered in 1 out of 36 children (aged 8 years). Autism has been detected in 4% of boys and 1% of girls. Since 2021, the growth in prevalence has increased by 22% and 37.9% of children have intellectual disability [20-27].

In the Republic of Kazakhstan, as reported by the Psychological, medical and Pedagogical Consultations Center (PMPC) by the time of January 1st, 2024, 16,710 children and adolescents with autism spectrum were counted, which is 8.2% of the total number of children with special educational needs. As a result, 7.6% (1,278) of young children, 39.6% (6,616) of preschool children, and 52.8% (8816) of school-age children. There are 2,943 children enrolled in special educational institutions, 4,181 in inclusive education centers, and 1,226 children studying at home conditions. In comparison with the data for 2023, the growth was up to 1,670 children.

Today, there are no official statistical data reflecting the intellectual development of children with autism spectrum disorders and those with concomitant disorders, and believed that from unofficial reports, more than 40% of children have intellectual disability.

Statistics for the Republic of Kazakhstan reflect the global trend of an increase in the number of children with ASD. The growth of statistical data indicates a significant leap in the early detection of autism spectrum disorders, which is a significant step toward the successful socialization of children.

Frequently, ASD and concomitant disorders often occur with indications of hearing, vision, musculoskeletal disorders, intellectual disabilities, mental health issues, severe speech disorders, as well as endocrine and somatic diseases. The range of neurological disorders in children with ASD is wide and covers early delayed and disharmonious psychomotor development, sensory disorders, motor stereotypes, speech and articulation disorders, motor and vocal tics [28-33]. While choosing the methodical part of diagnostic and correctional care for children with ASD, it is necessary to remember the above features that lead to slow and disharmonious psychomotor development.

Currently, one of the priority requests of parents (legal representatives) is the introduction of children with autism spectrum disorders into the educational system of both special and secondary schools. The basis for the successful adaptation of students to the requirements of society and the staff of an educational organization is communication skills [34-49].

Thus, creating conditions for the development of communication capabilities and the gain of social skills for children with autism spectrum disorders, becoming a priority task for educational organizations.

Consequently, the process of education and socialization of children with ASD requires a thorough study of existing

problems: the current models of education and socialization of children with mental health and intellectual disability mechanically transferred to children with ASD without taking into account the peculiarities of their development. Lack of experience at the level of primary and secondary educational systems, no approaches have been developed for designing variable educational programs for children with ASD in conditions of special and inclusive education, complicated by concomitant disorders. Furthermore, the study of the socialization issues for children with ASD complicated by concomitant disorders, and possible ways to solve problems are justified by the stable growth of this group of children.

Materials and Methods

The study aims to analyze the effectiveness of the socialization model for children with autism spectrum disorders complicated by concomitant disorders.

Furthermore, the article used such a research method as a pedagogical experiment, divided into three categories such as ascertaining, formative, and control stages.

The pedagogical experiment was carried out in 3 general education schools (schools No. 25, 27, 5, and 12 children in total) and 6 special schools (No. 4 and No. 6 in the city of Karaganda, No. 10 in the city of Balkhash, No. 9 in the city of Saran, No. 5 in the city of Temirtau, No. 7 city of Shakhtinsk, 52 children in total) in the Karaganda region. The total number of children with ASD complicated by concomitant disorders amounted to 64 individuals, among them 20 children with communication and social interaction disorders and mental retardation (group 1); 20 school students - with communication and social interaction disorders and mild intellectual disabilities (group 2); 24 - with communication and social disorders interactions and moderate intellectual disabilities (group 3). It is important to note that in the experiment, the pilot group.

During the pedagogical experiment, four levels of socialization skills were identified in children with ASD complicated by concomitant disorders. First, maladaptive (low or no motivation to learn, often with facts of undesirable behavior, low level of the cognitive sphere), minimal (low motivation to learn, superficial or formal attitude to communication, demonstration of auto-aggression, aggression, self-stimulation, situational understanding of speech). Intermediate (formation of motivation for learning, rare facts of undesirable behavior, socially acceptable behavioral skills). Lastly, optimal (average level of motivation to learn, absence of facts of undesirable behavior, sufficient level of cognitive development, proficiency in verbal speech or means of alternative communication).

To determine the initial level of socialization, several diagnostic tools were used, where research methods of the emotional-volitional and cognitive spheres were applied and interpreted taking into account the communicative and intellectual characteristics of each child with ASD complicated by concomitant disorders. The leading diagnostic tools were

methods of observation, questioning, and questioning of parents (and legal representatives).

The level of socialization of children with ASD is determined by comparing the results obtained during the diagnosis. According to the results, the level of socialization of children with ASD was determined as (maladaptive, minimal, intermediate, or optimal). During the control stages of the pedagogical experiment, diagnostic methods were used by authors such as N. G. Luskanova (2012), A. I. Barkan (1987), A. Abrams (2016), A. V. Khaustova (2014), A. R. Luria (1995), J. Ravena (1936), K. Machover and F. Gudinafa (2002).

The socialization diagnostics of children with ASD complicated by concomitant disorders using the above methods at the confirming stage showed the following distribution of children according to the selected four levels. Each level is characterized by a different degree of severity. Thus, at the maladaptive and minimal levels, a low degree of severity is the best result for children in the group. Respectively, a high degree of severity is the statement of the worst results. At the intermediate and optimal level, on the contrary, a low degree implies the worst results and a high degree implies the best results for children (Table 1).

Results and Discussion

Table 1. Distribution of children by socialization levels at the ascertaining stage of the experiment

Group of examinees	Maladaptive level: degree of severity			Minimum level: degree of severity			Intermediate level: degree of severity			Optimal level: degree of severity		
	Low	Med	High	Low	Med.	High	Low	Med.	High	Low	Med.	High
Group 1 (n=20)	5%	10%	0%	20%	10%	15%	5%	5%	0%	0%	5%	25%
Group 2 (n=20)	10%	10%	30%	35%	5%	10%	0%	0%	0%	0%	0%	0%
Group 3 (n=24)	8%	8%	42%	38%	4%	0%	0%	0%	0%	0%	0%	0%

Analyzing the experiment results discovered that the optimal level of socialization was revealed in six children only in the first group – children with a combination of ASD and mental retardation. These children learn at the basic level of education, receive special psychological and pedagogical assistance from psychologists, speech pathologists, and speech therapists in a special boarding school for a long time.

An intermediate level was recorded in two children also only in the first group, whose intellectual state corresponded to the borderline, between mental retardation and mild intellectual impairment. Such children had an average level of motivation, mechanically, but followed the rules of behavior in the classroom, showed a situational interest in communication, understood speech and simple instructions. In addition, children possessed either alternative communication skills or verbal speech at the stage of formation (corresponded to the general underdevelopment of level I speech).

The minimum level was found in all three groups of 29 examinees. Teachers noted in their behavior, facts of undesirable behavior, auto-stimulation, aggression and auto-aggression, low memory level (children could not remember and reproduce

words, objects) and verbal speech was absent, situational understanding of speech, partial formation of self-service skills and neatness [50].

The maladaptive level was detected in three groups of 27 subjects. The children's behavior is characterized by aggression, auto-aggression, auto-stimulation, low levels of verbal and non-verbal intelligence, and short-term mechanical memory. Such children did not accept the rules of behavior in the group (they ran away from the classroom, violated school discipline, there were facts of aggression towards students and teachers), gross violations of educational activities or lack of formation were noted. The basic skills of neatness and self-service, are at the stage of formation or partially formed [51].

The allocation of children within each group according to different levels and degrees of severity and the results of the confirming stage of the experiment are presented in quantitative indicators in Table 1.

After the experiment aimed at increasing the level of socialization in children with ASD complicated by concomitant disorders, a repeated diagnosis was carried out based on the developed Model, which showed the following changes, shown in Table 2.

Table 2. Allocation of children with ASD by levels of socialization at the control stage of the experiment

Group of examinees	Maladaptive level: degree of severity			Minimum level: degree of severity			Intermediate level: degree of severity			Optimal level: degree of severity		
	Low	Med	High	Low	Med.	High	Low	Med.	High	Low	Med.	High
Group 1 (n=20)	0	5%	0	15%	15%	0	5%	0	25%	0	10%	25%
Group 2 (n=20)	0	5%	30%	35%	10%	0	0	0%	20%	0	0	0
Group 3 (n=24)	4%	4%	38%	13%	33%	0	0	0	8%	0	0	0

A high degree of severity of the optimal level was found in seven children of the first group. Behavioral changes: cases of undesirable behavior were no longer recorded, acceptable norms of behavior in the classroom formed, understanding of speech improved, the level of development of nonverbal and verbal intelligence increased, and a steady interest in communication appeared. The rest of the children who were at this stage of socialization began consciously performing with the norms of behavior in the classroom, and their level of motivation increased [52, 53].

Twelve children from three groups reached the intermediate level of varying severity. Such dynamics occurred due to a decrease in the number of recorded facts of undesirable behavior, aggression, and auto-aggression; more acceptable social behavior skills were established. Speech in some children began to correspond to the general underdevelopment of speech of I-II levels and I. The nature of the implementation of techniques that determine verbal and (or) non-verbal intelligence has moved to a medium-low level (but still within the framework of their general intellectual development) [54].

The minimum level of varying severity was found in 26 children from different three groups. There are some positive qualitative changes in children's behavior. The improvement occurred due to the gradual formation of skills of neatness and self-service, the time of "retention" in the classroom increased by more than 10-15 minutes, the first prerequisites for the possession of non-

verbal means of communication appeared, anxiety, aggression, and auto-aggression decreased, etc [55].

The maladaptive level is still observed in 19 children. By the end of the study, one of the students having not mastered the general educational curriculum for children with mental retardation transferred on the recommendation of the PMPC. In order to, a make simpler training program without focusing on the requirements of the state mandatory standard of education (for children with mild intellectual disabilities). In children of this level, the dynamics in the motivational-value, cognitive-evaluative, socio-functional, and communicative components of social adaptation were unstable and insignificant [56, 57].

The allocation of children within each group according to different levels and severity, the results of the control stage of the experiment presented in quantitative indicators in **Table 2**.

On the other hand, a comparative assessment of the levels of socialization (maladaptive, minimal, intermediate, and optimal) was carried out in accordance with the levels of the Socialization Model in the confirming and control stages of the experiment (**Tables 3-6**).

The obtained data was statistically processed using the IBM SPSS Statistics 15 application program. The presence of a common intergroup variance was confirmed by single-factor analysis of variance (ANOVA). The intergroup variance is considered statistically significant at the $p < 0.01$ level. The results of a comparative analysis of the data on the determination of the maladaptive level are presented in **Table 3**.

Table 3. Comparative analysis of the results for determining the maladaptive level

Group of examinees		Maladaptive level					
		Ascertaining experiment			Final experiment		
		High	Medium	Low	High	Medium	Low
Group 1 (n=20)	M±m	19,35±1,5	6,0±1,1	4,0±0,8	12,8±2,1*	4,0±0,6*	8,2±1,1*
Group 2 (n=20)	M±m	37,4±5,6	6,2±0,9	3,4±1,1	27,7±1,1*	9,5±1,2*	9,9±1,3*
Group 3 (n=24)	M±m	4,4±0,8	4,8±1,3	1,9±0,8	2,3±1,0	2,9±0,7*	3,2±1,1*

Note: * - significance level $p < 0.01$

When analyzing the data obtained in the final experiment in group 1, children with impaired communication and social interaction with mild intellectual impairment had a high maladaptive level and made up to 12.8 ± 2.1 , average - 4.0 ± 0.6 , low - 8.2 ± 1.1 ($p < 0.01$, respectively). In the second group, children with impaired communication and social interaction

with an average intellectual disability had a high maladaptive 27.7 ± 1.1 , average - 9.5 ± 1.2 , and low - 9.9 ± 1.3 . Whereas, in the third group, only the average and low maladaptive levels of 2.9 ± 0.7 , and 3.2 ± 1.1 (respectively) were determined in patients with mental retardation. **Table 4** shows the results of a comparative analysis of the definition of the minimum level.

Table 4. Comparative analysis of the results for determining the minimum level

Group of examinees		Minimal level					
		Ascertaining experiment			Final experiment		
		High	Medium	Low	High	Medium	Low
Group 1 (n=20)	M±m	9,25±1,0	1,8±0,7	13,5±1,3	11,4±1,2*	5,1±0,9*	9,5±1,2*
Group 2 (n=20)	M±m	3,8±0,8	5,1±0,8	16,5±1,4	7,1±1,4*	8,1±1,0*	10,1±1,5*
Group 3 (n=24)	M±m	13,7±1,6	6,6±1,2	7,6±0,9	4,1±1,3*	2,4±0,7*	2,7±1,1

Note: * - significance level $p < 0.01$

When determining the confidence level in group 1, children with impaired communication and social interaction with mild intellectual impairment had a high minimum level equal to 11.4 ± 1.2 , average -5.1 ± 0.9 , and low -9.5 ± 1.2 ($p < 0.01$, respectively). In the second group, children with impaired communication and social interaction with an average intellectual

disability had a high minimum level of 7.1 ± 1.4 , an average of 8.1 ± 1.0 , and a low of 10.1 ± 1.5 . Whereas, in the third group, only high and average minimum levels of 4.1 ± 1.3 , 2.4 ± 0.7 (respectively) were determined in patients with mental retardation. **Table 5** reports the results of a comparative analysis of the intermediate-level definition.

Table 5. Comparative analysis of the results for determining the intermediate-level

Group of examinees		Intermediate level					
		Ascertaining experiment			Final experiment		
		High	Medium	High	High	Medium	Low
Group 1 (n=20)	M±m	4,0±0,8	4,8±0,6	3,7±0,8	7,6±1,1*	7,7±0,9*	1,5±0,6*
Group 2 (n=20)	M±m	3,2±1,3	2,5±0,8	1,5±0,6	4,5±0,8*	5,9±1,0	3,1±1,2
Group 3 (n=24)	M±m	8,4±0,9	13,2±0,8	11,9±0,*8	13,0±1,1*	15,0±0,7*	7,8±1,3*

The reliability level in group 1 in children with impaired communication and social interaction with mild intellectual impairment was set at a high intermediate level and amounted to 7.6 ± 1.1 , average -7.7 ± 0.9 , low -1.5 ± 0.6 ($p < 0.01$, respectively). In the second group, children with impaired communication and social interaction with an average intellectual

disability had a high intermediate level of 4.5 ± 0.8 ($p < 0.01$). Whereas, in the third group, only a high intermediate level of 13.0 ± 1.1 , average -15.0 ± 0.7 , low -7.8 ± 1.3 ($p < 0.01$) was determined in patients with mental retardation. **Table 6** shows the results of a comparative analysis of determining the optimal level.

Table 6. Comparative analysis of the results to determine the optimal level

Group of examinees		Optimal level					
		Ascertaining experiment			Final experiment		
		High	Medium	High	High	Medium	Low
Group 1 (n=20)	M±m	3,3±3,8	1,5±0,7	1,7±0,7	3,6±0,9*	2,6±0,7*	1,2±0,4*
Group 2 (n=20)	M±m	1,9±2,2	1,4±0,6	1,5±0,6	2,1±0,6*	2,0±0,7*	1,5±0,6*
Group 3 (n=24)	M±m	7,7±1,1	5,15±0,7	8,9±1,0	12,1±3,9	10,8±1,1*	1,9±0,7*

Note: * - significance level $p < 0.01$

When determining the confidence level in group 1, children with impaired communication and social interaction with mild intellectual impairment had a high optimal level and amounted to 3.6 ± 0.9 , average -2.6 ± 0.7 , low -1.2 ± 0.4 ($p < 0.01$, respectively). In the second group, children with impaired communication and social interaction with an average intellectual disability had a high optimal level of 2.1 ± 0.6 ($p < 0.01$), an average of 2.0 ± 0.7 , and a low of 1.5 ± 0.6 . Whereas, in the third group, only the average optimal level of 10.8 ± 1.1 was determined in patients with mental retardation, and the low level was 1.9 ± 0.7 ($p < 0.01$).

Thus, the presence of a common intergroup variance is confirmed by the method of univariate analysis of variance (ANOVA) and the reliability of the results of experimental studies. There is a reason to reject the null hypothesis and accept an alternative one, which leads to an unambiguous conclusion. The level of socialization of children with ASD, complicated by concomitant disorders of the pilot group is higher, due to the use at the formative stage of the experiment of a model aimed at increasing children's independence, ability to establish social

relations, and participation in educational and extracurricular activities.

Conclusion

Thus, the recorded changes in the socialization level of children in the pilot group are positive. During the formative stage of the pedagogical experiment, the components of the socialization model of children with ASD complicated by concomitant disorders used the method of applied behavior analysis (ABA), the method of environmental adaptation, the TEACCH method, and the PECS system. Additionally the methods of social stories, sensory integration, and active inclusion in the process of children's socialization with ASD, as well as their parents and neuro-typical peers. While working with children, different tools adapted to the special educational needs of each child (visual schedule, visual instructions, cards, tokens, etc.) and used format of information delivering as educational, extracurricular, correctional, and developmental activities.

The research was carried out in the areas of correcting maladaptive behavior, forming a stereotype of educational

behavior, communication skills, and social interaction skills, also expanding ideas about self-analyzing and social environment, and developing social orientation. It is based on the implementation of the principles of the socialization model. The principle of focusing on the special educational needs of children made it possible to achieve a deep individualization process. Moreover, the principle of consistency in assessing the needs of children made it possible to respond promptly to the influence of various environmental factors. The interaction and coordination principles of the specialist's actions of the psychological and pedagogical support service, teachers, and parents (legal representatives) contributed to building a clear idea of actions. The principles of content variability and individualization of the psycho-corrective work, education, and priorities of the formation of life-long competencies influenced the achievement of each child.

Consequently, introducing the socialization model concluded that the model allows to normalize and improve the quality of life of most children with autism spectrum disorders (ASD) complicated by concomitant disorders. Important to note that positive and qualitative changes in the process of socialization depend not only on the created exact conditions that contribute to increasing the level of independence. Also, by the ability to establish social relations and participation in educational and extracurricular activities, but also largely on the nature and severity of the types of developmental disorders associated with ASD. In such cases, children need to prolong the period of study, for a longer learning and upbringing process to obtain a stable desired result. In the absence of positive dynamics or with insignificant results, recommended, based on international experience, to consider the introduction of "co-teaching", the inclusion of teaching assistants in the staff of special schools, which is due to the need for children in a deeply individualized support of the educational process. At present, Kazakhstan has already gained positive experience in accompanying children with teaching assistants in secondary schools, which, therefore, needs to transfer to the conditions of a special school.

Acknowledgments: We would like to thank the management and teaching staff of educational institutions in the cities and towns of Karaganda, Balkhash, Saran, Temirtau, and Shaktinsk for their participation in the study. We are also grateful to Gulmira Ablayevna Tussupbekova, Associate Professor of the Department of Biophysics, Biomedicine, and Neuroscience at Al-Farabi Kazakh National University; Valentina Vladimirovna Bobrova, Alfina Nigamatzyanovna Sakayeva, Assistant Professors at Karaganda University named after Academician E.A. Buketov; Zhanar Askerbekovna Yerzholova, and Bakhytgul Nurtayevna Bakmaganbetova, specialists from the Psychological, Medical, and Pedagogical Consultations No. 1 in Karaganda.

Conflict of interest: None

Financial support: The article was prepared with support of grant funding from the Ministry of Education and Science of the Republic of Kazakhstan for 2023-2025 on the topic AP19678254 "Modeling the Process of Socialization of Children with Autism Spectrum Disorders

Complicated by Concomitant Disorders in Special and Inclusive Education".

Ethics statement: The research was carried out in accordance with the ethical standards of the Republic of Kazakhstan; the research was conducted with the voluntary consent of parents and/or their legal representatives.

References

1. Meshcheriakov B, Zinchenko V. A large psychological dictionary. SPb.: Praim-Evroznak; 2004. [in Russian].
2. Shaw KA, Williams S, Patrick ME, Valencia-Prado M, Durkin MS, Howerton EM, et al. Prevalence and early identification of autism spectrum disorder among children aged 4 and 8 years — autism and developmental disabilities monitoring network, 16 sites, United States, 2022. *MMWR Surveill Summ.* 2025;74(No. SS-2):1-22. doi:10.15585/mmwr.ss7402a1
3. Daniel KS, Jiang Q, Wood MS. The Increasing prevalence of autism spectrum disorder in the U.S. and Its implications for pediatric micronutrient status: a narrative review of case reports and series. *Nutrients.* 2025;17(6):990. doi:10.3390/nu17060990
4. Issac A, Halemani K, Shetty A, Thimmappa L, Vijay V, Koni K, et al. The global prevalence of autism spectrum disorder in children: a systematic review and meta-analysis. *Osong Public Health Res Perspect.* 2025;16(1):3-27.
5. Omori M. Increased observation of predictable visual stimuli in children with potential autism spectrum disorder. *Sci Rep.* 2025;15(1):4572. doi:10.1038/s41598-025-89171-1
6. Chung S, Son JW. Visual perception in autism spectrum disorder: a review of neuroimaging studies. *Soa Chongsonyon Chongsin Uihak.* 2020;31(3):105-20. doi:10.5765/jkacap.200018
7. Kochhar P, Arora I, Bellato A, Ropar D, Hollis C, Groom MJ. A comparison of visual attention to pictures in the Autism Diagnostic Observation Schedule in children and adolescents with ADHD and/or autism. *Front Psychiatry.* 2024;15:1378593. doi:10.3389/fpsy.2024.1378593
8. Kaye AD, Allen KE, Smith Iii VS, Tong VT, Mire VE, Nguyen H, et al. Emerging treatments and therapies for autism spectrum disorder: a narrative review. *Cureus.* 2024;16(7):e63671. doi:10.7759/cureus.63671
9. World Health Organization. Autism spectrum disorders [Internet]. Available from: <https://www.who.int/ru/news-room/factsheets/detail/autism-spectrum-disorders>
10. Lord C, Brugha TS, Charman T, Cusack J, Dumas G, Frazier T, et al. Autism spectrum disorder. *Nat Rev Dis Primers.* 2020;6(1):5. doi:10.1038/s41572-019-0138-4
11. Qin L, Wang H, Ning W, Cui M, Wang Q. New advances in the diagnosis and treatment of autism spectrum

- disorders. *Eur J Med Res.* 2024;29(1):322. doi:10.1186/s40001-024-01916-2
12. Hodges H, Fealko C, Soares N. Autism spectrum disorder: definition, epidemiology, causes, and clinical evaluation. *Transl Pediatr.* 2020;9(Suppl 1):S55-S65. doi:10.21037/tp.2019.09.09
 13. Alnusayri AMF. Autism spectrum disorders (ASD) diagnosis and management; current trends and future direction: a review article. *Int J Clin Exp Med.* 2021;14(9):2271-80.
 14. Zareei M, Tohidast SA, Salmani M. Parents' perceptions of the problems in children with autism spectrum disorders: a qualitative study. *Iran Rehabil J.* 2022;20(4):611-22. doi:10.32598/irj.20.4.1772.1
 15. Syriopoulou-Delli CK. Advances in Autism Spectrum Disorder (ASD) diagnostics: from theoretical frameworks to ai-driven innovations. *Electronics.* 2025;14(5):951. doi:10.3390/electronics14050951
 16. Ousley O, Cermak T. Autism spectrum disorder: defining dimensions and subgroups. *Curr Dev Disord Rep.* 2014;1:20-8. doi:10.1007/s40474-013-0003-1
 17. Jensen AR, Lane AL, Werner BA, McLees SE, Fletcher TS, Frye RE. Modern biomarkers for autism spectrum disorder: future directions. *Mol Diagn Ther.* 2022;26(5):483-95. doi:10.1007/s40291-022-00600-7
 18. Al Khateeb JM, Kaczmarek L, Al Hadidi MS. Parents' perceptions of raising children with autism spectrum disorders in the United States and Arab countries: a comparative review. *Autism.* 2019;23(7):1645-54. doi:10.1177/1362361319833929
 19. Thayer N, Marsack-Topolewski C, Wilson K. Parental perceptions of community and professional attitudes toward autism. *J Autism Dev Disord.* 2024;1-6. doi:10.1007/s10803-024-06554-5
 20. Morales Hidalgo P, Voltas Moreso N, Canals Sans J. Autism spectrum disorder prevalence and associated sociodemographic factors in the school population: EPINED study. *Autism.* 2021;25(7):1999-2011. doi:10.1177/13623613211007717
 21. Salari N, Rasoulpoor S, Rasoulpoor S, Shohaimi S, Jafarpour S, Abdoli N, et al. The global prevalence of autism spectrum disorder: a comprehensive systematic review and meta-analysis. *Ital J Pediatr.* 2022;48(1):112. doi:10.1186/s13052-022-01310-w
 22. Kiselev Y, Handal M, Hjellvik V, Reichborn-Kjennerud T, Stoltenberg C, Suren P, et al. Nationwide study of neuropsychiatric comorbidity and medicines use in children with autism spectrum disorder in Norway. *Front Psychiatry.* 2020;11:596032. doi:10.3389/fpsy.2020.596032
 23. Jonsdottir SL, Saemundsen E, Thorarinsdottir EA, Rafnsson V. Evaluating screening for autism spectrum disorder using cluster randomization. *Sci Rep.* 2024;14(1):6855. doi:10.1038/s41598-024-57656-0
 24. Salehi M, Jaka S, Lotfi A, Ahmad A, Saecidi M, Gunturu S. Prevalence, socio-demographic characteristics, and co-morbidities of autism spectrum disorder in US children: insights from the 2020–2021 national survey of children's health. *Children.* 2025;12(3):297. doi:10.3390/children12030297
 25. Talantseva OI, Romanova RS, Shurdova EM, Dolgorukova TA, Sologub PS, Titova OS, et al. The global prevalence of autism spectrum disorder: a three-level meta-analysis. *Front Psychiatry.* 2023;14:1071181. doi:10.3389/fpsy.2023.1071181
 26. Khachadourian V, Mahjani B, Sandin S, Kolevzon A, Buxbaum JD, Reichenberg A, et al. Comorbidities in autism spectrum disorder and their etiologies. *Transl Psychiatry.* 2023;13(1):71. doi:10.1038/s41398-023-02374-w
 27. Maniram J, Oosthuizen F, Karrim SB. An overview of pharmacotherapy in the management of children with autism spectrum disorder at a public hospital in KwaZulu-Natal. *Child Psychiatry Hum Dev.* 2024;55:1655-63. doi:10.1007/s10578-023-01514-z
 28. Zavadenko NN, Pechatnikova NL, Simashkova Z, Zavadchenko AN, Orlova KA. Neurological disorders of children with autism. *Ros Vest Perinatol Pediatr.* 2015;2:14-20. [in Russian].
 29. Mekhdi M, Shopsha EL. Medical aspects of accompanying children with autism spectrum disorders. *Hum Balk Res.* 2021;2:19-21. [in Russian].
 30. Niukikten Ch. Pediatric behavioral neurology. *Terevinf.* 2010;2:336. [in Russian].
 31. Chaidez V, Hansen RL, Hertz-Picciotto I. Gastrointestinal problems in children with autism, developmental delays or typical development. *J Autism Dev Disord.* 2014;44:1117-27.
 32. Manelis NG, Nikitina IV, Ferroi LM, Komarova OP. Sensory characteristics of children with autism spectrum disorders. Assistance strategies. *Methodical Manual.* 2018. [in Russian].
 33. Tkachuk EA, Martynovich NN, Rychkova LV, Poliakov VM. Problems of diagnosis of autism spectrum disorders. *Revive.* 2020;4:35-43. [in Russian].
 34. Morozov SA, Morozova TI, Morozova SS. Autism: Experience and Possible Solutions to Current Problems. A Look from Russia. *Autism and Developmental Disorders.* 2021;19(3):42-8. (in Russian). doi:10.17759/autdd.2021190305
 35. Yano N, Hosokawa K. The importance of comprehensive support based on the three pillars of exercise, nutrition, and sleep for improving core symptoms of autism spectrum disorders. *Front Psychiatry.* 2023;14:1119142. doi:10.3389/fpsy.2023.1119142
 36. Doda V, Kennedy C, Kaur M. Policies for individuals with autism: gaps, research, and recommendations. *Cureus* 2024;16(1):e51875. doi:10.7759/cureus.51875
 37. Georgiou I, Parlalis SK. Social services and support structures for adults with autism spectrum disorder (ASD) and their parents' quality of life in Cyprus. In: Ciciurkaite G, Brown RL, eds. *Disability and the changing contexts of*

- family and personal relationships. Research in social science and disability. Vol. 15. Leeds: Emerald Publishing Limited; 2024. p. 103-20. doi:10.1108/S1479-354720240000015008
38. Evans K, Whitehouse AJO, D'Arcy E, Hayden-Evans M, Wallace K, Kuzminski R, et al. Perceived support needs of school-aged young people on the autism spectrum and their caregivers. *Int J Environ Res Public Health*. 2022;19(23):15605. doi:10.3390/ijerph192315605
 39. Dedmore LB. Adult independent living outcomes for persons with autism spectrum disorders [master's thesis]. 2013. doi:10.33015/dominican.edu/2013.edu.19
 40. Sorensen C, Zarrett N. Benefits of physical activity for adolescents with autism spectrum disorders: a comprehensive review. *Rev J Autism Dev Disord*. 2014;1:344-53. doi:10.1007/s40489-014-0027-4
 41. Zou M, Han Y, Jin M, Xie W, Shang C, Li T, et al. Supporting access to comprehensive services: a scoping review of national policies related to autism in mainland China. *Autism*. 2025;29(7):1646-60. doi:10.1177/13623613241311736
 42. Weiss JA, Robinson S, Riddell RP, Flora D. Understanding stability and change in perceived social support in parents of autistic children and adolescents. *Front Rehabil Sci*. 2021;2:679974. doi:10.3389/fresc.2021.679974
 43. Jia M, Zhang J, Pan J, Hu F, Zhu Z. Benefits of exercise for children and adolescents with autism spectrum disorder: a systematic review and meta-analysis. *Front Psychiatry*. 2024;15:1462601. doi:10.3389/fpsy.2024.1462601
 44. Kou R, Li Z, Li M, Zhou R, Zhu F, Ruan W, et al. Comparative effectiveness of physical exercise interventions on sociability and communication in children and adolescents with autism: a systematic review and network meta-analysis. *BMC Psychol*. 2024;12(1):712. doi:10.1186/s40359-024-02210-w
 45. Ranieri A, Mennitti C, Falcone N, La Monica I, Di Iorio MR, Tripodi L, et al. Positive effects of physical activity in autism spectrum disorder: how influences behavior, metabolic disorder and gut microbiota. *Front Psychiatry*. 2023;14:1238797. doi:10.3389/fpsy.2023.1238797
 46. Correale C, Borgi M, Cirulli F, Laghi F, Trimarco B, Ferraro M, et al. The impact of health and social services on the quality of life in families of adults with autism spectrum disorder (ASD): a focus group study. *Brain Sciences*. 2022;12(2):177. doi:10.3390/brainsci12020177
 47. Fulceri F, Gila L, Caruso A, Micai M, Romano G, Scattoni ML. Building bricks of integrated care pathway for autism spectrum disorder: a systematic review. *Int J Mol Sci*. 2023;24(7):6222. doi:10.3390/ijms24076222
 48. Shurtz L, Schwartz C, DiStefano C, McPartland JC, Levin AR, Dawson G, et al. Concomitant medication use in children with autism spectrum disorder: data from the autism biomarkers consortium for clinical trials. *Autism*. 2023;27(4):952-66. doi:10.1177/13623613221121425
 49. Watanabe S, Masamura N, Satoh S, Hirao T. Evaluating the effectiveness of DNA barcoding for insect identification: a comprehensive review. *Entomol Lett*. 2024;4(2):34-41. doi:10.51847/ZVNniNFsOR
 50. Dipalma G, Inchingolo AD, Fiore A, Balestriere L, Nardelli P, Casamassima L, et al. Comparative effects of fixed and clear aligner therapy on oral microbiome dynamics. *Asian J Periodontics Orthod*. 2022;2:33-41. doi:10.51847/mK28wdKCIX
 51. Pisano M, Sangiovanni G, Frucci E, Scorziello M, Benedetto GD, Iandolo A. Assessing the reliability of electronic apex locators in different apical foramen configurations. *Asian J Periodontics Orthod*. 2023;3:1-5. doi:10.51847/qOUk00kkRZ
 52. Bolay Ş, Öztürk E, Tuncel B, Ertan A. Studying fracture strength of root-treated and reconstructed teeth with two types of post and core. *Ann J Dent Med Assist*. 2024;4(2):1-6. doi:10.51847/i57dzmzc2A
 53. Bulusu A, Cleary SD. Comparison of dental caries in autistic children with healthy children. *Ann J Dent Med Assist*. 2023;3(2):14-9. doi:10.51847/wa2pZXE4RJ
 54. AlHussain BS, AlFayez AA, AlDuhaymi AA, AlMulhim EA, Assiri MY, Ansari SH. Impact of different antibacterial substances in dental composite materials: a comprehensive review. *Int J Dent Res Allied Sci*. 2022;2(1):1-7. doi:10.51847/jg2xu2PbJK
 55. Maneca ASB, Alqahtani AD, Alhazzaa AK, Albalawi AO, Alotaibi AK, Alanazi TF. Systematic review of the microbiological impact of sodium hypochlorite concentrations in endodontic treatment. *Int J Dent Res Allied Sci*. 2024;4(2):9-15. doi:10.51847/PH80PpWOX7
 56. Fernandes AL, Malik JB, Ansari SR, Murali S, Thirupathii J. Saudi dentists' knowledge and approaches to managing tooth wear: a cross-sectional survey-based analysis. *Turk J Public Health Dent*. 2022;2(2):1-12. doi:10.51847/p7ulFD4XZm
 57. Ravoori S, Sekhar PR, Pachava S, Pavani NPM, Shaik PS, Ramanarayana B. Perceived stress and depression among oral cancer patients - a hospital based cross-sectional study. *Turk J Public Health Dent*. 2024;4(1):1-5. doi:10.51847/FoK9xAl1JW