

The first boundary value problem for the fractional diffusion equation in a degenerate angular domain

M.I. Ramazanov¹, A.V. Pskhu², M.T. Omarov^{1,*}

¹*Institute of Applied Mathematics, Karaganda Buketov University, Karaganda, Kazakhstan;*

²*Institute of Applied Mathematics and Automation of Kabardin-Balkar Scientific Center of RAS, Nalchik, Russia
(E-mail: ramamur@mail.ru, pskhu@list.ru, madiomarovt@gmail.com)*

This article addresses the problems observed in branching fractal structures, where super-slow transport processes can occur, a phenomenon described by diffusion equations with a fractional time derivative. The characteristic feature of these processes is their extremely slow relaxation rate, where a physical quantity changes more gradually than its first derivative. Such phenomena are sometimes categorized as processes with “residual memory”. The study presents a solution to the first boundary problem in an angular domain degenerating into a point at the initial moment of time for a fractional diffusion equation with the Riemann-Liouville fractional differentiation operator with respect to time. It establishes the existence theorem of the problem under investigation and constructs a solution representation. The need for understanding these super-slow processes and their impact on fractal structures is identified and justified. The paper demonstrates how these processes contribute to the broader understanding of fractional diffusion equations, proving the theorem’s existence and formulating a solution representation.

Keywords: partial differential equation, fractional calculus, angular domain, kernel, weak singularity, parabolic cylinder, Carleman-Vekua equation, general solution, unique solution, Riemann-Liouville fractional operator.

2020 Mathematics Subject Classification: 35R11.

Introduction

The paper discusses an equation of the form:

$$\left(\frac{\partial^\alpha}{\partial t^\alpha} - \frac{\partial^2}{\partial x^2}\right) u(x, t) = f(x, t), \quad (0 < \alpha < 1), \tag{1}$$

where $\frac{\partial^\alpha}{\partial t^\alpha}$ is a fractional derivative of an order α with respect to the variable t , starting from the point $t = 0$. This type of fractional differentiation is defined by the Riemann-Liouville operator:

$${}_a g^{(\nu)}(x) \equiv {}_a D_x^\nu g(x) = \begin{cases} \frac{1}{\Gamma(-\nu)} \int_a^x (x - \xi)^{-\nu-1} g(\xi) d\xi, & \nu < 0, \\ \frac{1}{\Gamma(1-\nu)} D_x \int_a^x (x - \xi)^{-\nu} g(\xi) d\xi, & 0 \leq \nu < 1, \\ \frac{1}{\Gamma(2-\nu)} D_x^2 \int_a^x (x - \xi)^{-\nu+1} g(\xi) d\xi, & 1 \leq \nu < 2, \\ \dots\dots\dots, & \dots\dots\dots \end{cases}$$

Fractional diffusion equations (where $0 < \alpha \leq 1$) have been extensively studied in recent years. This surge in interest is due to their widespread applications in physics and modeling, as referenced in sources [1–5]. The primary methodologies for exploring diffusion-wave equations are detailed in publications [6–24], while monographs [25] and [26] provide a comprehensive bibliography on the subject.

*Corresponding author. E-mail: madiomarovt@gmail.com

This research was funded by the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan (Grant No. AP23488729).

Received: 18 October 2023; Accepted: 04 December 2023.

© 2024 The Authors. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Nearly all studies related to equation (1) have focused on initial and boundary problems in both limited and unlimited cylindrical areas. Specifically, the first boundary problem for the fractional diffusion equation in a rectangular area was examined in [17,18]. In publication [27], the first boundary problem for the fractional diffusion-wave equation in a non-cylindrical area was solved. However, the area where the solution is sought does not degenerate into a point at the initial moment in time.

The aim of this study is to solve the first boundary problem for equation (1) in a domain that is not cylindrical, but rather angular, and degenerates into a point at the initial moment in time.

In relation to the boundary value problems for the heat conduction equation with a diffusion coefficient α set to 1:

$$\left(\frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2}\right) u(x, t) = f(x, t),$$

these problems have been investigated in non-cylindrical domains by several authors [28–32]. It is important to underline that boundary value problems for the Laplace equation in domains with evolving boundaries are distinct from the classical ones defined in fixed cylindrical domains. The reason is that the dimensions of the domain where the solution is sought are time-dependent, which makes these problems unsuitable for classical variable separation and integral transformation methods.

The potential theory approach allows reformulating the boundary value problem into a Volterra system of second kind integral equations. In such cases, if the domain's boundary does not exist at the initial time, then the corresponding system of integral equations can be solved by the method of successive approximations due to the weak singularity of their kernels. In contrast, if the boundary exists at the initial time, the integral equations of the boundary value problem might admit additional solutions, and the implementation of the Picard method encounters certain mathematical complexities. Similar issues occur for boundary value problems of the Dirichlet problem for the Laplace equation in non-cylindrical domains that originate at the initial moment in time.

1 Problem Statement

To determine a regular solution for the fractional time-derivative heat equation:

$$\left(\frac{\partial^\alpha}{\partial t^\alpha} - \frac{\partial^2}{\partial x^2}\right) u(x, t) = f(x, t), \quad (0 < \alpha < 1),$$

within the domain

$$D = \{(x, t) : 0 < x < t, 0 < t < \infty\},$$

that adheres to the boundary conditions:

$$u(0, t) = 0, \quad u(t, t) = 0, \quad 0 < t < \infty. \quad (2)$$

We denote $u(x, t)$ as a regular solution of equation (1) in domain D such that:

$$t^{1-\gamma}u(x, t) \in C(\overline{D})$$

for some $\gamma > 0$. Additionally, the solution $u(x, t)$ must be continuous within D and possess a continuous partial derivative with respect to x , and its second-order derivative with respect to x , ${}_0D_t^\nu u(x, y)$, must be continuous in the variable t at fixed x inside the domain D and up to the boundary set $\{0 < x < t\}$, with $u(x, t)$ fulfilling equation (1) at every point in D .

2 Main Result

Definitions are introduced as follows:

$$\beta = \frac{\alpha}{2}, \quad \omega_{\beta,\mu}(x, t) = t^{\mu-1}W\left(-\beta, \mu; -\frac{|x|}{t^\beta}\right),$$

$$\omega(x, t) = \omega_{\beta,0}(x, t),$$

in which

$$W(-\beta, \mu; z) = \sum_{k=0}^{\infty} \frac{z^k}{k! \Gamma(\mu - \beta k)}$$

represents the Wright function, as discussed in [33].

The following statement holds true:

Theorem 1. Let the conditions be satisfied: $t^{1-\gamma}g_i(t) \in C[0, T]$, $i = 1, 2$, for some $\gamma > 0$, and $t^{1-\mu}f(x, t) \in C(\overline{D})$, $\mu \geq 0$, if $f(x, t)$ satisfies the Holder condition with respect to the variable x .

Then the solution to problem (1)-(2) exists and can be expressed as

$$u(x, t) = \int_0^t \psi_1(\tau)\omega(x, t - \tau)d\tau + \int_0^t \psi_2(\tau)\omega(T - x, t - \tau)d\tau + F(x, t), \tag{3}$$

here

$$F(x, t) = \frac{1}{2} \int_0^t \int_0^\tau f(s, \tau)\omega_{\beta,\mu}(x - s, t - \tau)dsd\tau,$$

and $\psi_1(t), \psi_2(t)$ from (3) are the solutions to the system of integral equations

$$\begin{cases} \psi_1(t) + \int_0^t \psi_2(\tau)\omega(\tau, t - \tau)d\tau = -F(0, t), \\ \psi_2(t) + \int_0^t \psi_1(\tau)\omega(t, t - \tau)d\tau + \int_0^t \psi_2(\tau)\omega(\tau - t, t - \tau)d\tau = -F(t, t). \end{cases} \tag{4}$$

From the first equation of this system (4), we obtain

$$\psi_1(t) = - \int_0^t \psi_2(\tau)\omega(t, \tau - t)d\tau - F(0, t),$$

and substituting $\psi_1(t)$ into the second equation of the system (4), we get [33]:

$$\begin{aligned} \psi_2(t) + \int_0^t \left(- \int_0^\tau \psi_2(\xi)\omega(\xi, \tau - \xi)d\xi - F(0, \tau) \right) \omega(t, t - \tau)d\tau + \\ + \int_0^t \psi_2(\tau)\omega(\tau - t, t - \tau)d\tau = -F(t, t). \end{aligned} \tag{5}$$

Substituting into the repeated integral and changing the order of integration as well as the dummy variables ξ and τ , from equation (5), we arrive at a special integral equation of the second kind in the form of a Volterra equation:

$$\psi_2(t) - \int_0^t \mathcal{K}(t, \tau)\psi_2(\tau)d\tau = \mathcal{F}(t), \tag{6}$$

here

$$\mathcal{K}(t, \tau) = \int_\tau^t \omega_{\beta,0}(\tau, \xi - \tau)\omega_{\beta,0}(t, t - \xi)d\xi - \omega_{\beta,0}(\tau - t, t - \tau), \tag{7}$$

and

$$\mathcal{F}(t) = \int_0^t F(0, \tau)\omega(t, \tau - t)d\tau - F(t, t). \tag{8}$$

To perform the calculation of integral (6) having (7) and (8), it is necessary to employ.

$$\int_{\tau}^t \omega_{\beta,0}(\tau, \xi - \tau)\omega_{\beta,0}(t, t - \xi)d\xi.$$

The convolution formula is applied to the Wright function as referenced in [33], and this is expressed through the function $\omega_{\beta,\mu}(x, t)$:

$$\int_0^y \omega_{\alpha,\delta}(x_1, \xi) \omega_{\alpha,\mu}(x_2, (y - \xi)) d\xi = \omega_{\alpha,\delta+\mu}(x_1 + x_2, y).$$

Then we obtain,

$$\begin{aligned} \int_{\tau}^t \omega_{\beta,0}(\tau, \xi - \tau)\omega_{\beta,0}(t, t - \xi)d\xi &= \|\xi - \tau = \eta\| = \\ &= \int_0^{t-\tau} \omega_{\beta,0}(\tau, \eta)\omega_{\beta,0}(t, (t - \tau) - \eta)d\eta = \\ &= \omega_{\beta,0}(t + \tau, t - \tau). \end{aligned}$$

Therefore, the conclusive kernel $\mathcal{K}_{\beta}(t, \tau)$ is determined by the following relation:

$$\mathcal{K}_{\beta}(t, \tau) = \omega_{\beta,0}(t + \tau, t - \tau) - \omega_{\beta,0}(\tau - t, t - \tau). \tag{9}$$

The second term of kernel (9) has a weak singularity, since the following estimate is valid for it:

$$\omega_{\beta,0}(\tau - t, t - \tau) \leq \frac{C(\beta)}{(t - \tau)^{\beta}}. \tag{10}$$

Indeed, by applying the estimate found in [26]:

$$\begin{aligned} |\omega_{\mu}(x, y)| &\leq C(\beta, \mu, \theta)|x|^{-\theta}y^{\beta\theta+\mu-1}, \\ \theta &\geq \begin{cases} 0, & (-\mu) \notin \mathbb{N} \cup \{0\} \\ -1, & (-\mu) \in \mathbb{N} \cup \{0\} \end{cases}, \end{aligned}$$

taking into account that $\mu = 0$, and choosing

$$\theta = -\frac{\beta}{1 - \beta} > -1.$$

This leads to the confirmation of inequality (10). Next, we aim to demonstrate the special nature of the kernel $\mathcal{K}_{\beta}(t, \tau)$.

Lemma. If $0 < \beta \leq 1/2$, the equality holds true

$$\lim_{t \rightarrow 0} \int_0^t \mathcal{K}_{\beta}(t, \tau)d\tau = 1. \tag{11}$$

Proof. Initially, when t is small, these inequalities are applicable:

$$\omega_{\beta,0}(t, t - \tau) \geq \omega_{\beta,0}(t + \tau, t - \tau) \geq \omega_{\beta,0}(2t, t - \tau).$$

Using equation [33]

$$D_{0t}^v \omega_{\beta,\mu}(x, y) = \omega_{\beta,\mu-v}(x, y)$$

these results in

$$\lim_{t \rightarrow 0} \int_0^t \omega_{\beta,0}(bt, t - \tau) d\tau = \lim_{t \rightarrow 0} \omega_{\beta,1}(bt, t) = 1, \quad b = 1, 2.$$

Therefore, considering inequality (10), we establish the validity of equality (11).

The kernel's properties make it unsuitable for solving the corresponding integral equation through the method of successive approximations. This limitation of the integral equation is due to the solution domain for the problem collapsing to a single point at the start. Otherwise, if this collapse didn't occur, the kernel for the integral equation would possess a weak singularity, enabling the use of Picard's method for finding a solution [33].

3 Solution of the special integral equation (6)

To solve the integral equation mentioned in equation (6), we apply the Carleman-Vekua method. This involves using a specific integral equation, which we refer to as the characteristic equation.

$$\psi_2(t) - \int_0^t \mathcal{K}_{\frac{1}{2}}(t, \tau) \psi_2(\tau) d\tau = \mathcal{Q}(t) \tag{12}$$

here

$$\mathcal{K}_{1/2}(t, \tau) = \frac{1}{2a\sqrt{\pi}} \left\{ \frac{t + \tau}{(t - \tau)^{\frac{3}{2}}} \exp\left(-\frac{(t + \tau)^2}{4a^2(t - \tau)}\right) + \frac{1}{(t - \tau)^{\frac{1}{2}}} \exp\left(-\frac{t - \tau}{4a^2}\right) \right\}. \tag{13}$$

Relation (13) can be verified directly using the following formula [34; 5.2.10(2)] for $(n = -2)$,

$$\sum_{k=0}^{\infty} \frac{(-1)^k x^k}{k! \Gamma[1 + (n - k)/2]} = \frac{1}{\sqrt{\pi}} 2^{(n+1)/2} e^{-x^2/8} D_{-n-1} \left(\frac{x\sqrt{2}}{2} \right),$$

here $D_{-n-1}(z)$ is the function of a parabolic cylinder.

At the same time, the kernel $\mathcal{K}_{\frac{1}{2}}(t, \tau)$ possesses a similar property as described in equation (11):

$$\lim_{t \rightarrow 0} \int_0^t \mathcal{K}_{\frac{1}{2}}(t, \tau) d\tau = 1.$$

This means that the kernel difference $\mathcal{K}_{\frac{1}{2}}(t, \tau) - \mathcal{K}_{\beta}(t, \tau) = \tilde{K}(t, \tau)$ has a weak singularity. We will employ the regularization method to solve the characteristic equation, known as the Carleman-Vekua equation, and to do so, we will express equation (12) in a particular form:

$$\psi_2(t) - \int_0^t \mathcal{K}_{\frac{1}{2}}(t, \tau) \psi_2(\tau) d\tau = \mathcal{Q}(t) - \int_0^t \tilde{K}(t, \tau) \psi_2(\tau) d\tau. \tag{14}$$

Assuming the right-hand side of this equality is temporarily known and denoting it by

$$\mathcal{Q}(t) = \mathcal{F}(t) - \int_0^t \tilde{K}(t, \tau) \psi_2(\tau) d\tau.$$

Equation (14) can be represented in the following form:

$$\mathbb{K}_{1/2}\psi_2 \equiv \psi_2(t) - \int_0^t \mathcal{K}_{\frac{1}{2}}(t, \tau)\psi_2(\tau)d\tau = Q(t). \quad (15)$$

In [35] it is shown that the general solution of equation (15) in the weight class of functions

$$\sqrt{t} \exp\left(-\frac{t}{4a^2}\right)\varphi(t) \in L_\infty(0, \infty)$$

has the form:

$$\mathbf{K}\psi_2 \equiv \psi_2(t) - [\mathbb{K}_{\frac{1}{2}}]^{-1} Q(t) = c_0\psi_0(t) \quad (16)$$

and the function

$$\psi_0(t) = \frac{1}{\sqrt{t}} \exp\left(-\frac{t}{4a^2}\right) + \frac{\sqrt{\pi}}{2a} \operatorname{erf}\left(\frac{\sqrt{t}}{2a}\right) + \frac{\sqrt{\pi}}{2a}$$

is the general solution of the corresponding homogeneous integral equation.

The integral equation (16) is already solvable by the method of successive approximations and the solution to the corresponding homogeneous equation will be determined by the equality:

$$\psi_{2,0}(t) = c_0[\mathbf{K}]^{-1}[\psi_0(t)].$$

Similarly, as in the work [33], it is proven that function (6) is a solution to equation (1) and satisfies conditions (2), thus proving the validity of Theorem 1.

Conclusion

It is shown that in a non-cylindrical domain that degenerates at the initial moment of time into a point, the first boundary value problem for a fractional diffusion equation with the Riemann-Liouville fractional differentiation operator with respect to a time variable is singular, that is, it may not have a unique solution.

Acknowledgments

This research was funded by the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan. (Grant No. AP23488729).

Author Contributions

All authors contributed equally to this work.

Conflict of Interest

The authors declare no conflict of interest.

References

- 1 Нахушев А.М. Дробное исчисление и его применение / А.М. Нахушев. — М.: Физматлит, 2003. — 272 с.
- 2 Учайкин В.В. Метод дробных производных / В.В. Учайкин. — Ульяновск: Артишок, 2008. — 512 с.

- 3 Mainardi F. Fractional calculus and waves in linear viscoelasticity. An introduction to mathematical models / F. Mainardi. — London: Imperial College Press, 2010. — 347 p. <https://doi.org/10.1142/p614>
- 4 Тарасов В.Е. Модели теоретической физики с интегро-дифференцированием дробного порядка / В.Е. Тарасов. — М.–Ижевск: Ин-т комп. исслед., 2011. — 568 с.
- 5 Atanackovic T.M. Fractional calculus with applications in mechanics. Vibrations and diffusion processes / T.M. Atanackovic, S. Pilipovic, B. Stankovic, D. Zorica. — London: John Wiley and Sons, Inc., Hoboken, NJ, 2014. — 315 p. <https://doi.org/10.1002/9781118577530>
- 6 Wyss W. The fractional diffusion equation / W. Wyss // J. Math. Phys. — 1986. — 27. — No. 11. — P. 2782–2785. <https://doi.org/10.1063/1.527251>
- 7 Schneider W.R. Fractional diffusion and wave equations / W.R. Schneider, W. Wyss // J. Math. Phys. — 1989. — 30. — No. 1. — P. 134–144. <https://doi.org/10.1063/1.528578>
- 8 Kochubei A.N. Diffusion of fractional order / A.N. Kochubei // Differential Equations. — 1990. — 26. — No. 4. — P. 485–492.
- 9 Fujita Ya. Integrodifferential equation which interpolates the heat equation and the wave equation / Ya. Fujita // Osaka J. Math. — 1990. — 27. — No. 2. — P. 309–321. <https://doi.org/10.18910/8094>
- 10 Геккиева С.Х. Краевая задача для обобщенного уравнения переноса с дробной производной по времени / С.Х. Геккиева // Докл. Адыгской (Черкесской) междунар. акад. наук. — 1994. — 1. — № 1. — С. 17,18.
- 11 Mainardi F. The fundamental solutions for the fractional diffusion-wave equation / F. Mainardi // Appl. Math. Lett. — 1996. — 9. — № 6. — P. 23–28. [https://doi.org/10.1016/0893-9659\(96\)00089-4](https://doi.org/10.1016/0893-9659(96)00089-4)
- 12 Шогенов В.Х. Обобщенное уравнение переноса и дробные производные / В.Х. Шогенов, С.К. Кумыкова, М.Х. Шхануков-Лафипев // Докл. НАН Украины. — 1997. — № 12. — С. 47–54.
- 13 Buckwar E. Invariance of a partial differential equation of fractional order under the Lie group of scaling transformations / E. Buckwar, Yu. Luchko. // J. Math. Anal. Appl. — 1998. — 227. — No. 1. — P. 81–97. <https://doi.org/10.1006/jmaa.1998.6078>
- 14 Gorenflo R. Wright functions as scale-invariant solutions of the diffusion-wave equation / R. Gorenflo, Yu. Luchko, F. Mainardi // J. Comput. Appl. Math. — 2000. — 118. — No. 1-2. — P. 175–191. [https://doi.org/10.1016/S0377-0427\(00\)00288-0](https://doi.org/10.1016/S0377-0427(00)00288-0)
- 15 Gorenflo R. Mapping between solutions of fractional diffusion-wave equations / R. Gorenflo, A. Iskenderov, Yu. Luchko // Fract. Calc. Appl. Anal. — 2000. — 3. — No. 1. — P. 75–86.
- 16 Agrawal. O.P. Solution for a fractional diffusion-wave equation defined in a bounded domain / O.P. Agrawal // Nonlinear Dynam. — 2002. — 29. — P. 145–155. <https://doi.org/10.1023/A:1016539022492>
- 17 Псху А.В. Решение первой краевой задачи для уравнения диффузии дробного порядка / А.В. Псху // Диф. ур. — 2003. — 39. — № 9. — С. 1286–1289.
- 18 Псху А.В. Решение краевых задач для уравнения диффузии дробного порядка методом функции Грина / А.В. Псху // Диф. ур. — 39. — № 10. — С. 1430–1433. <https://doi.org/10.1023/B:DIEQ.0000012703.45373.a>
- 19 Eidelman S. D. Cauchy problem for fractional diffusion equations / S.D. Eidelman, A.N. Kochubei // J. Differential Equations. — 2004. — 199. — No. 2. — P. 211–255. <https://doi.org/10.1016/j.jde.2003.12.002>

- 20 Ворошилов А.А. Задача Коши для диффузионно-волнового уравнения с частной производной Капуто / А.А. Ворошилов, А.А. Килбас // Диф. ур. — 2006. — 42. — № 5. — С. 599–609. <https://doi.org/10.1134/S0012266106050041>
- 21 Yang. Y. Spectral collocation method for the time-fractional diffusion-wave equation and convergence analysis / Y. Yang, Y. Chen, Y. Huang, H. Wei // Computers and Mathematics with Applications. — 2017. — 73. — No. 6. — P. 1218–1232. <https://doi.org/10.1016/j.camwa.2016.08.017>
- 22 Псху А.В. Фундаментальное решение диффузионно-волнового уравнения дробного порядка / А.В. Псху // Изв. РАН. Сер. матем. — 2009. — 73. — № 2. — С. 141–182. <https://doi.org/10.4213/im2429>
- 23 Luchko Yu. Initial-boundary-value problems for the one-dimensional time-fractional diffusion equation // Fract. Calc. Appl. Anal. — 2012. — 15. — No. 1. — P. 141–160. <https://doi.org/10.2478/s13540-012-0010-7>
- 24 Мамчуев М.О. Необходимые нелокальные условия для диффузионно-волнового уравнения / М.О. Мамчуев // Вестн. Самар. гос. ун-та. Естественнонауч. сер. — 2014. — 118. — № 7. — С. 45–59. <https://doi.org/10.18287/2541-7525-2014-20-7-45-59>
- 25 Kilbas A.A. Theory and applications of fractional differential equations / A.A. Kilbas, H.M. Srivastava, J.J. Trujillo. — Elsevier, 2006. — 523 p. [https://doi.org/10.1016/s0304-0208\(06\)x8001-5](https://doi.org/10.1016/s0304-0208(06)x8001-5)
- 26 Псху А.В. Уравнения в частных производных дробного порядка / А.В. Псху. — М.: Наука, 2005. — 272 с.
- 27 Gevrey M. Sur les équations aux dérivées partielles du type parabolique / M. Gevrey // J. Math. Pures Appl. — 1913. — 6. — No. 9. — P. 305–476.
- 28 Mamchuev M.O. Fourier Problem for Fractional Diffusion–Wave Equation / M.O. Mamchuev, A.M. Mamchuev // Lobachevskii Journal of Mathematics. — 2023. — 44. — No. 2. — P. 620–628. <https://doi.org/10.1134/S1995080223020257>
- 29 Liu J. A new fractional derivative for solving time fractional diffusion wave equation / J. Liu, X. Yang, Y. Feng, L. Geng // Mathematical Methods in the Applied Sciences. — 2022. — 46. — No. 1. — P. 267–272. <https://doi.org/10.1002/mma.8509>
- 30 Kharin S. Mathematical model of electrical contact bouncing / S. Kharin // AIP Conference Proceedings. — 2015. — 1676. — No. 1. — 020019. <https://doi.org/10.1063/1.4930445>
- 31 Дженалиев М.Т. Нагруженные уравнения как возмущения дифференциальных уравнений / М.Т. Дженалиев, М.И. Рамазанов. — Алматы: Ғылым, 2010. — 334 с.
- 32 Pskhu A.V. Boundary value problem for fractional diffusion equation in a curvilinear angle domain / A.V. Pskhu, M.I. Ramazanov, N.K. Gulmanov, S.A. Iskakov // Bulletin of the Karaganda university. Mathematics series. — 2022. — № 1(105). — P. 83–95. <https://doi.org/10.31489/2022M1/83-95>
- 33 Псху А.В. Первая краевая задача для дробного диффузионно-волнового уравнения в нецилиндрической области / А.В. Псху // Изв. РАН. Сер. матем. — 2017. — 81. — № 6. — С. 158–179. <https://doi.org/10.4213/im8520>
- 34 Прудников А.П. Интегралы и ряды: справоч.: [В 3-х т.] — Т. 2. Специальные функции / А.П. Прудников, Ю.А. Брычков, О.И. Маричев. — 2-е изд., испр. — М.: Физматлит, 2003. — 664 с.
- 35 Jenaliyev M. On a Volterra equation of the second kind with “incompressible” kernel / M. Jenaliyev, M. Amangaliyeva, M. Kosmakova, M. Ramazanov // Advances in Difference Equations. — 2015. — 71. — P. 1–14. <https://doi.org/10.1186/s13662-015-0418-6>

Бұрыштық жойылмалы аядағы бөлшекті диффузия теңдеуі үшін бірінші шекаралық есеп

М.И. Рамазанов¹, А.В. Псху², М.Т. Омаров¹

¹Қолданбалы математика институты, Академик Е.А. Бөкетов атындағы Қарағанды университеті, Қарағанды, Қазақстан;

²РГА Кабардин-Балқар ғылыми орталығының Қолданбалы математика және автоматтандыру институты, Нальчик, Ресей

Мақала тармақталған фракталды құрылымдарда байқалатын мәселелерді қарастырады, мұнда уақыт бойынша бөлшектік туындылары бар диффузиялық теңдеулермен сипатталатын өте баяу транс-порттық процестер болуы мүмкін. Осы процестердің ерекше белгісі — олардың өте баяу релаксация жылдамдығы, мұнда физикалық шама оның бірінші туындысынан гөрі біртіндеп өзгереді. Мұндай құбылыстар кейде «қалдық жады» бар процестер ретінде жіктеледі. Зерттеуде уақыт бойынша Риман-Лиувилль бөлшектік дифференциалдау операторы бар бөлшектік диффузиялық теңдеу үшін бұрыштық облыста, бастапқы уақыт моментінде нүктеге дегенерацияланған бірінші шекаралық есептің шешімі ұсынылған. Онда зерттелетін есептің бар екендігі туралы теорема анықталған және есептің шешімі көрсетілген. Мақалада осындай өте баяу процестерді және олардың фракталды құрылымдарға әсерін түсінудің қажеттілігі атап өтілген. Жұмыс бөлшектік диффузиялық теңдеулердің кеңірек түсінілуіне осы процестердің қалай ықпал ететінін көрсетеді, теореманың бар екендігін дәлелдейді және есептің шешімін тұжырымдайды.

Кілт сөздер: дербес туынды теңдеу, бөлшек есептеу, бұрыштық облыс, ядро, әлсіз ерекшелік, параболикалық цилиндр, Карлеман-Векуа теңдеуі, жалпы шешім, жалғыз шешім, Риман-Лиувилльдің бөлшекті операторы.

Первая краевая задача для дробного диффузионного уравнения в угловой вырождающейся области

М.И. Рамазанов¹, А.В. Псху², М.Т. Омаров¹

¹Институт прикладной математики, Карагандинский университет имени академика Е.А. Букетова, Караганда, Казахстан;

²Институт прикладной математики и автоматизации, Кабардино-Балкарский научный центр РАН, Нальчик, Россия

Статья рассматривает проблемы, наблюдаемые в ветвящихся фрактальных структурах, где могут происходить сверхмедленные транспортные процессы; явление, описываемое диффузионными уравнениями с дробной производной по времени. Характерной особенностью этих процессов является их крайне медленная скорость релаксации, при которой физическая величина изменяется более постепенно, чем её первая производная. Такие явления иногда классифицируются как процессы с «остаточной памятью». В исследовании представлено решение первой краевой задачи в угловой области, вырождающейся в точку в начальный момент времени, для дробного диффузионного уравнения с оператором дробного дифференцирования Римана-Лиувилля по времени. В нем устанавливается теорема существования исследуемой задачи и строится представление решения. Авторами подчёркивается необходимость понимания этих сверхмедленных процессов и их влияния на фрактальные структуры. Работа демонстрирует, как эти процессы способствуют более широкому пониманию дробных диффузионных уравнений, доказывая существование теоремы и формулируя представление решения.

Ключевые слова: уравнение в частных производных, дробное исчисление, угловая область, ядро, слабая особенность, параболический цилиндр, уравнение Карлемана-Векуа, общее решение, единственное решение, дробный оператор Римана-Лиувилля.

References

- 1 Nakhushev, A.M. (2003). *Drobnoe ischislenie i ego primenenie [Fractional calculus and its application]*. Moscow: Fizmatlit [in Russian].
- 2 Uchaikin, V.V. (2008). *Metod drobnykh proizvodnykh [Fractional derivative method]*. Ulianovsk: Artishok [in Russian].
- 3 Mainardi, F. (2010). *Fractional calculus and waves in linear viscoelasticity. An introduction to mathematical models*. London. Imperial College Press. <https://doi.org/10.1142/p614>
- 4 Tarasov, V.E. (2011). *Modeli teoreticheskoi fiziki s integro-differentsirovaniem drobnogo poriadka [Models of theoretical physics with fractional order integro-differentiation]*. Institut kompiuternykh issledovaniy Moscow—Izhevsk: Institut kompiuternykh issledovaniy [in Russian].
- 5 Atanackovic, T.M., Pilipovic, S., Stankovic, B., & Zorica, D. (2014). *Fractional calculus with applications in mechanics. Vibrations and diffusion processes*. London. John Wiley and Sons, Inc., Hoboken, NJ. <https://doi.org/10.1002/9781118577530>
- 6 Wyss, W. (1986). The fractional diffusion equation. *J. Math. Phys.*, 27(11), 2782–2785. <https://doi.org/10.1063/1.527251>
- 7 Schneider, W.R., & Wyss, W. (1989). Fractional diffusion and wave equations. *J. Math. Phys.*, 30(1), 134–144. <https://doi.org/10.1063/1.528578>
- 8 Kochubei, A.N. (1990). Diffusion of fractional order. *Differential Equations*, 26(4), 485–492.
- 9 Fujita, Ya. (1990). Integrodifferential equation which interpolates the heat equation and the wave equation. *Osaka J. Math.*, 27(2), 309–321. <https://doi.org/10.18910/8094>
- 10 Gekkieva, S.Kh. (1994). Kraevaia zadacha dlia obobshchennogo uravneniia perenosa s drobnou proizvodnoi po vremeni [Boundary value problem for the generalized transport equation with fractional time derivative]. *Doklady Adygskoi (Cherkesskoi) mezhdunarodnoi akademii nauk — Reports of the Adyghe (Circassian) International Academy of Sciences*, 1(1), 17,18 [in Russian].
- 11 Mainardi, F. (1996). The fundamental solutions for the fractional diffusion-wave equation. *Appl. Math. Lett.*, 9(6), 23–28. [https://doi.org/10.1016/0893-9659\(96\)00089-4](https://doi.org/10.1016/0893-9659(96)00089-4)
- 12 Shogenov, V.Kh., Kumyкова, S.K., & Shkhanukov-Lafishev, M.Kh. (1997). Obobshchennoe uravnenie perenosa i drobnye proizvodnye [Generalized transport equation and fractional derivatives]. *Doklady Natsionalnoi akademii nauk Ukrainy — Reports of the NAS of Ukraine*, (12), 47–54 [in Russian].
- 13 Buckwar, E., & Luchko, Yu. (1998). Invariance of a partial differential equation of fractional order under the Lie group of scaling transformations. *J. Math. Anal. Appl.*, 227(1), 81–97. <https://doi.org/10.1006/jmaa.1998.6078>
- 14 Gorenflo, R., Luchko, Yu., & Mainardi, F. (2000). Wright functions as scale-invariant solutions of the diffusion-wave equation. *J. Comput. Appl. Math.*, 118(1-2), 175–191. [https://doi.org/10.1016/S0377-0427\(00\)00288-0](https://doi.org/10.1016/S0377-0427(00)00288-0)
- 15 Gorenflo, R., Iskenderov, A., & Luchko, Yu. (2000). Mapping between solutions of fractional diffusion-wave equations. *Fract. Calc. Appl. Anal.*, 3(1), 75–86.
- 16 Agrawal, O.P. (2002). Solution for a fractional diffusion-wave equation defined in a bounded domain. *Nonlinear Dynam.*, 29(1), 145–155. <https://doi.org/10.1023/A:1016539022492>
- 17 Pskhu, A.V. (2003). Reshenie pervoi kraevoi zadachi dlia uravneniia diffuzii drobnogo poriadka [Solution of the first boundary value problem for the fractional order diffusion equation]. *Differentsialnye uravneniia — Differential equations*, 39(9), 1286–1289 [in Russian].

- 18 Pskhu, A.V. (2003). Reshenie kraevykh zadach dlia uravneniia diffuzii drobnogo poriadka metodom funktsii Grina [Solving boundary value problems for the fractional order diffusion equation using the Green's function method]. *Differentsialnye uravneniia – Differential equations*, 39(10), 1430–1433 [in Russian]. <https://doi.org/10.1023/B:DIEQ.0000012703.45373.aa>
- 19 Eidelman, S.D., & Kochubei, A.N. (2004). Cauchy problem for fractional diffusion equations. *J. Differential Equations*, 199(2), 211–255. <https://doi.org/10.1016/j.jde.2003.12.002>
- 20 Voroshilov, A.A., & Kilbas, A.A. (2006). Zadacha Koshi dlia diffuzionno-volnovogo uravneniia s chastnoi proizvodnoi Kaputo [Cauchy problem for the diffusion-wave equation with Caputo's partial derivative]. *Differentsialnye uravneniia – Differential equations*, 42(5), 599–609 [in Russian]. <https://doi.org/10.1134/S0012266106050041>
- 21 Yang, Y., Chen, Y., Huang, Y., & Wei, H. (2017). Spectral collocation method for the time-fractional diffusion-wave equation and convergence analysis. *Computers and Mathematics with Applications*, 73(6), 1218–1232. <https://doi.org/10.1016/j.camwa.2016.08.017>
- 22 Pskhu, A.V. (2009). Fundamentalnoe reshenie diffuzionno-volnovogo uravneniia drobnogo poriadka. [Fundamental solution of the diffusion-wave equation of fractional order] *Izvestiia RAN. Serii matematicheskaiia – News of the Russian Academy of Sciences. Mathematical series*, 73(2), 141–182 [in Russian]. <https://doi.org/10.4213/im2429>
- 23 Luchko, Yu. (2012). Initial-boundary-value problems for the one-dimensional time-fractional diffusion equation. *Fract. Calc. Appl. Anal.*, 15(1), 141–160. <https://doi.org/10.2478/s13540-012-0010-7>
- 24 Mamchuev, M.O. (2014). Neobkhodimye nelokalnye usloviia dlia diffuzionno-volnovogo uravneniia [Necessary nonlocal conditions for the diffusion-wave equation]. *Vestnik Samarskogo gosudarstvennogo universiteta. Estestvennonauchnaia serii – Vestnik of Samara University. Natural Science Series*, 118(7), 45–59 [in Russian]. <https://doi.org/10.18287/2541-7525-2014-20-7-45-59>
- 25 Kilbas, A.A., Srivastava, H.M., & Trujillo, J.J. (2006). *Theory and applications of fractional differential equations*. Elsevier. [https://doi.org/10.1016/s0304-0208\(06\)x8001-5](https://doi.org/10.1016/s0304-0208(06)x8001-5)
- 26 Pskhu, A.V. (2005). *Uravneniia v chastnykh proizvodnykh drobnogo poriadka [Fractional partial differential equations]*. Moscow: Nauka [in Russian].
- 27 Gevrey, M. (1913). Sur les équations aux dérivées partielles du type parabolique [On partial differential equations of the parabolic type]. *J. Math. Pures Appl.*, 6(9), 305–476 [in French].
- 28 Mamchuev, M.O., & Mamchuev, A.M. (2023). Fourier Problem for Fractional Diffusion–Wave Equation. *Lobachevskii Journal of Mathematics*, 44(2), 620–628. <https://doi.org/10.1134/S1995080223020257>
- 29 Liu, J., Yang, X., Feng, Y., & Geng, L. (2022). A new fractional derivative for solving time fractional diffusion wave equation. *Mathematical Methods in the Applied Sciences*, 46(1), 267–272. <https://doi.org/10.1002/mma.8509>
- 30 Kharin, S. (2015). Mathematical model of electrical contact bouncing. *AIP Conference Proceedings*, 1676(1). <https://doi.org/10.1063/1.4930445>
- 31 Dzenaliev, M.T., & Ramazanov, M.I. (2010). *Nagruzhennye uravneniia kak vozmushcheniia differentsialnykh uravnenii [Loaded equations as perturbations of differential equations]*. Almaty: Gylym [in Russian].
- 32 Pskhu, A.V., Ramazanov, M.I., Gulmanov, N.K., & Iskakov, S.A. (2022). Boundary value problem for fractional diffusion equation in a curvilinear angle domain. *Bulletin of the Karaganda university. Mathematics series*, 1(105), 83–95. <https://doi.org/10.31489/2022M1/83-95>

- 33 Pskhu, A.V. (2017). Pervaia kraevaia zadacha dlia drobnogo diffuzionno-volnovogo uravneniia v netsilindricheskoi oblasti [The first boundary value problem for the fractional diffusion-wave equation in a non-cylindrical domain.]. *Izvestiia RAN. Seriya matematicheskaia — News of the Russian Academy of Sciences. Mathematical series*, 81 (6), 158–179 [in Russian]. <https://doi.org/10.4213/im8520>
- 34 Prudnikov, A.P., Brychkov, Yu.A., & Marichev, O.I. (2003). *Integraly i riady: spravochnik: [v 3-kh tomakh]. Tom 2. Spetsialnye funktsii [Integrals and series: handbook: in volume 2. Special functions.]* 2-e izdanie, ispravlennoe. Moscow: Fizmatlit [in Russian].
- 35 Jenaliyev, M., Amangaliyeva, M., Kosmakova, M., & Ramazanov, M. (2015). On a Volterra equation of the second kind with “incompressible” kernel. *Advances in Difference Equations*, 71, 1–14. <https://doi.org/10.1186/s13662-015-0418-6>.

*Author Information**

Murat Ibraevich Ramazanov — Doctor of physical and mathematical sciences, Professor, Institute of Applied Mathematics, Karaganda Buketov University, 28 Universitetskaya street, Karaganda, 100028, Kazakhstan; e-mail: ramamur@mail.ru; <https://orcid.org/0000-0002-2297-5488>.

Arsen Vladimirovich Pskhu — Doctor of physical and mathematical sciences, Docent, A. D. of IAMA KBSC RAS, Head of the Department of Fractional Calculus, The Institute of Applied Mathematics and Automation of Kabardin-Balkar Scientific Center of RAS, st. Shortanova, 89 A, Nalchik, 360000, Kabardin-Balkar Republic, Russia; e-mail: pskhu@list.ru; <https://orcid.org/0000-0002-0506-3516>.

Madi Tulegenovich Omarov (*corresponding author*) — Master of natural sciences, teacher, Institute of Applied Mathematics, Karaganda Buketov University, 28 Universitetskaya street, Karaganda, 100028, Kazakhstan; e-mail: madiomarovt@gmail.com; <https://orcid.org/0000-0002-9026-5912>.

*The author’s name is presented in the order: First, Middle and Last Names.