

Lower and Upper Bounds for the Norm of Multipliers of Multiple Trigonometric Fourier Series in Lebesgue Spaces

E. D. Nursultanov and N. T. Tleukhanova

UDC 517.51

Let $1 < p \leq q < \infty$, and let \mathbb{T}^n be the n -dimensional torus. We say that a sequence $\lambda = \{\lambda_k\}_{k \in \mathbb{Z}^n}$ of complex numbers is a *multiplier of trigonometric Fourier series* from $L_p(\mathbb{T}^n)$ into $L_q(\mathbb{T}^n)$ if for an arbitrary function $f \in L_p(\mathbb{T}^n)$ with Fourier series $\sum_{k \in \mathbb{Z}^n} \hat{f}(k)e^{ikx}$ there exists a function $f_\lambda \in L_q(\mathbb{T}^n)$ with Fourier series $\sum_{k \in \mathbb{Z}^n} \lambda_k \hat{f}(k)e^{ikx}$ and if $T_\lambda, T_\lambda f = f_\lambda$, is a bounded operator from $L_p(\mathbb{T}^n)$ into $L_q(\mathbb{T}^n)$. The set \mathfrak{m}_p^q of these multipliers is a normed linear space with the norm

$$\|\lambda\|_{\mathfrak{m}_p^q} = \|T_\lambda\|_{L_p \rightarrow L_q}.$$

In this paper, we find lower and upper bounds for the norm of multipliers $\lambda \in \mathfrak{m}_p^q$. The methods used in our proofs are based on the results in [1, 2] establishing new inequalities describing the relationship between metric properties of a function and the summability of its Fourier coefficients with respect to the trigonometric system.

The following result by Hörmander is well known [3]: if $1 < p \leq 2 \leq q < \infty$ and $1/r = 1/p - 1/q$, then the embedding $l_{r\infty}(\mathbb{Z}^n) \hookrightarrow \mathfrak{m}_p^q$ holds, that is,

$$\|\lambda\|_{\mathfrak{m}_p^q} \leq c \sup_{e \in E} \frac{1}{|e|^{1-1/r}} \left| \sum_{m \in e} \lambda_m \right|.$$

Here $l_{r\infty}(\mathbb{Z}^n)$ is a discrete Lorentz space, E is the set of all finite subsets in \mathbb{Z}^n , and $|e|$ is the cardinality of a set e .

The assertion below is a refinement of Hörmander's theorem for multiple series ($n \geq 2$).

Theorem 1. *Suppose that $1 < p \leq 2 \leq q < \infty$, $p' = p/(p-1)$, and (j_1, \dots, j_n) is an arbitrary permutation of the sequence $1, \dots, n$. Then the inequality*

$$\|\lambda\|_{\mathfrak{m}_p^q} \leq C \sup_{k \in \mathbb{N}^n} \frac{1}{(\prod_{i=1}^n k_i)^{1/q+1/p'}} \sum_{m_1=1}^{k_1} \dots \sum_{m_n=1}^{k_n} \lambda_{m_1 \dots m_n}^{*j_1 \dots *j_n}$$

holds, where the sequence $\{\lambda_{m_1 \dots m_n}^{*j_1 \dots *j_n}\}_{m \in \mathbb{N}^n}$ is obtained from $\{|\lambda_r|\}_{r \in \mathbb{Z}^n}$ by passing consecutively to the non-decreasing rearrangement with respect to the indices r_{j_1}, \dots, r_{j_n} and C is a constant depending only on the parameters p and q and the dimension n .

Let $B = \{m_0, m_0 + 1, \dots, m_0 + l\}$ be an interval in \mathbb{Z} , and let $d \in \mathbb{N}$, $d > l$. A set of the form $I = \bigcup_{k=0}^N [B + kd] = \bigcup_{k=0}^N \{m + kd : m \in B\}$ will be called a *harmonic interval* in \mathbb{Z} . Accordingly, a set of the form $I = I_1 \times \dots \times I_n$, where the I_i are harmonic intervals in \mathbb{Z} , will be called a harmonic interval in \mathbb{Z}^n .

Theorem 2. *Let $1 < p \leq 2 \leq q < \infty$, let $p' = p/(p-1)$, and let M_0 be the set of all harmonic intervals in \mathbb{Z}^n . If $\lambda = \{\lambda_k\}_{k \in \mathbb{Z}^n} \in \mathfrak{m}_p^q$, then*

$$\sup_{Q \in M_0} \frac{1}{|Q|^{1/q+1/p'}} \left| \sum_{m \in Q} \lambda_m \right| \leq C \|\lambda\|_{\mathfrak{m}_p^q}.$$

Definition. We say that a sequence $\{a_k\}_{k \in \mathbb{Z}^n}$ of complex numbers is *monotone in the extended sense* if there exists a number $c > 0$ such that

$$|a_k| \leq \frac{c}{|Q_k|} \left| \sum_{r \in Q_k} a_r \right|$$

for every $k \in \mathbb{Z}^n$, where $Q_k = \{r \in \mathbb{Z}^n : 0 \leq |r_j| \leq |k_j|, j = 1, \dots, n\}$.

If a sequence $\lambda = \{\lambda_k\}_{k \in \mathbb{N}^n}$ is monotone or quasimonotone with respect to each k_j , then it is monotone in the extended sense.

Corollary 1. Let $1 < p \leq 2 \leq q < \infty$ and $p' = p/(p-1)$. Suppose that $\lambda = \{\lambda_k\}_{k \in \mathbb{Z}^n}$ is a monotone sequence in the extended sense. Then a necessary and sufficient condition for the relation $\lambda \in \mathbf{m}_p^q$ to hold is

$$F(\lambda) = \sup_{m \in \mathbb{N}^n} \frac{1}{(m_1 \cdots m_n)^{1/q+1/p'}} \left| \sum_{k_1=-m_1}^{m_1} \cdots \sum_{k_n=-m_n}^{m_n} \lambda_{k_1 \dots k_n} \right| < \infty.$$

In this case, $\|\lambda\|_{\mathbf{m}_p^q} \sim F(\lambda)$.

Consider the Shapiro sequence $\{\varepsilon_k\}_{k=1}^\infty$ (see [4]) defined by the recursion formulas $\varepsilon_1 = 1$, $\varepsilon_{2k} = \varepsilon_k$, $\varepsilon_{2k+1} = (-1)^k \varepsilon_k$, $k \in \mathbb{N}$. It consists of the numbers ± 1 and satisfies the estimate

$$\left| \sum_{m=0}^{N-1} \varepsilon_m e^{imx} \right| \leq cN^{1/2}, \quad x \in [0, 2\pi).$$

Let $\varepsilon = \{\varepsilon_k\}_{k \in \mathbb{N}^n}$, where $\varepsilon_k = \varepsilon_{k_1} \cdots \varepsilon_{k_n}$.

Theorem 3. Suppose that either $1 < p \leq q \leq 2$ or $2 \leq p \leq q < \infty$, $\varepsilon = \{\varepsilon_k\}_{k \in \mathbb{N}^n}$ is the Shapiro sequence, (j_1, \dots, j_n) is an arbitrary permutation of the sequence $1, \dots, n$, and

$$\frac{1}{r} = \begin{cases} 1/2 + 1/q & \text{for } 2 \leq p \leq q < \infty, \\ 1/p' + 1/2 & \text{for } 1 < p \leq q \leq 2. \end{cases}$$

Then

$$C^{-1}(r, n) \sup_{\substack{r \in \mathbb{Z}^n \\ m \in \mathbb{N}^n}} \frac{1}{(m_1 \cdots m_n)^{1/r}} \left| \sum_{1 \leq k \leq m} \varepsilon_k \lambda_{k+r} \right| \leq \|\lambda\|_{\mathbf{m}_p^q} \leq C(r, n) \sup_{m \in \mathbb{N}^n} \frac{1}{(m_1 \cdots m_n)^{1/r}} \sum_{1 \leq k \leq m} \lambda_{k_1 \dots k_n}^{*j_1 \dots *j_n}. \quad (1)$$

The upper and lower bounds in Theorem 3 do not depend on the parameter p for $2 \leq p \leq q < \infty$ and on the parameter q for $1 < p \leq q \leq 2$. This means that if these parameters are varied, then the class \mathbf{m}_p^q varies with preservation of relation (1).

This assertion solves the problem of finding sufficient conditions for λ to belong to the space of multipliers $\mathbf{m}_p = \mathbf{m}_p^p$ essentially depending on the parameter p . In this connection, we note that, according to [5], the classical theorems on multipliers are somewhat unnatural, since the sufficient conditions in them usually do not depend on p .

Corollary 2. Let $\varepsilon = \{\varepsilon_k\}_{k \in \mathbb{N}^n}$ be the Shapiro sequence. Suppose that either $1 < p \leq q \leq 2$ or $2 \leq p \leq q < \infty$. If the sequence $\{\varepsilon_k \lambda_k\}_{k \in \mathbb{N}^n}$ is monotone in the extended sense, then the relation $\{\lambda_k\}_{k \in \mathbb{N}^n} \in \mathbf{m}_p^q$ is equivalent to the inequality

$$\sup_{m \in \mathbb{N}^n} \frac{1}{(m_1 \cdots m_n)^{1/r}} \sum_{k_1=1}^{m_1} \cdots \sum_{k_n=1}^{m_n} |\lambda_{k_1 \dots k_n}| < \infty,$$

where the number $1/r$ is defined in Theorem 3.

There are as "many" sequences satisfying the assumptions of Corollary 2 as sequences monotone in the extended sense. For example, if $\lambda = \{\lambda_k\}$ is monotone in the extended sense, then $\lambda_\varepsilon = \{\varepsilon_k \lambda_k\}$ satisfies the assumptions of Corollary 2, since $\{\varepsilon_k^2 \lambda_k\}_{k \in \mathbb{N}^n} = \{\lambda_k\}_{k \in \mathbb{N}^n}$.

References

1. E. D. Nursultanov, *Mat. Sb.*, **189**, No. 3, 83–102 (1998).
2. E. D. Nursultanov, *East J. App.*, No. 3, 243–275 (1998).
3. L. Hörmander, *Acta Math.*, **104**, 93–140 (1960).
4. J. Brillhart and L. Carlitz, *Proc. Amer. Math. Soc.*, **25**, 114–118 (1970).
5. E. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton Mathematical Series, No. 30, Princeton University Press, Princeton, 1970.

Translated by V. E. Nazaikinskii

Репозиторий КАРГУ