

K.S. Panchenko^{1*}, M.M. Silantieva¹, D.V. Sokolova²

¹Altai State University, Barnaul, Russia;

²Federal Research Center "All-Russian Institute of Plant Genetic Resources named after N.I. Vavilov", St. Petersburg, Russia

*Corresponding author: kseniya.potapova.00@yandex.ru

Resource potential of amaranth and possibilities of its cultivation in the conditions of the south of Western Siberia

Amaranth, with more than 60 species, was introduced to Europe in the 16th century and is currently gaining popularity due to its high content of protein, vitamins and antioxidants. Its importance has increased in recent decades, both globally and in the conditions of Siberia, where it has become an object of selective breeding. The article provides a brief historical sketch on the study and breeding of new amaranth varieties in the south of Western Siberia. In addition, the biological features of amaranth (*Amaranthus* L.) and its importance as a valuable food and fodder crop are considered. The analysis of breeding traits allowed us to identify the most promising samples for successful cultivation in the south of Western Siberia: amaranth panicle (PC-318 and PC-391) and dark amaranth (PC-42). They are characterized by the shortest growing season and have greater resistance to adverse factors. The results of the study show that amaranth can become an important crop for Western Siberia, providing high yield and adaptability in difficult climatic conditions.

Keywords: squalene, growing season length, *Amaranthus cruentus* L., *Amaranthus hypochondriacus* L., *Amaranthus caudatus* L., *Amaranthus tricolor* L., introducer, ontogenesis stages.

Introduction

Amaranth (*Amaranthus* L.) is a genus of plants in the Amaranth family (Amaranthaceae Juss.) with more than 60 species. The greatest diversity of forms, species and varieties is represented on the territory of South America and Mexico, which is its homeland. Northern India and China are considered to be the second center of form formation. Amaranth was introduced to Europe only in the 16th century by the Spaniards for ornamental purposes and only two centuries later it was cultivated for seeds and animal feed.

Amaranth is of great importance in the food industry and fodder production. Plants contain a large amount of protein, balanced and unique in amino acid composition, valuable oils, pectin. Daily consumption of 100 g of amaranth seeds provides the human body with essential amino acids for 140 % of the adequate daily intake. It is important to note the presence of antioxidants in the seeds and green mass: carotenoids, amaranthine, squalene and vitamins B₂, C. The ascorbic acid content in different varieties of amaranth can be up to 50 mg/100 g [1–4]. In fodder production, amaranth is valued for its high potassium content and a large amount of easily digestible plant protein (1.0-1.5 tons/ha) [5]. The most valuable species in economic terms are: amaranth panicle (*Amaranthus cruentus* L.), amaranth sad (*A. hypochondriacus* L.), amaranth caudate (*A. caudatus* L.), and amaranth tricolor (*A. tricolor* L.).

In Russia, fodder species of amaranth became known in the 30s of the XX century, when Nikolai Ivanovich Vavilov brought seeds from an expedition to South America. He actively advocated for the wider use of amaranth in agriculture because of the large number of advantages of this crop. Even in difficult climatic conditions, high yields were noted because the C-4 type of photosynthesis provides amaranth with rapid growth and drought resistance [6, 7]. The resumption of amaranth research as a valuable food, fodder and ornamental crop occurred only at the end of the XX century and is currently being actively pursued [8]. Due to the difficult weather and climatic conditions of Siberia, the assortment of fodder crops in the region was rather scarce. To solve this problem, comprehensive research on the evaluation of non-traditional crops and selection of promising introducers was launched. Since the mid-1980s, active study of agrobiological features of the "forgotten" high-protein crop in Siberia began using seed samples transferred from VIR (Federal Research Center All-Russian Institute of Plant Genetic Resources named after N.I. Vavilov). Thus, at the workshop "Results of research and applied work with amaranth for 1987-1988", the staff of Altai State University Marina Mikhailovna Yablokova, Raisa Nikolaevna Afonina and Elena Vasilievna Repetunova presented a report on the successful experience of growing amaranth in the conditions of the Altai Territory [9].

In the conditions of the Tomsk and Novosibirsk regions, scientists under the leadership of Valentina Pavlovna Rykova also identified promising early maturing samples of the collection, which in the zone of risky agriculture give high yields [10, 11].

A little later, in 1995, the Federal Research Center IRCIG SB RAS together with the Omsk Agrarian Scientific Center registered the first in Siberia broad-plastic and early maturing amaranth variety “Cherginsky” [12].

Since the early 2000s, Svetlana Ivanovna Mikhailova, Tatiana Petrovna Astafurova, and Anastasia Anatolievna Burenina have been actively studying amaranth biology at Tomsk State University [13–17]. In 2006, the State Register of Breeding Achievements Allowed for Use in the Russian Federation included the variety “Yantar” developed by the Federal Altai Scientific Center for Agrobiotechnology and FIC ICIG SB RAS. The variety has a short vegetation period (106 days) and has a special value in fodder production. It significantly surpasses oilseed radish, rape, peas and beans in terms of yield of fodder mass and quality of plant raw material [18, 19].

In 2019, scientists of the Kuzbass State Agricultural Academy showed interest in grain forms of amaranth and optimized the technology of their cultivation in the conditions of the West Siberian forest-steppe in the Kemerovo region. In particular, the dependence of the quality of green mass and grain yield on the biological characteristics of the variety and such an agrotechnical technique as row spacing width was determined [20]. Currently, the staff of the Siberian Botanical Garden of TSU is implementing a project on cultivation of agricultural plants on carbonaceous farms in the sub-taiga zone of Western Siberia and, in addition to the range of traditional crops, offers amaranth as a promising introducer [21]. Thus, the interest in amaranth as a highly productive agricultural crop with a valuable protein composition has been growing rapidly in recent years.

The purpose of this work was the primary evaluation of a number of breeding and significant traits of plants of the amaranth genus to identify the most promising species and varieties for cultivation in the south of Western Siberia.

Experimental

The objects of the study were 7 amaranth cultivars of different origins: 1) *Amaranthus cruentus* L. — PK-318 (Russia), PK-391 (Brazil); 2) Amaranth hybrid (*A. hybridus* L. convar *erythrostr.*) — PK-96 (Germany); 3) Amaranth caudatus (*A. caudatus* L.) — PK-146 (Germany), PK-150 (Greece); 4) Amaranth *tricolor* (*A. tricolor* L.) — PK-168 (Nepal); 5) Dark amaranth (*A. hypochondriacus* L.) — PK-429 (Mexico).

Seed material was obtained from the world collection of plant genetic resources of All-Russian Institute of Plant Industry named after N. Vavilov, St. Petersburg. To perform the experimental part of the work we used “Methodology of tests for distinctiveness, uniformity and stability of amaranth Federal State Center “State sort committee” (2007).

Pre-sowing tillage was carried out in spring using a gasoline cultivator Huter GMC-6.5. Sowing was carried out on June 3. Seed sowing rate was 0.5 kg/ha, sowing density was 20 plants/m². Repetition of the experiment was fourfold, experimental plot area was 1 m². Sowing was carried out manually in a wide-row method. Seeds were mixed with washed coarse sand. Later, we additionally thinned the seedlings and regularly loosened the soil during the growing season to break the soil crust and kill weeds. Starting from the emergence of seedlings, phenological phases were noted and morphological parameters were measured. Seeds were harvested and threshed manually.

Results and Discussion

In the vegetation period of amaranth there are 5 phenological phases: sprouting, growth, bottoming, flowering and fruiting. On average, the totality of all development periods is 110–140 days and varies depending on the variety and species. Seeds begin to germinate when the soil warms up to +10...+12°C, but the seedlings are extremely unstable to any frost. Therefore, sowing amaranth in the south of Western Siberia should be carried out when the threat of return frosts has finally passed — in the last decade of May or early June. Sprouts appear on the 6th-7th day and their development in the first three weeks is extremely slow due to a small supply of nutrients because of the size of the seeds. However, due to a special type of photosynthesis, the plants begin to grow rapidly after the start of the growth phase and the formation of green phytomass. The budding phase lasts 1-2 weeks on average, energy is redistributed to the formation of inflorescences — panicles, and growth in height is somewhat slower. Then, flower blossoming and seed ripening take place [22].

Amaranth plants have a fairly high seed production coefficient, but not every variety has seeds that reach physiological maturity before the onset of frost and cold weather. Therefore, breeding selection of this crop is aimed at developing early-ripening, cold- and drought-resistant varieties with high yield values. Among the species and varieties studied, Russian (PK-318, “Frant”) and Brazilian (PK-391) amaranth varieties, as well as dark amaranth from Mexico (PK-429) showed excellent performance. The vegetation period for representatives of these species was 95–100 days, with the first seeds beginning to form on the 60–65th day (Fig.). Hybrid amaranth (PK-96, Germany) should be considered later maturing. In this species, the budding phase occurred on 93–95 days from seedlings, and the first seeds on single specimens matured on 105–110 days. *Amaranthus caudatus* (PK-146, Germany) had a similar growing season, but was twice affected by harmful insects and mold fungi. Tricolor amaranth (PC-168, Nepal) and Greek tail amaranth (PC-150) did not have time to form seeds before the first frosts and finished the vegetation at the budding and flowering stage. These species have too long vegetation period and their cultivation in local conditions is inexpedient.



Figure. Amaranth panicle (PK-318, Russia) during growth and flowering phases

In addition to the growing season, an important economic and valuable feature of amaranth is the low stature of plants. Private farms do not have special harvesting machines and use combines like KSK-100, KPKU-75, etc. For morphological description of the studied species we used the materials of “Methods of testing for distinctiveness, homogeneity and stability of amaranth” of Federal state center “State sort committee” (2007). The obtained data are presented in Table.

Morphological traits of the studied amaranth breeding samples

PC-	Species name	Average plant height in the middle of vegetation, cm	Average plant height at the end of vegetation, cm	Presence of side shoots	Stem thickness at the end of vegetation, cm	Coloration of leaf lamina, presence of "spot"	Inflorescence coloration
318	<i>Amaranthus cruentus</i> L.	110	190	-	1,9	red, no.	red
96	<i>Amaranthus hybridus</i> L.convar <i>erythrost.</i>	136	223	-	2,2	green, V-shaped "spot" in the center	red
146	<i>Amaranthus caudatus</i> L.	126	190	-	1,9	green, no	green
150	<i>Amaranthus caudatus</i> L.	106,5	160	+	2,5	green, no	green
168	<i>Amaranthus tricolor</i> L.	57,5	100	-	1,2	green, egg-shaped "spot" in the center	red
391	<i>Amaranthus cruentus</i> L.	71	119	-	1,2	green, no	green
429	<i>Amaranthus hypochondriacus</i> L.	76,5	146	-	1	green, no	green with red spots
Statistical characteristics*							
	\bar{x}	97.6	161.1				
	σ	29.7	43.3				
	Cv	30.4	26.9				
	SDx	11.2	16.4				
Note. * \bar{x} — mean, σ — standard deviation, CV — coefficient of variation, %, SDx — standard error of the experiment							

According to the results of the experiment of the first year, hybrid amaranth plants (PK-96, Germany) were the tallest — up to 223 cm. The average height of amaranth panicle PK-318 (Russia) at the time of harvesting was 190 cm, dark amaranth PK-429 (Brazil) — 146 cm. The lowest of the listed varieties was the height of Mexican amaranth panicle PC-391 (Mexico) — 119 cm.

All the studied samples with a vegetation period suitable for local conditions can be proposed for cultivation not only as food and fodder plants, but also for ornamental purposes. In addition, the leaves of the red-colored amaranth variety (PK-318, Russia) can be used to produce fermented tea, which is of great importance in functional nutrition [23].

Conclusion

Thus, at the primary stage of study of a number of samples from the amaranth collection of "All-Russian Institute of Plant Industry named after N. Vavilov" the most promising species, varieties and breeding samples for cultivation in climatic conditions of the south of Western Siberia were identified: amaranth panicle (PK-318, "Frant", Russia; PK-391, Brazil), dark amaranth (PK-429, Mexico).

Acknowledgements

The authors would like to thank the reviewers for their contribution to the peer review of this article.

References

- 1 Гинс М.С. Биологически активные вещества амаранта. Амарантин: свойства, механизмы действия и практическое использование / М.С. Гинс. — М.: Изд-во РУДН, 2002. — 182 с.
- 2 Леушкина Е.В. Растения семейства амарантовых — перспективные источники водорастворимых витаминов / Е.В. Леушкина, Л.В. Донченко // Сб. Всерос. науч. конф. «Теория и практика современной аграрной науки». — Новосибирск: Золотой колос, 2018. — 371 с.

- 3 Клешнева Е.В. Растения амарантовых — эффективный источник биологически ценного белка в обогащенных продуктах питания / Е.В. Клешнева, Л.В. Донченко, Т.В. Щеколдина // Современные аспекты производства и переработки сельскохозяйственной продукции. — Краснодар: КубГАУ, 2017. — 764 с.
- 4 Moazma S. An overview of the nutritional and therapeutic properties of amaranth / S. Moazma, S. Farhan, A. Muhammad, R. Amara, A. Aliha, S. Sadia, H. Muzzamal, A. Hafiz, A.R. Muhammad, M. Haroon, Al Jb. Entessar // International Journal of Food Properties, 2024. — No 27(1). — P. 263–272. DOI: 10.1080/10942912.2024.2304266.
- 5 Белоножкина Т.Г. Амарант — культура больших возможностей для ЦЧЗ России / Т.Г. Белоножкина // Актуальные проблемы инноваций с нетрадиционными природными ресурсами и создания функциональных продуктов: Материалы II Росс. науч.-практ. конф. — М., 2003. — С. 33–35.
- 6 Вавилов Н.И. Проблема новых культур / Н.И. Вавилов // Избранные труды. — Т. 5. — М., 1965. — С. 537–563.
- 7 Магомедов И.М. К вопросу об истории интродукции амаранта в СССР и РФ / И.М. Магомедов // Процветание науки. — 2022. — № 9 (15).
- 8 Магомедов И.М. Амарант — прошлое, настоящее и будущее / И.М. Магомедов, Т.В. Чиркова // Успехи современного естествознания. — 2015. — № 1. — С. 1108–1113.
- 9 Яблокова М.М. Опыт возделывания амаранта в условиях Алтайского края / М.М. Яблокова, Р.Н. Афонина, Е.В. Репетунова // Итоги научно-исследовательских и прикладных работ с культурой амаранта за 1987-1988 гг.: тез. докл. Раб. совещ. — Л., 1989. — С. 40-41.
- 10 Рыкова В.П. Новые кормовые растения — резерв кормопроизводства / В.П. Рыкова // Сб докл. «Рациональное использование природных ресурсов Сибири». — Томск: Изд-во Том. ун-та, 1989. — С. 178.
- 11 Рыкова В.П. Багряная нива амаранта / В.П. Рыкова // Красное знамя. — Томск, 1990. — С. 2.
- 12 Железнов В.А. Сравнительный анализ самоопыленных линий амаранта при его интродукции в Западной Сибири / А.В. Железнов, Н.Б. Железнова, Н.В. Бурмакина, Л.П. Солоненко // Сиб. вестн. сельскохоз. науки. — 2008. — № 7(187). — С. 10–16.
- 13 Михайлова С.И. Особенности роста и размножения амаранта на юге Томской области / С.И. Михайлова, А.А. Буренина, Т.П. Астафурова // Роль ботанических садов в сохранении биоразнообразия. — Ростов на/Д., 2002. — С. 214–217.
- 14 Михайлова С.И. Морфофизиологические особенности видов р. *Amaranthus* L., интродуцированных в Томской области / С.И. Михайлова, А.А. Буренина, Т.П. Астафурова // Новые и нетрадиционные растения и перспективы их использования: материалы V Междунар. симпоз. — М., 2003. — Т. 1. — С. 64–66.
- 15 Михайлова С.И. Интродукционные исследования амаранта в Томском госуниверситете / С.И. Михайлова, Т.П. Астафурова, А.А. Буренина // Современные проблемы и достижения аграрной науки в животноводстве, растениеводстве и экономике: сб. тр. Регион. науч.-практ. конф. — Томск, 2012. — Вып. 14. — С. 87–92.
- 16 Астафурова Т.П. Амарант — перспективная кормовая культура для возделывания в подтаежной зоне Западной Сибири / Т.П. Астафурова, С.И. Михайлова, А.А. Буренина // Сиб. вестн. Сельскохоз. науки. — 2004. — № 1 (151). — С. 56–60.
- 17 Буренина А.А. Перспективные сорта амаранта зернового и овощного направления в условиях Томской области / А.А. Буренина, Н.В. Сотникова, С.И. Михайлова, Т.П. Астафурова // Сб. тр. Регион. науч.-практ. конф. «Современные проблемы и достижения аграрной науки в животноводстве, растениеводстве и экономике». — Томск, 2006. — Вып. 9. — С. 17–20.
- 18 Шукис Е.Р. Совершенствование видового и сортового состава зернобобовых и кормовых культур в условиях Алтайского края / Е.Р. Шукис, С.К. Шукис // Достижения науки и техники АПК. — 2008. — № 11. — С. 38–40.
- 19 Шукис Е.Р. Особенности работы с кормовыми культурами / Е.Р. Шукис, С.К. Шукис // Актуальные вопросы АПК Сибири: итоги и перспективы: сб. науч. работ (к 65-летию Алтайского НИИ сельского хозяйства). — Барнаул: Федерал. гос. бюджет. науч. учрежд. «Алтайский научно-исследовательский институт сельского хозяйства», 2015. — С. 201–219.
- 20 Агронет — отраслевая сеть инноваций в АПК. — 2019. — [Электронный ресурс]. — Режим доступа: <https://apknnet.ru>
- 21 Новости Томского государственного университета. — 2022. — [Электронный ресурс]. — Режим доступа: <https://news.tsu.ru>
- 22 Романов Д.В. Способы и нормы посева амаранта метельчатого на тёмно-серых лесных почвах Курской области: автореф. дис. ... канд. сельскохоз. наук / Д.В. Романов. — Курск, 2000. — 18 с.
- 23 Соколова Д.В. Патент № 2793627 С1 Российская Федерация, МПК А23F 3/00. Способ производства ферментированного чая из красноокрашенных сортов амаранта. № 2022103757, опубл. 04.04.2023.

К.С. Панченко, М.М. Силантьева, Д.В. Соколова
**Амаранттың ресурстық әлеуеті және оны
Батыс Сібірдің оңтүстігінде өсіру мүмкіндігі**

Алпыстан астам түрі бар амарант Еуропаға XVI ғасырда енгізілген және қазіргі уақытта ақуыздың, витаминдердің және антиоксиданттардың көп болуына байланысты танымал болып келеді. Оның маңызы соңғы онжылдықтарда әлемдік масштабта да, Сібірде де өсті, онда ол селекциялық іріктеу объектісіне айналды. Мақалада Батыс Сібірдің оңтүстігінде амаранттың жаңа сорттарын зерттеу және өсіру туралы қысқаша тарихи очерк берілген. Сонымен қатар, амаранттың биологиялық ерекшеліктері (*Amaranthus* L.) және оның құнды азық-түлік және жем-шөп дақылдары ретіндегі маңызы қарастырылған. Селекциялық белгілерді талдау Батыс Сібірдің оңтүстігінде табысты өсіру үшін ең перспективалы үлгілерді анықтауға мүмкіндік берді: шашақты амарант (ПК-318 және ПК-391) және күнгірт амарант (ПК-42). Олар ең қысқа вегетациялық кезеңмен сипатталады және қолайсыз факторларға үлкен төзімділікке ие. Зерттеу нәтижелері амаранттың Батыс Сібір үшін маңызды ауыл шаруашылығы дақылына айналуы мүмкін екенін көрсетті, ол жоғары өнімділік пен күрделі климаттық жағдайларда бейімделуді қамтамасыз етеді.

Кілт сөздер: сквален, вегетациялық кезеңнің ұзақтығы, *Amaranthus cruentus* L., *Amaranthus hypochondriacus* L., *Amaranthus caudatus* L., *Amaranthus tricolor* L., жерсіндірілген түр, онтогенез кезеңдері.

К.С. Панченко, М.М. Силантьева, Д.В. Соколова

**Ресурсный потенциал амаранта и возможности его выращивания
в условиях юга Западной Сибири**

Амарант, насчитывающий более 60 видов, завезен в Европу в XVI веке и в настоящее время набирает популярность благодаря высокому содержанию белка, витаминов и антиоксидантов. Его значение возросло в последние десятилетия, как в мировом масштабе, так и в условиях Сибири, где он стал объектом селекционного отбора. В статье приведен краткий исторический очерк по изучению и выведению новых сортов амаранта на юге Западной Сибири. Кроме того, рассмотрены биологические особенности амаранта (*Amaranthus* L.) и его значимость как ценной пищевой и кормовой культуры. Анализ селекционных признаков позволил выделить наиболее перспективные образцы для успешного возделывания в условиях юга Западной Сибири: амарант метельчатый (ПК-318 и ПК-391) и амарант темный (ПК-42). Они характеризуются наиболее коротким вегетационным периодом и обладают большей устойчивостью к неблагоприятным факторам. Результаты исследования показывают, что амарант может стать важной сельскохозяйственной культурой для Западной Сибири, обеспечивая высокую урожайность и адаптивность в сложных климатических условиях.

Ключевые слова: сквален, длина вегетационного периода, *Amaranthus cruentus* L., *Amaranthus hypochondriacus* L., *Amaranthus caudatus* L., *Amaranthus tricolor* L., интродуцент, стадии онтогенеза.

References

- 1 Gins, M.S. (2002). *Biologicheski aktivnye veshchestva amaranta. Amarantin: svoystva, mekhanizmy deystviia i prakticheskoe ispolzovanie* [Biologically active substances of amaranth. Amaranth: properties, mechanisms of action and practical use]. Moscow [in Russian].
- 2 Leushkina, E.V., & Donchenko, L.V. (2018). Rasteniia semeystva amarantovykh — perspektivnye istochniki vodorastvorimykh vitaminov [Plants of the amaranth family are promising sources of water-soluble vitamins]. *Sbornik Vserossiiskoi nauchnoi konferentsii «Teoriia i praktika sovremennoi agrarnoi nauki» — Book of abstracts of All-Russian scientific conference “Theory and practice of modern agrarian science”*. Novosibirsk: Zolotoi kolos [in Russian].
- 3 Kleshneva, E.V., Donchenko, L.V., & Shchekoldina, T.V. (2017). Rasteniia amarantovykh — effektivnyi istochnik biologicheskoi tseennogo belka v obogashchennykh produktakh pitaniia [Amaranth plants are an effective source of biologically valuable protein in fortified foods]. *Sovremennye aspekty proizvodstva i pererabotki sel'skokhoziaistvennoi produktsii — The Modern Aspects of Production and Processing of Agricultural Products*. Krasnodar: KubGAU [in Russian].
- 4 Moazma, S., Farhan, S., Muhammad, A., Amara, R., Aliha, A., Sadia, S., Muzzamal, H., Hafiz, A., Muhammad, A.R., Haroon, M., & Entessar, Al.Jb. (2024). An overview of the nutritional and therapeutic properties of amaranth. *International Journal of Food Properties*, 27 (1), 263–272. DOI: 10.1080/10942912.2024.2304266.
- 5 Belonozhkina, T.G. (2003). Amaranth — kultura bolshikh vozmozhnostei dlia TsChZ Rossii [Amaranth is a culture of great opportunities for the Central Economic District of Russia]. *Aktualnye problemy innovatsii s netraditsionnymi prirodnymi resursami i*

sozdaniia funktsionalnykh produktov: materialy II Rossiiskoi nauchno-prakticheskoi konferentsii — Actual problems of innovations with non-traditional natural resources and creation of functional products: materials of the II Russian Scientific and Practical Conference. Moscow, 33–35 [in Russian].

6 Vavilov, N.I. (1965). Problema novykh kultur [The problem of new cultures]. *Izbrannye trudy — Selected works*. Moscow [in Russian].

7 Magomedov, I.M. (2022). K voprosu ob istorii introduktsii amaranta v SSSR i RF [On the history of the introduction of amaranth in the USSR and the Russian Federation]. *Prosvetanie nauki — Prosperity of science*, 9 (15) [in Russian].

8 Magomedov, I.M. (2015). Amarant — proshloe, nastoiashchee i budushchee [Amaranth — past, present and future]. *Uspekhi sovremennogo estestvoznaniia — Advances in modern natural science*, 1, 1108–1113 [in Russian].

9 Yablokova, M.M., Afonina, R.N., & Repetunova, E.V. (1989). Opyt vozdel'yvaniia amaranta v usloviakh Altaiskogo kraia [Experience of amaranth cultivation in the Altai Territory]. *Itogi nauchno-issledovatel'skikh i prikladnykh rabot s kulturoi amaranta za 1987-1988 gg.»: tezisy dokladov Rabochego soveshchaniia — Results of research and applied works with amaranth culture for 1987-1988: abstracts of reports of the workshop*. Leningrad, 40-41 [in Russian].

10 Rykova, V.P. (1989). Novye kormovye rasteniia — rezerv kormoproizvodstva [New forage plants are a reserve of forage production]. *Sbornik dokladov Ratsionalnoe ispolzovanie prirodnykh resursov Sibiri — Book of abstracts: Rational use of natural resources of Siberia*. Tomsk: Izdatel'stvo Tomskogo universiteta [in Russian].

11 Rykova, V.P. (1990). Bagrianaia niva amaranta [The scarlet field of amaranth]. *Krasnoe znamia — Red Banner*. Tomsk [in Russian].

12 Zheleznov, V.A. (2008). Sravnitelnyi analiz samoopylennykh linii amaranta pri ego introduktsii v Zapadnoi Sibiri [Comparative analysis of self-pollinated amaranth lines during its introduction in Western Siberia]. *Sibirskii vestnik sel'skokhoziaistvennoi nauki — Siberian Bulletin of agricultural science*, 7(187), 10–16 [in Russian].

13 Mikhailova, S.I. (2002). Osobennosti rosta i razmnozheniia amaranta na yuge Tomskoi oblasti [Features of amaranth growth and reproduction in the south of the Tomsk region]. *Rol botanicheskikh sadov v sokhraneniі bioraznoobraziia — Role of botanical gardens in biodiversity conservation. Rostov-na-Donu* [in Russian].

14 Mikhailova, S.I. (2003). Morfofiziologicheskie osobennosti vidov r. *Amaranthus* L., introdutsirovannykh v Tomskoi oblasti [Morphophysiological features of R. *amaranthus* L. species introduced in the Tomsk region]. *Novye i netraditsionnye rasteniia i perspektivy ikh ispolzovaniia: materialy V Mezhdunarodnogo simpoziuma — New and non-traditional plants and prospects for their use: proceedings of the V International Symposium*, 1, 64–66 [in Russian].

15 Mikhailova, S.I. (2012). Introduktsionnye issledovaniia amaranta v Tomskom gosuniversitete [Introduction studies of amaranth at Tomsk State University]. *Sovremennye problemy i dostizheniia agrarnoi nauki v zhivotnovodstve, rasteniievodstve i ekonomike: sbornik trudov Regionalnoi nauchno-prakticheskoi konferentsii — New and non-traditional plants and prospects of their utilization: proceedings of the V International Symposium*. Moscow, 14, 87–92 [in Russian].

16 Astafurova, T.P. (2004). Amarant — perspektivnaia kormovaia kultura dlia vozdel'yvaniia v podtaezhnoi zone Zapadnoi Sibiri [Amaranth is a promising forage crop for cultivation in the subtaiga zone of Western Siberia]. *Sibirskii vestnik sel'skokhoziaistvennoi nauki — Siberian Bulletin of agricultural science*, 1(151), 56–60 [in Russian].

17 Burenina, A.A. (2006). Perspektivnye sorta amaranta zernovogo i ovoshchnogo napravleniia v usloviakh Tomskoi oblasti [Promising varieties of amaranth of grain and vegetable direction in the conditions of the Tomsk region]. *Sbornik trudov Regionalnoi nauchno-prakticheskoi konferentsii «Sovremennye problemy i dostizheniia agrarnoi nauki v zhivotnovodstve, rasteniievodstve i ekonomike» — Proceedings of the regional scientific-practical conference “Modern problems and achievements of agrarian science in animal husbandry, crop production and economics”*. Tomsk, 9, 17–20 [in Russian].

18 Shukis, E.R. (2008). Sovershenstvovanie vidovogo i sortovogo sostava zernobobovykh i kormovykh kultur v usloviakh Altaiskogo kraia [Improvement of the species and varietal composition of leguminous and forage crops in the conditions of the Altai Territory]. *Dostizheniia nauki i tekhniki Agropromyshlennogo kompleksa — Achievements of science and techniques of agroindustrial complex*, 11, 38–40 [in Russian].

19 Shukis, E.R. (2015). Osobennosti raboty s kormovymi kulturami [Features of working with forage crops]. *Aktualnye voprosy Agropromyshlennogo kompleksa Sibiri: itogi i perspektivy: Sbornik nauchnykh rabot — Actual issues of agro-industrial complex of Siberia: results and prospects: Collection of scientific works*. Barnaul, 201–209 [in Russian].

20 (2019). Agronet — otraslevaia set innovatsii v Agropromyshlennom komplekse [Agronet — sectoral network of innovations in agribusiness]. Retrieved from <https://apknet.ru> [in Russian].

21 (2022). Novosti Tomskogo gosudarstvennogo universiteta [News of Tomsk state university]. Retrieved from <https://news.tsu.ru> [in Russian].

22 Romanov, D.V. (2000). Sposoby i normy poseva amaranta metelchatogo na temno-serykh lesnykh pochvakh Kurskoi oblasti [Methods and patterns of development of the paniculate amaranth on dark gray forest soils of the Kursk region]. *Extended abstract of candidate's thesis*. Kursk [in Russian].

23 Sokolova, D.V. (2023). Patent No. 2793627 C1 Rossiiskaia Federatsiia, MPK A23F 3/00. Sposob proizvodstva fermentirovannogo chaia iz krasnookrashennykh sortov amaranta [Patent No. 2793627 C1 Russian Federation. Method of production of fermented tea from red-colored amaranth varieties]: No 2022103757, publ. 04.04.2023 [in Russian].

Information about the authors

Panchenko Ksenia Sergeevna — PhD-student, Lecturer, Department of Botany, Institute of Biology and Biotechnology, Altai State University, Barnaul, Russian; e-mail: *kseniya.potapova.00@yandex.ru*;

Silantieva Marina Mikhailovna — Doctor of biological science, Professor, Head of Botany Department, Institute of Biology and Biotechnology, Altai State University, Barnaul, Russia; e-mail: *msilan@mail.ru*;

Sokolova Diana Viktorovna — PhD., Senior researcher, Curator of beet, spinach and amaranth collections, Department of Genetic Resources of Vegetable and Melon Crops, Federal Research Center “All-Russian Institute of Plant Genetic Resources named after N.I. Vavilov”, St. Petersburg, Russia; e-mail: *dianasokol@bk.ru*.

Buketov University