

Numerical solution of source identification multi-point problem of parabolic partial differential equation with Neumann type boundary condition

C. Ashyralyev^{1,2,3,*}, T.A. Ashyralyeva⁴

¹Bahcesehir University, Istanbul, Turkey;

²Khoja Akhmet Yassawi International Kazakh-Turkish University, Turkestan, Kazakhstan;

³National University of Uzbekistan named after Mirzo Ulugbek, Tashkent, Uzbekistan;

⁴Yagshygeldi Kakayev International University of Oil and Gas, Ashgabat, Turkmenistan
(E-mail: charyar@gmail.com, tazegulashyralyewa@gmail.com)

We study a source identification boundary value problem for a parabolic partial differential equation with multi-point Neumann type boundary condition. Stability estimates for the solution of the overdetermined mixed BVP for multi-dimensional parabolic equation were established. The first and second order of accuracy difference schemes for the approximate solution of this problem were proposed. Stability estimates for both difference schemes were obtained. The result of numerical illustration in test example was given.

Keywords: inverse problem, source identification, parabolic equation, difference scheme, stability, nonlocal condition, boundary value problem, well-posedness, stability estimates, mixed problem.

2020 Mathematics Subject Classification: 34B10, 35K10, 49K40.

Introduction

Various techniques can be used to solve source identification problems (SIPs) for parabolic equations. These may include optimization algorithms, regularization methods, or numerical techniques such as finite element and finite difference methods (see [1–28] a references therein).

In papers [3, 18], SIP for abstract differential equation with self-adjoint positive definite operator A

$$\frac{dv(t)}{dt} + Av(t) = p + f(t), \quad 0 < t < 1, \quad (1)$$

$$v(0) = \varphi, \quad v(1) = \psi \quad (2)$$

in a Hilbert space H was investigated. In paper [3], for solution of SIP (1), (2), stability estimates in the Hölder norms were obtained.

Some applications to boundary value problems (BVPs) for partial differential equation (PDE) and approximate solutions were studied in [8, 12].

Let $s_1, \mu_1, s_2, \mu_2, \dots, s_r, \mu_r$ be given numbers so that

$$\sum_{k=1}^r |\mu_k| < 1, \quad 0 \leq s_1 < s_2 < \dots < s_r < 1 \quad (3)$$

*Corresponding author. E-mail: charyar@gmail.com

This research was funded by the grant no. AP19676663 of the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan.

Received: 18 December 2023; Accepted: 20 May 2024.

© 2024 The Authors. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

and elements $\varphi, \psi \in H$ and function $f : [0, 1] \rightarrow H$ are given. In paper [15] SIP to find a pair (p, v) for equation

$$\frac{dv(t)}{dt} + Av(t) = p + f(t), 0 < t < 1,$$

with multi-point nonlocal conditions

$$v(0) = \sum_{k=1}^r \mu_k v(s_k) + \psi, \tag{4}$$

$$v(1) = \varphi \tag{5}$$

was studied and stability estimates for the solution were given in the following theorem.

Theorem 1. [15] Assume that conditions (3) for interior points and coefficients are valid, $\varphi \in H$, $\psi \in D(A)$, $f \in C^\alpha(H)$ ($\alpha \in (0, 1)$) are given. Then, for the solution $(v(t), p)$ of SIP (1), (4), (5) the stability estimates

$$\|v\|_{C(H)} \leq M \left[\|\varphi\|_H + \|\psi\|_H + \|f\|_{C(H)} \right]$$

and

$$\|p\|_H \leq M \left[\|A\varphi\|_H + \|A\psi\|_H + \frac{1}{\alpha} \|f\|_{C^\alpha(H)} \right]$$

are fulfilled, where $M \in R^+$ does not depend on f, ψ, φ and α . Here $C(H)$, $C^\alpha(H)$ and $C_1^\alpha(H)$ are the Banach spaces of H -valued functions $u(t)$ with the corresponding norms

$$\begin{aligned} \|u\|_{C(H)} &= \max_{0 \leq t \leq 1} \|u(t)\|_H, \|u\|_{C^\alpha(H)} = \|u\|_{C(H)} + \sup_{0 \leq t < t+\tau \leq 1} \frac{\|u(t+\tau)\|_H - \|u(t)\|_H}{\tau^\alpha}, \\ \|u\|_{C_1^\alpha(H)} &= \|u\|_{C(H)} + \sup_{0 \leq t < t+\tau \leq 1} \frac{(1-t)^\alpha \|u(t+\tau)\|_H - \|u(t)\|_H}{\tau^\alpha}, \end{aligned} \tag{6}$$

respectively.

1 SI parabolic problem with multi-point boundary conditions

Now, we study a source identification (SI) BVP for the multi-dimensional PDE.

Let $\Omega = (0, 1)^n \subset \mathbb{R}^n$ with boundary $S = \partial\Omega$, $\bar{\Omega} = \Omega \cup S$.

Denote by $L_2(\Omega)$ and $W_2^2(\Omega)$ the Hilbert spaces of integrable functions $u(y)$, defined on Ω , equipped with the corresponding norms

$$\begin{aligned} \|u\|_{L_2(\Omega)} &= \left\{ \int_{y \in \Omega} |u(y)|^2 dy_1 \dots dy_n \right\}^{\frac{1}{2}}, \\ \|u\|_{W_2^2(\Omega)} &= \left\{ \int_{y \in \Omega} \left(|u(y)|^2 + \sum_{i=1}^n \sum_{j=1}^n |u_{y_i y_j}(y)|^2 \right) dy_1 \dots dy_n \right\}^{\frac{1}{2}}. \end{aligned}$$

Let $\varphi \in L_2(\Omega)$, $\psi \in W_2^2(\Omega)$, $f \in C^\alpha(L_2(\Omega))$ be given functions, and $a_i : \Omega \rightarrow R^+$ be known smooth function for any index $i = 1, \dots, n$.

In $[0, 1] \times \Omega$, we study multi-dimensional SIP for parabolic PDE with multi-point boundary and nonlocal conditions

$$\begin{cases} v_t(t, x) - \sum_{i=1}^n (a_i(x) v_{x_i}(t, x))_{x_i} + \sigma v(t, x) = f(t, x) + p(x), \\ x = (x_1, \dots, x_n) \in \Omega, 0 < t < 1, \\ \frac{\partial}{\partial n} v(t, x) = 0, x \in S, 0 < t < 1, \\ v(0, x) = \sum_{k=1}^r \mu_k v(s_k, x) + \psi(x), v(1, x) = \varphi(x), x \in \bar{\Omega}, \end{cases} \tag{7}$$

where \vec{n} is the normal vector to Ω at corresponding boundary point.

The differential expression

$$A^x u(x) = - \sum_{i=1}^n (a_i(x) u_{x_i}(x))_{x_i} + \sigma u(x)$$

defines SAPD operator A^x , which acts on the Hilbert space $L_2(\Omega)$ with the domain

$$D(A^x) = \left\{ u \mid u \in W_2^2(\Omega), \frac{\partial u}{\partial \vec{n}}(x) = 0 \text{ on } S \right\}.$$

So, the SI BVP (7) for the multi-dimensional parabolic PDE can be replaced with the abstract problem (1), (4), (5) for $H = L_2(\Omega)$. By using stability estimates of Theorem 1, we obtain the following stability estimates for solution of BVP (7).

Theorem 2. Suppose that conditions (3) are satisfied, $\varphi, \psi \in W_2^2(\Omega)$ and $f \in C^\alpha(L_2(\Omega))$. Then, for the solution of multi-dimensional SIP for parabolic PDE (7), the following estimates are valid

$$\begin{aligned} \|p\|_{L_2(\Omega)} &\leq M \left[\|\varphi\|_{W_2^2(\Omega)} + \|\psi\|_{W_2^2(\Omega)} + \frac{1}{\alpha} \|f\|_{C^\alpha(L_2(\Omega))} \right], \\ \|v\|_{C(L_2(\Omega))} &\leq M \left[\|\varphi\|_{L_2(\Omega)} + \|\psi\|_{L_2(\Omega)} + \|f\|_{C(L_2(\Omega))} \right], \end{aligned}$$

where positive number M is independent of f, ψ, φ and α .

2 First and second order of ADSs

We will use the set of uniform grid points

$$[0, 1]_\tau = \{t_k = k\tau, k = 0, 1, \dots, N, N\tau = 1\}.$$

To discretize problem (7) we use algorithm with two steps. Firstly, we define grid spaces

$$\begin{aligned} \tilde{\Omega}_h &= \{x = x_m = (h_1 m_1, \dots, h_n m_n); m = (m_1, \dots, m_n), \\ & m_j = 0, \dots, N_j, h_j N_j = 1, j = 1, \dots, n\}, \\ \Omega_h &= \tilde{\Omega}_h \cap \Omega, S_h = \tilde{\Omega}_h \cap S. \end{aligned}$$

Introduce difference operator A_h^x by formula

$$A_h^x v^h(x) = - \sum_{i=1}^n \left(a_i(x) v_{\bar{x}_i}^h(x) \right)_{x_i, j_i} + \sigma v^h(x),$$

which acts in space of grid functions $v^h(x)$ and satisfies the condition $Dv^h(x) = 0$ for all $x \in S_h$.

Applying A_h^x , we arrive at the multi-point nonlocal BVP for some infinite system of ordinary differential equations. Secondly, by using Equation (26) [15; p.1922], we get the first order of accuracy difference scheme (ADS)

$$\begin{cases} \tau^{-1} \left(v_k^h(x) - v_{k-1}^h(x) \right) + A_h^x v_k^h(x) = f^h(t_k, x) + p^h(x), & 1 \leq k \leq N, x \in \tilde{\Omega}_h, \\ v_N^h(x) = \varphi^h(x), v_0^h(x) = \sum_{i=1}^r \mu_i v_{l_i}^h(x) + \psi^h(x), & x \in \tilde{\Omega}_h. \end{cases} \quad (8)$$

By using Equations (37)–(39) [15; p. 1925], we get the second order of ADS

$$\begin{cases} \tau^{-1} \left(v_k^h(x) - v_{k-1}^h(x) \right) + A_h^x \left(I + \frac{\tau A_h^x}{2} \right) v_k^h(x) \\ = \left(I + \frac{\tau A_h^x}{2} \right) \left(f^h(t_{k-\frac{\tau}{2}}, x) + p^h(x) \right), \quad 1 \leq k \leq N, \quad x \in \tilde{\Omega}_h, \\ v_N^h(x) = \varphi^h(x), \quad x \in \tilde{\Omega}_h, \\ v_0^h(x) = \sum_{i=1}^r \left\{ \mu_i (1 - \rho_i) v_{i_i}^h(x) + \mu_i \rho_i v_{i_i+1}^h(x) \right\} + \psi^h(x), \quad x \in \tilde{\Omega}_h. \end{cases} \quad (9)$$

Denote by $L_{2h} = L_2(\tilde{\Omega}_h)$ and $W_{2h}^2 = W_2^2(\tilde{\Omega}_h)$, the spaces of the grid functions $u^h(x) = \{u(h_1 m_1, \dots, h_n m_n)\}$ defined on $\tilde{\Omega}_h$, equipped with the corresponding norms

$$\begin{aligned} \|u^h\|_{L_{2h}} &= \left(\sum_{x \in \tilde{\Omega}_h} |u^h(x)|^2 h_1 \cdots h_n \right)^{1/2}, \\ \|u^h\|_{W_{2h}^2} &= \|u^h\|_{L_{2h}} + \left(\sum_{x \in \tilde{\Omega}_h} \sum_{r=1}^n |(u^h(x))_{x_r \bar{x}_r, m_r}|^2 h_1 \cdots h_n \right)^{1/2}, \end{aligned}$$

and by $\mathcal{C}_\tau(L_{2h}) = \mathcal{C}([0, 1]_\tau, L_{2h})$, the Banach space of L_{2h} -valued grid functions $u^\tau = \{u_k\}_1^N$ with the suitable norm $\|u^\tau\|_{\mathcal{C}_\tau(L_{2h})} = \max_{1 \leq k \leq N} \|u_k\|_{L_{2h}}$.

Let $\mathcal{C}^\alpha(L_{2h}) = \mathcal{C}^\alpha([0, 1]_\tau, L_{2h})$ and $\mathcal{C}_\tau^\alpha(L_{2h}) = \mathcal{C}_\tau^\alpha([0, 1]_\tau, L_{2h})$ be correspondingly Hölder and weighted Hölder spaces with the corresponding norms defined by (6) for $H = L_{2h}$.

Theorem 3. Suppose that τ and $|h| = \sqrt{h_1^2 + \cdots + h_n^2}$ are sufficiently small positive numbers, $\varphi^h \in L_{2h}$, $\psi^h \in W_{2h}^2$ and $\{f_k^h\}_1^N \in \mathcal{C}_\tau^\alpha(L_{2h})$. Then, for the solution of difference schemes (DSs) (8) and (9), the following stability estimates hold

$$\begin{aligned} \|p^h\|_{\mathcal{C}_\tau(L_{2h})} &\leq M \left[\|\varphi^h\|_{L_{2h}} + \|\psi^h\|_{W_{2h}^2} + \frac{1}{\alpha} \left\| \{f_k^h\}_1^N \right\|_{\mathcal{C}_\tau^\alpha(L_{2h})} \right], \\ \left\| \{v_k^h\}_1^N \right\|_{\mathcal{C}_\tau(L_{2h})} &\leq M \left[\|\varphi^h\|_{L_{2h}} + \|\psi^h\|_{L_{2h}} + \left\| \{f_k^h\}_1^N \right\|_{\mathcal{C}_\tau(L_{2h})} \right], \end{aligned}$$

where M is independent of $\{f_k^h\}_1^N$, $\psi^h(x)$, $\varphi^h(x)$ and τ .

The proof of Theorem 3 based on Theorems 3.1 and 3.2 of paper [15] on stability estimate for solutions of corresponding DSs for approximate solution of abstract SIP (1), (4), (5) and the theorem on the coercivity inequality for the solution of the elliptic difference problem in L_{2h} .

3 Numerical analysis

For test example, we consider the SIP

$$\begin{cases} v_t(t, x) - (3 + 2 \cos x)v_{xx}(t, x) + 2 \sin x \cdot v_x(t, x) + v(t, x) = f(t, x) + p(x), \\ 0 < x < \pi, \quad 0 < t < 1, \\ v(1, x) = \varphi(x), \\ v(0, x) = v(\frac{1}{3}, x) + \psi(x), \quad 0 \leq x \leq \pi, \\ v_x(t, 0) = 0, \quad v_x(t, \pi) = 0, \quad 0 \leq t \leq 1 \end{cases} \quad (10)$$

for one-dimensional parabolic PDE. Here

$$f(t, x) = (e^{-t} - e^{-1})(3 \cos x + 2 \cos 2x) - e^{-1} \cos x, \quad 0 < x < \pi, \quad 0 < t < 1,$$

$$\varphi(x) = \cos x, \quad \psi(x) = \left(1 - e^{-\frac{1}{3}}\right) \cos x, \quad 0 \leq x \leq \pi.$$

It is easy to check that the pair $(e^{-1}(4 \cos x + 2 \cos 2x), e^{-t} \cos x)$ is the exact solution of problem (10).

An algorithm of finding the solution of problem (10) contains three stages. In the first stage, we find the solution of auxiliary BVP

$$\begin{cases} u_t(t, x) - (3 + 2 \cos x)u_{xx}(t, x) + 2 \sin x \cdot u_x(t, x) + u(t, x) \\ = (3 + 2 \cos x) \cos x - 2 \sin x \cdot \sin x + \cos x + f(t, x), \quad 0 < x < \pi, \quad 0 < t < 1, \\ u(1, x) - u(\frac{1}{3}, x) = \psi(x), \quad 0 \leq x \leq \pi, \\ u_x(t, 0) = 0, \quad u_x(t, \pi) = 0, \quad 0 \leq t \leq 1. \end{cases} \quad (11)$$

Then, in the second stage, we find $p(x)$ by

$$p(x) = -(3 + 2 \cos x)u_{xx}(1, x) + 2 \sin x \cdot u_x(1, x) + u(1, x).$$

In the third stage, we put $p(x)$ in the right side of equation (10) and solve that problem for $v(t, x)$.

Introduce the set of grid points

$$[0, 1]_\tau \times [0, \pi]_h = \{(t_k, x_n) \mid t_k = k\tau, \quad k = 1, \dots, N-1, \quad N\tau = 1, \\ x_n = nh, \quad n = 1, \dots, M-1, \quad Mh = \pi\}.$$

We use notation $l = [\frac{\gamma}{\tau}]$ for greatest integer function of $\frac{\gamma}{\tau}$ and $\rho = \frac{\gamma}{\tau} - l$.

So, we get the first order of ADS for SIP (10)

$$\begin{cases} \frac{v_n^k - v_n^{k-1}}{\tau} - \frac{(3+2 \cos x_n)(v_{n+1}^k - 2v_n^k + v_{n-1}^k)}{h^2} + \frac{\sin(x_n)(v_{n+1}^k - v_{n-1}^k)}{h} + v_n^k \\ = f(t_k, x_n) + p(x_n), \quad k = 1, \dots, N, \quad n = 1, \dots, M-1, \\ v_n^N = \varphi_n, \quad v_n^0 - \rho v_n^1 = \psi_n, \quad n = 0, \dots, M, \\ v_0^k = v_1^k, \quad v_M^k = v_{M-1}^k, \quad k = 0, \dots, N. \end{cases}$$

Later, $p(x_n)$ can be obtained by

$$p(x_n) = -\frac{(3 + 2 \cos(x_n))(u_{n+1}^N - 2u_n^N + u_{n-1}^N)}{h^2} + \frac{\sin(x_n)(u_{n+1}^N - u_{n-1}^N)}{h} + u_n^N, \quad (12)$$

where $\{u_n^k\}$ is solution of the difference problem

$$\begin{cases} \frac{u_n^k - u_n^{k-1}}{\tau} - \frac{(3 + 2 \cos(x_n))(u_{n+1}^k - 2u_n^k + u_{n-1}^k)}{h^2} \\ + \frac{\sin(x_n)(u_{n+1}^k - u_{n-1}^k)}{h} + u_n^k = f(t_k, x_n) - \frac{(3 + 2 \cos(x_n))(\varphi_{n+1} - 2\varphi_n + \varphi_{n-1})}{h^2} \\ + \frac{\sin(x_n)(\varphi_{n+1} - \varphi_{n-1})}{h} + \varphi_n, \quad k = 1, \dots, N, \quad n = 1, \dots, M-1, \\ u_n^0 - u_n^l = \psi_n, \quad n = 0, \dots, M, \\ u_0^k - u_M^k = 0, \quad u_M^k - u_{M-1}^k = 0, \quad k = 0, \dots, N, \end{cases} \quad (13)$$

which is the first order of ADS for approximate solution of the nonlocal BVP (11).

For computational reasons it is convenient to write (13) in the following matrix form

$$A_n u_{n+1} + B_n u_n + C_n u_{n-1} = I\theta_n, \quad n = 1, \dots, M-1, \\ u_0 = u_1, \quad u_M = u_{M-1}. \quad (14)$$

Here, θ_n is column vector, A_n, B_n, C_n, I are square matrices with $(N + 1)$ rows and columns:

$$A_n = \begin{bmatrix} 0 & \dots & 0 & 0 \\ & & & 0 \\ a_n R & & & \vdots \\ & & & 0 \end{bmatrix}, \quad C_n = \begin{bmatrix} 0 & \dots & 0 & 0 \\ & & & 0 \\ c_n R & & & \vdots \\ & & & 0 \end{bmatrix},$$

$$B_n = \begin{bmatrix} 1 & 0 & 0 & 0 & \dots & -1 & \dots & 0 & 0 & 0 & 0 \\ b_n & d & 0 & 0 & \dots & 0 & \dots & 0 & 0 & 0 & 0 \\ 0 & b_n & d & 0 & \dots & 0 & \dots & 0 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \dots & \dots & \dots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & 0 & \dots & 0 & b_n & d & 0 \\ 0 & 0 & 0 & 0 & \dots & 0 & \dots & 0 & 0 & b_n & d \end{bmatrix},$$

$$a_n = -(3 + 2 \cos(x_n))h^{-2} + \sin(x_n)h^{-1}, d = \frac{1}{\tau},$$

$$b_n = 1 + d + 2(3 + 2 \cos(x_n))h^{-2},$$

$$c_n = -(3 + 2 \cos(x_n))h^{-2} - \sin(x_n)h^{-1},$$

$$\theta_n = \begin{bmatrix} \theta_n^0 \\ \vdots \\ \theta_n^N \end{bmatrix}, \quad u_{n\pm 1} = \begin{bmatrix} u_{n\pm 1}^0 \\ \vdots \\ u_{n\pm 1}^N \end{bmatrix}_{(N+1) \times 1}, \quad u_n = \begin{bmatrix} u_n^0 \\ \vdots \\ u_n^N \end{bmatrix}_{(N+1) \times 1}.$$

R is the $N \times N$ identity matrix, as well as

$$\theta_n^0 = \psi_n, \quad n = 1, \dots, M - 1,$$

$$\theta_n^k = f(t_k, x_n) - \frac{(3+2 \cos(x_n))(\varphi_{n+1}-2\varphi_n+\varphi)}{h^2} + \frac{\sin(x_n)(\varphi_{n+1}-\varphi_{n-1})}{h} + \varphi_n,$$

$$k = 1, \dots, N, \quad n = 1, \dots, M - 1.$$

We search solution of (14) by recurrence formula

$$u_n = \alpha_{n+1}u_{n+1} + \beta_{n+1}, \quad n = M - 1, \dots, 1,$$

where α_n and β_n ($n = 1, \dots, M - 1$) are column vectors with $(N + 1)$ elements. For the solution of difference equation (14) we use the following formulas for α_n, β_n

$$\alpha_n = -(B_n + C_n \alpha_{n-1})^{-1} A_n,$$

$$\beta_n = (B_n + C_n \alpha_{n-1})^{-1} (R \theta_n - C_n \beta_{n-1}), \quad n = 1, \dots, M - 1,$$

where α_1 is the $(N + 1) \times (N + 1)$ identity matrix and β_1 is the column vector with $(N + 1)$ zeros. u_M is computed by formula

$$u_M = (A_M + B_M + C_M \alpha_M)^{-1} (I \theta_M - C_M \beta_M).$$

Second, applying appropriate approximation formulas for derivatives in the nonlocal BVP (10), we get the second order of ADS in t and x

$$\left\{ \begin{array}{l} \frac{v_n^k - v_{n-1}^{k-1}}{\tau} + \frac{q_2(v_{n+1}^k - v_{n-1}^k)}{2h} + \frac{q_3(v_{n+1}^k - 2v_n^k + v_{n-1}^k)}{h^2} \\ + \frac{\tau q_0(v_{n+2}^k - 3v_{n+1}^k + 3v_n^k - v_{n-1}^k)}{h^3} + \frac{\tau q_1(v_{n+2}^k - 4v_{n+1}^k + 6v_n^k - 4v_{n-1}^k + v_{n-2}^k)}{h^4} \\ = \theta_n^k + p(x_n) - \frac{\tau}{2} \cdot \frac{(3+2 \sin x_n)(p(x_{n+1}) - 2p(x_n) + p(x_{n-1}))}{h^2} \\ - \frac{\tau}{2} \cdot \frac{\cos(x_n)(p(x_{n+1}) - p(x_{n-1}))}{h} + \frac{\tau p(x_n)}{2}, \\ k = 1, \dots, N, \quad n = 2, \dots, M - 2, \\ -3v_0^k + 4v_1^k - v_2^k = 0, \quad -3v_M^k + 4v_{M-1}^k - v_{M-2}^k = 0, \\ 10v_0^k - 15v_1^k + 6v_2^k - v_3^k = 0, \\ 10v_M^k - 15v_{M-1}^k + 6v_{M-2}^k - v_{M-3}^k = 0, \quad k = 0, \dots, N, \quad n = 0, \dots, M, \\ v_n^N = \varphi_n, \quad v_n^0 - (1 - \rho)v_n^l - \rho v_n^{l+1} = \psi(x_n), \quad n = 0, \dots, M \end{array} \right.$$

for the approximate solution of the nonlocal BVP (10).

Later, we calculate $p(x_n)$ by using (12), where $\{u_n^k\}$ is solution of the difference problem

$$\left\{ \begin{array}{l} \frac{u_n^k - u_{n-1}^{k-1}}{\tau} + \frac{q_2(u_{n+1}^k - u_{n-1}^k)}{2h} + \frac{q_3(u_{n+1}^k - 2u_n^k + u_{n-1}^k)}{h^2} + \frac{\tau}{2} \frac{q_0(u_{n+2}^k - 2u_{n+1}^k + 2u_{n-1}^k - u_{n-2}^k)}{2h^3} \\ + \frac{\tau}{2} \frac{q_1(u_{n+2}^k - 4u_{n+1}^k + 6u_n^k - 4u_{n-1}^k + u_{n-2}^k)}{h^4} = \theta_n^k, \quad k = 1, \dots, N, \quad n = 2, \dots, M - 2, \\ -3u_0^k + 4u_1^k - u_2^k = 0, \quad -3u_M^k + 4u_{M-1}^k - u_{M-2}^k = 0, \\ 10u_0^k - 15u_1^k + 6u_2^k - u_3^k = 0, \\ 10u_M^k - 15u_{M-1}^k + 6u_{M-2}^k - u_{M-3}^k = 0, \quad k = 0, \dots, N, \\ u_n^0 - (1 - \rho)u_n^l - \rho u_n^{l+1} = \psi(x_n), \quad n = 0, \dots, M, \end{array} \right. \tag{15}$$

which is the second order of ADS for nonlocal BVP (11).

For computational reasons it is convenient to rewrite the system (15) in the following matrix form

$$\begin{aligned} A_n u_{n+2} + B_n u_{n+1} + C_n u_n + D_n u_{n-1} + E_n u_{n-2} &= I \theta_n, \quad n = 2, \dots, M - 2, \\ -3u_0 + 4u_1 - u_2 &= \vec{0}, \quad -3u_M + 4u_{M-1} - u_{M-2} = \vec{0}, \\ 10u_0 - 15u_1 + 6u_2 - u_3 &= \vec{0}, \quad 10u_M - 15u_{M-1} + 6u_{M-2} - u_{M-3} = \vec{0}, \end{aligned} \tag{16}$$

where θ_n is column vector, $A_n, B_n, C_n, D_n, E_n, I$ are $(N + 1) \times (N + 1)$ square matrices, R is $N \times N$ identity matrix,

$$\begin{aligned} A_n &= \begin{bmatrix} 0 & \dots & 0 & 0 \\ & & & 0 \\ e_n R & & & \vdots \\ & & & 0 \end{bmatrix}, \quad B_n = \begin{bmatrix} 0 & \dots & 0 & 0 \\ & & & 0 \\ y_n R & & & \vdots \\ & & & 0 \end{bmatrix}, \\ D_n &= \begin{bmatrix} 0 & \dots & 0 & 0 \\ & & & 0 \\ z_n R & & & \vdots \\ & & & 0 \end{bmatrix}, \quad E_n = \begin{bmatrix} 0 & \dots & 0 & 0 \\ & & & 0 \\ w_n R & & & \vdots \\ & & & 0 \end{bmatrix}, \\ C_n &= \begin{bmatrix} 1 & 0 & 0 & \dots & -(1 - \rho) & \rho & \dots & 0 & 0 & 0 \\ r_n & d & 0 & \dots & 0 & 0 & \dots & 0 & 0 & 0 \\ 0 & r_n & d & \dots & 0 & 0 & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \dots & \dots & \dots & \dots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & 0 & \dots & r_n & d & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 & \dots & 0 & r_n & d \end{bmatrix}, \quad \theta_n = \begin{bmatrix} \theta_n^0 \\ \vdots \\ \theta_n^N \end{bmatrix}, \end{aligned}$$

$$\begin{aligned} e_n &= \frac{\tau q_0}{4h^3} + \frac{\tau q_1}{2h^4}, \quad y_n = \frac{q_2}{2h} + \frac{1}{h^2} q_3 - \frac{\tau q_0}{2h^3} - \frac{2\tau q_1}{h^4}, \\ r_n &= 1 + \frac{1}{\tau} + \frac{\tau}{2} - \frac{2}{h^2} q_3 + \frac{3\tau q_1}{h^4}, \\ z_n &= -\frac{q_2}{2h} + \frac{1}{h^2} q_3 + \frac{\tau q_0}{h^3} - \frac{2\tau q_1}{h^4}, \\ w_n &= -\frac{\tau q_0}{4h^3} + \frac{\tau q_1}{2h^4}, \quad n = 2, \dots, M-2. \end{aligned}$$

We search solution of linear system equation (16) in the next form

$$\begin{aligned} u_n &= \alpha_{n+1}u_{n+1} + \beta_{n+1}u_{n+2} + \gamma_{n+1}, \quad n = M-2, \dots, 0, \\ u_M &= D_M^{-1}((3I - 2\alpha_{M-2})\gamma_{M-1} - 3\gamma_{M-2}), \\ u_{M-1} &= D_M^{-1}[(4I - \alpha_{M-2})\gamma_{M-1} - \gamma_{M-2}], \\ D_M &= (\beta_{M-2} + 5I) - (4I - \alpha_{M-2})\alpha_{M-1}, \end{aligned}$$

where

$$\gamma_0 = \gamma_1 = \vec{0}, \quad \alpha_0 = \frac{4}{3}I, \quad \beta_0 = -\frac{1}{3}I, \quad \alpha_1 = \frac{8}{5}I, \quad \beta_1 = -\frac{3}{5}I$$

$$\gamma_{M-2} = \gamma_{M-3} = \vec{0}, \quad \alpha_{M-2} = 4I, \quad \beta_{M-2} = -3I, \quad \alpha_{M-3} = \frac{8}{5}I, \quad \beta_{M-3} = -\frac{3}{5}I,$$

and

$$\begin{aligned} F_n &= (C_n + D_n\alpha_{n-1} + E_n\beta_{n-2} + E_n\alpha_{n-2}\alpha_{n-1}), \quad n = 2, \dots, M-4. \\ \alpha_n &= -F_n^{-1}(B_n + D_n\beta_{n-1} + E_n\alpha_{n-2}\beta_{n-1}), \quad \beta_n = -F_n^{-1}A_n, \\ \gamma_n &= -F_n^{-1}(I\varphi_n - D_n\gamma_{n-1} - E_n\alpha_{n-2}\gamma_{n-1} - E_n\gamma_{n-2}), \end{aligned}$$

$$\begin{aligned} Q_{11} &= -3B_{M-2} - 8C_{M-2} - 8D_{M-2}\alpha_{M-3} - 3D_{M-2}\beta_{M-3} \\ &\quad - 8E_{M-2}\alpha_{M-4}\alpha_{M-3} - 3E_{M-2}\alpha_{M-4}\beta_{M-3} - 8E_{M-2}\beta_{M-4}, \\ Q_{12} &= A_{M-2} + 4B_{M-2} + 9C_{M-2} + 9D_{M-2}\alpha_{M-3} + 4D_{M-2}\beta_{M-3} \\ &\quad + 9E_{M-2}\alpha_{M-4}\alpha_{M-3} + 4E_{M-2}\alpha_{M-4}\beta_{M-3} + 9E_{M-2}\beta_{M-4}, \\ Q_{21} &= A_{M-1} - 3C_{M-1} - 8D_{M-1} - E_{M-1}(8\alpha_{M-3} + 3\beta_{M-3}), \\ Q_{22} &= B_{M-1} + 4C_{M-1} + 9D_{M-1} + E_{M-1}(9\alpha_{M-3} + 4\beta_{M-3}), \\ G_1 &= I\varphi_{M-2} - D_{M-2}\gamma_{M-3} - E_{M-2}\alpha_{M-4}\gamma_{M-3} - E_{M-2}\gamma_{M-3}, \\ G_2 &= I\varphi_{M-1} - E_{M-1}\gamma_{M-3}, \\ u_M &= (Q_{11} - Q_{12}Q_{22}^{-1}Q_{21})^{-1}(G_1 - Q_{12}Q_{22}^{-1}G_2), \\ u_{M-1} &= Q_{22}^{-1}(G_2 - Q_{21}u_M). \end{aligned}$$

Numerical illustration is carried out by using MATLAB program. Solutions of DSs are computed for different values of (N, M) . v_n^k and u_n^k correspond to the corresponding numerical values of $v(t, x)$ and $u(t, x)$ at $(t, x) = (t_k, x_n)$ and p_n represents the numerical value of $p(x)$ at point $x = x_n$. The errors are computed by

$$\begin{aligned} Ev_M^N &= \max_{0 \leq k \leq N} \left(\sum_{n=1}^{M-1} |v(t_k, x_n) - v_n^k|^2 h \right)^{\frac{1}{2}}, \\ Eu_M^N &= \max_{0 \leq k \leq N} \left(\sum_{n=1}^{M-1} |u(t_k, x_n) - u_n^k|^2 h \right)^{\frac{1}{2}}, \\ Ep_M &= \left(\sum_{n=1}^{M-1} |p(x_n) - p_n|^2 h \right)^{\frac{1}{2}}. \end{aligned}$$

Tables 1 and 2 illustrate the errors between the exact and approximate solutions of DSs for various N and M , respectively. It can be seen from output results that the second order of ADS is more accurate than the first order of ADS. The error analysis shown in Tables 1 and 2 indicate that both DSs have correct convergence rates.

Table 1

Mesh grid absolute value of difference between exact solution and solution of first order of ADS

N=M	20	40	80	160
Ev_M^N	0.034277	0.016674	0.008483	0.004278
Ep_M	0.086716	0.043925	0.022123	0.011104
Ev_M^N	0.152320	0.075113	0.037321	0.018601

Table 2

Mesh grid absolute value of difference between exact solution and solution of second order of ADS

N=M	20	40	80	160
Ev_M^N	0.020123	0.004141	0.000920	0.000217
Ep_M	0.08946	0.024373	0.006796	0.001926
Ev_M^N	0.089678	0.018803	0.004188	0.000969

Conclusion

In this work, SIP for a multi-dimensional parabolic partial differential equation with multi-point nonlocal boundary condition was studied. Stability estimates for solution of inverse problem were obtained. Well-posedness of three SIPs for the reverse parabolic partial differential equations was established.

Author Contributions

All authors contributed equally to this work.

Conflict of Interest

The authors declare no conflict of interest.

References

- 1 Andreev, V.K., & Stepanova, I.V. (2021). Inverse problem for source function in parabolic equation at Neumann boundary Conditions. *Journal of Siberian Federal University. Mathematics & Physics*, 14(4), 445–451. <https://doi.org/10.17516/1997-1397-2021-14-4-445-451>
- 2 Ashyraliyev, M., & Ashyralyeva, M. (2024). Stable difference schemes for hyperbolic–parabolic equations with unknown parameter. *Bol. Soc. Mat. Mex.*, 30, 14. <https://doi.org/10.1007/s40590-023-00585-1>
- 3 Ashyralyev, A. (2011). On the problem of determining the parameter of a parabolic equation. *Ukrainian Mathematical Journal*, 62(9), 1397–1408. <https://doi.org/10.1007/s11253-011-0438-7>
- 4 Ashyralyev, A., & Ashyralyev, C. (2022). On the stability of parabolic differential and difference equations with a time-nonlocal condition. *Computational Mathematics and Mathematical Physics*, 62(6), 962–973. <https://doi.org/10.1134/S0965542522060021>
- 5 Ashyralyev, A., & Erdogan, A.S. (2021). Parabolic time dependent source identification problem with involution and Neumann condition. *Bulletin of the Karaganda University. Mathematics Series*, 2(102), 5–15. <https://doi.org/10.31489/2021M2/16-24>
- 6 Ashyralyev, A., & Sobolevskii, P.E. (1994). *Well-posedness of parabolic difference equations*, 69 of *Operator Theory: Advances and Applications*. Birkhäuser, Basel, Switzerland.

- 7 Ashyralyev, A., Ashyraliyev, M., & Ashyralyyeva, M.A. (2020). Identification problem for telegraph-parabolic equations. *Computational Mathematics and Mathematical Physics*, 60(8), 1294–1305. <https://doi.org/10.1134/S0965542520080035>
- 8 Ashyralyev, A., Erdogan, A.S., Demirdag, O. (2012). On the determination of the right-hand side in a parabolic equation. *Appl. Numer. Math.*, 62, 1672–168. <https://doi.org/10.1016/j.apnum.2012.05.008>
- 9 Ashyralyev, A., Ashyralyyev, C., & Ahmed, A.M.S. (2023). Numerical solution of the boundary value problems for the parabolic equation with involution. *Bulletin of the Karaganda University. Mathematics Series*, 1(109), 48–57. <https://doi.org/10.31489/2023M1/48-57>
- 10 Ashyralyev, A., Erdogan, A.S., & Sarsenbi, A. (2020). A note on the parabolic identification problem with involution and Dirichlet condition. *Bulletin of the Karaganda University. Mathematics Series*, 3(99), 130–139. <https://doi.org/10.31489/2020M3/130-139>
- 11 Ashyralyyev, C. (2020). Stability of Rothe difference scheme for the reverse parabolic problem with integral boundary condition. *Math. Meth. Appl. Sci.*, 43, 5369–5379. <https://dx.doi.org/10.1002/mma.6278>
- 12 Ashyralyyev, C., & Demirdag, O. (2012). The difference problem of obtaining the parameter of a parabolic equation. *Abstr. Appl. Anal.*, 2012, 1–14. <https://doi.org/10.1155/2012/603018>
- 13 Ashyralyyev, C., Dural, A., & Sozen, Y. (2012). Finite difference method for the reverse parabolic problem. *Abstr. Appl. Anal.*, 2012, 1–17. <https://doi.org/10.1155/2012/294154>
- 14 Ashyralyyev, C., Dural, A., & Sozen, Y. (2012). Finite difference method for the reverse parabolic problem with Neumann condition. *AIP Conf. Proceedings*, 1470, 102–105. <https://dx.doi.org/10.1063/1.4747650>
- 15 Ashyralyyev, C. & Akkan, P. (2020). Source identification problem for a parabolic equation with multi-point nonlocal boundary condition. *Numerical Functional Analysis and Optimization*, 41(16), 1913–1935. <https://dx.doi.org/10.1080/01630563.2020.1802748>
- 16 Ashyralyyev, C., & Gonenc, A. (2021). Crank–Nicolson difference scheme for reverse parabolic nonlocal problem with integral and Neumann boundary conditions. *Int. J. Appl. Math.*, 34(2), 273–282. <https://dx.doi.org/10.12732/ijam.v34i2.5>
- 17 Azizbayov, E.I., & Mehraliyev, Y.T. (2017). Solvability of nonlocal inverse boundary-value problem for a second-order parabolic equation with integral conditions. *Electron. J. Differ. Equ.*, 2017(125), 1–14.
- 18 Eidel'man, Y.S. (1991). An inverse problem for an evolution equation. *Math. Notes*, 49(5), 535–540. <https://doi.org/10.1007/BF01142653>
- 19 Imanbetova, A.B., Sarsenbi, A., & Seilbekov, B. (2024). On solvability of the inverse problem for a fourth-order parabolic equation with a complex-valued coefficient. *Bulletin of the Karaganda University. Mathematics Series*, 1(113), 60–72. <https://doi.org/10.31489/2024M1/60-72>
- 20 Imanuvilov, O., & Yamamoto, M. (2024). Inverse parabolic problems by Carleman estimates with data taken at initial or final time moment of observation. *Inverse Problems and Imaging*, 18(2), 366–387. <https://doi.org/10.3934/ipi.2023036>
- 21 Kamynin, V.L. (2022). The inverse problem of recovering the source function in a multidimensional nonuniformly parabolic equation. *Math. Notes*, 112, 412–423. <https://doi.org/10.1134/S0001434622090103>
- 22 Kozhanov, A.I. (2017). Parabolic equations with unknown time-dependent coefficients. *Comput. Math. Math. Phys.*, 57(6), 961–972. <https://doi.org/10.1134/S0965542517060082>
- 23 Kozhanov, A.I., Abulkayirov, U.U., & Ashurova, G.R. (2022). Inverse problems of determining coefficients of time type in a degenerate parabolic equation. *Bulletin of the Karaganda University*.

- Mathematics Series*, 2(106), 128–142. <https://doi.org/10.31489/2022M2/128-142>
- 24 Pyatkov, S.G., & Samkov, M.L. (2017). Solvability of some inverse problems for the nonstationary heat-and-mass-transfer system. *Journal of Mathematical Analysis and Applications*, 446(2), 1449–1465. <https://doi.org/10.1016/j.jmaa.2016.09.040>
- 25 Sadybekov, M., Oralsyn, G., & Ismailov, M. (2018). Determination of a time-dependent heat source under not strengthened regular boundary and integral overdetermination conditions. *Filomat*, 32(3), 809–814. <http://dx.doi.org/10.2298/FIL1803809S>
- 26 Sazaklioglu, A.U. (2022). On the numerical solutions of some identification problems for one- and multidimensional parabolic equations backward in time. *Applied Numerical Mathematics*, 181, 76–93. <https://doi.org/10.1016/j.apnum.2022.05.015>
- 27 Sazaklioglu, A.U. (2024). An iterative numerical method for an inverse source problem for a multidimensional nonlinear parabolic equation. *Applied Numerical Mathematics*, 198, 428–447. <http://dx.doi.org/10.1016/j.apnum.2024.02.001>
- 28 Sobolevskii, P.E. (1975). *Raznostnye Metody Priblizhennogo Resheniia Differentsialnykh Uravnenii [Difference Methods for the Approximate Solution of Differential Equations]*. Voronezh: Izdatelstvo VGU [in Russian].

*Author Information**

Charyyar Ashyralyyev (*corresponding author*) — Doctor of physical and mathematical sciences, Professor of the Department of Mathematics, Faculty of Engineering and Natural Sciences, Bahcesehir University, Istanbul, Turkey; Khoja Akhmet Yassawi International Kazakh-Turkish University, Turkestan, Kazakhstan; National University of Uzbekistan named after Mirzo Ulugbek, Tashkent, Uzbekistan; e-mail: charyyar@gmail.com; <https://orcid.org/0000-0002-6976-2084>

Tazegul Aydogdyevna Ashyralyyeva — Lecturer, Department of Higher Mathematics, Yagshygeldi Kakayev International University of Oil and Gas, Ashgabat, Turkmenistan; e-mail: tazegulashyralyyewa@gmail.com; <https://orcid.org/0009-0007-6341-9334>

*The author's name is presented in the order: First, Middle and Last Names.