

L.Kh. Gadzova*

*Institute of Applied Mathematics and Automation KBSC RAS, Nalchik, Russia
(E-mail: macaneeva@mail.ru)*

Generalized boundary value problem for a linear ordinary differential equation with a discretely distributed fractional differentiation operator

This paper formulates and solves a generalized boundary value problem for a linear ordinary differential equation with a discretely distributed fractional differentiation operator. The fractional derivative is understood as the Gerasimov–Caputo derivative. The boundary conditions are given in the form of linear functionals, which makes it possible to cover a wide class of linear local and non-local conditions. A representation of the solution is found in terms of special functions. A necessary and sufficient condition for the solvability of the problem under study is obtained, as well as conditions under which the solvability condition is certainly satisfied. The theorem of existence and uniqueness of the solution is proved.

Keywords: fractional differentiation operator, Caputo derivative, boundary value problem, functional, Wright function.

Introduction and statement of the problem

In the interval $0 < x < 1$, let us consider the equation

$$\sum_{j=1}^m \beta_j \partial_{0x}^{\alpha_j} u(x) + \lambda u(x) = f(x), \quad (1)$$

where $\alpha_j \in (1, 2)$, $\lambda, \beta_j \in \mathbb{R}$, $\beta_1 > 0$, $\alpha_1 > \alpha_2 > \dots > \alpha_m$, $\partial_{0x}^{\gamma} u(x)$ is the Caputo derivative [1; 11]:

$$\partial_{sx}^{\gamma} u(x) = \text{sign}^n(x-s) D_{sx}^{\gamma-n} u^{(n)}(x), \quad n-1 < \gamma \leq n, \quad n \in \mathbb{N}, \quad (2)$$

and D_{sx}^{γ} is the Riemann–Liouville fractional integro-differentiation operator of order γ with respect to the variable x [1; 9], which is defined by the formula

$$D_{sx}^{\gamma} u(x) = \frac{\text{sign}(x-s)}{\Gamma(-\gamma)} \int_s^x \frac{u(t) dt}{|x-t|^{\gamma+1}}, \quad \gamma < 0,$$

$$D_{sx}^{\gamma} u(x) = u(x), \quad \gamma = 0,$$

$$D_{sx}^{\gamma} u(x) = \text{sign}^n(x-s) \frac{d^n}{dx^n} D_{sx}^{\gamma-n} u(x), \quad n-1 < \gamma \leq n, \quad n \in \mathbb{N}.$$

Operator (2) is also known in the literature as the Gerasimov–Caputo operator [2, 3].

At present, differential equations of fractional order are being extensively studied in connection with practical applications in various areas of physics and mathematical modeling. The theory of fractional differential equations has proven itself well in the study of «classical» viscoelastic models. All this is supported by new applied problems [4–8].

*Corresponding author.

E-mail: macaneeva@mail.ru

One of the first works devoted to fractional calculus and its applications is the monograph [9]. The main theoretical results and solution methods are reflected in the works [7], [6], and [10].

Linear fractional ordinary differential equations were studied by many authors; a detailed bibliography on this subject can be found in [1, 5, 6, 11]. A significant contribution to the study of fractional differential equations was made by the authors of [12–15].

Differential equations with discretely distributed differentiation operator

$$\sum_{k=1}^m \lambda_k \frac{\partial^{\sigma_k}}{\partial y^{\sigma_k}}$$

can be treated as an operator

$$\int_{\alpha}^{\beta} \left(\lambda(y, t) \frac{\partial^t}{\partial y^t} \right) d\mu(t)$$

with a measure concentrated on a discrete set [16, 17].

Differential equations with discretely distributed differentiation operators and continuously distributed differentiation operators were studied in [18–20], where equations with discretely distributed differentiation operators were used to search for approximate solutions of equations with continuously distributed differentiation operators. We also note the papers [16], [17], [21], [22], where equations with fractional discretely distributed differentiation operators were studied.

In this paper, we investigate a generalized boundary value problem (in the terminology of M.A. Naimark) for equation (1) [22; 16]. An explicit representation of the solution of the problem under study is constructed, a condition for unique solvability is found, and a uniqueness theorem for the solution is proved. We specified boundary conditions in the form of linear functionals, which makes it possible to cover a wide class of linear local and nonlocal conditions. Various boundary value problems for equation (1) were studied in the works [23–25]. Note also that in work [26] a generalized boundary value problem for an ordinary differential equation of fractional order with general conditions was investigated.

A *regular solution* to equation (1) is said to be a function $u = u(x)$ that has an absolutely continuous first-order derivative on the closed interval $[0, 1]$ and satisfies equation (1) for all $x \in (0, 1)$.

Problem. Find a regular solution to equation (1) in the interval $(0, 1)$, which satisfies the conditions

$$\ell_0[u] = u_0, \tag{3}$$

$$\ell_1[u] = u_1, \tag{4}$$

where u_0, u_1 are given real numbers, ℓ_0, ℓ_1 are linear bounded functionals in $C^1[0, 1]$.

Notation and auxiliary statements

We use the following notation (see [27])

$$G_m^\mu(x) = G_m^\mu(x; \nu_1, \dots, \nu_m; \gamma_1, \dots, \gamma_m) \equiv \int_0^\infty e^{-t} S_m^\mu(x; \nu_1 t, \dots, \nu_m t; \gamma_1, \dots, \gamma_m) dt,$$

$$\nu_1 = -\frac{\lambda}{\beta_1}, \quad \nu_j = -\frac{\beta_j}{\beta_1}, \quad \gamma_1 = \alpha_1, \quad \gamma_j = \alpha_1 - \alpha_j, \quad (j = \overline{2, m}),$$

$$S_m^\mu(x; z_1, \dots, z_m; \gamma_1, \dots, \gamma_m) = (h_1 * h_2 * \dots * h_m)(x),$$

by

$$(g * h)(x) = \int_0^x g(x-t)h(t)dt$$

we denote the Laplace convolution of the functions $g(x)$ and $h(x)$,

$$h_j = h_j(x) \equiv x^{\mu_j-1} \phi(\gamma_j, \mu_j; z_j x^{\gamma_j}),$$

where

$$\phi(\rho, \zeta; z) = \sum_{k=0}^{\infty} \frac{z^k}{k! \Gamma(\rho k + \zeta)}$$

is the Wright function (see [28]).

Further, we assume that the parameters $G_m^\mu(x)$ range over

$$x > 0, \quad z_i \in \mathbb{R}, \quad \gamma_j > 0, \quad \mu_j > 0.$$

We note that the function $G_m^\mu(x)$ is independent of the distribution of the numbers $\mu_j > 0$, but only depends on their sum $\mu = \sum_{j=1}^m \mu_j$.

The following equalities for the function $G_m^\mu(x)$ (see [24]) hold:

$$G_m^\mu(x) = O(x^{\mu-1}) \quad \text{при } x \rightarrow 0,$$

$$D_{0x}^\nu G_m^\mu(x) = G_m^{\mu-\nu}(x), \quad \text{если } \mu > \nu, \tag{5}$$

$$G_m^\mu(x) - \sum_{j=1}^m \nu_j D_{0x}^{-\gamma_j} G_m^\mu(x) = \frac{x^{\mu-1}}{\Gamma(\mu)}, \quad \mu > 0. \tag{6}$$

In particular, from equalities (5) and (6) we obtain the formula (see [25])

$$\sum_{j=1}^m \beta_j G_m^{\mu-\gamma_j}(x) + \lambda G_m^\mu(x) = \frac{\beta_1 x^{\mu-\alpha_1-1}}{\Gamma(\mu - \alpha_1)}, \quad \mu > \alpha_1.$$

We also need the following auxiliary statement proved in (see [26]).

It should be noted that hereinafter the ℓ functionality is applied to the function depending on x .

Lemma. Let $K(x, t) \in C([0, 1] \times [0, 1])$ and $\frac{\partial}{\partial x} K(x, t) \in C([0, 1] \times [0, 1])$, ℓ – linear bounded functional in space $C^1[0, 1]$. Then the following relation is true

$$\ell \left[\int_0^1 K(x, t) dt \right] = \int_0^1 \ell[K(x, t)] dt. \tag{7}$$

Main result

Theorem. Let a function $f(x)$ satisfy the conditions

$$x^{1-\mu} f(x) \in C[0, 1], \quad f(x) = D_{0x}^{\alpha_1-2} g(x), \quad g(x) \in L[0, 1], \quad \mu > 0.$$

and the inequality

$$\det A = \ell_0[\mathcal{W}_2(x)] \ell_1[\mathcal{W}_3(x)] - \ell_0[\mathcal{W}_3(x)] \ell_1[\mathcal{W}_2(x)] \neq 0 \tag{8}$$

be fulfilled. Then a function $u(x)$ defined by the relation

$$u(x) = \int_0^1 f(t)\mathcal{T}(x,t)dt + \overline{\mathcal{W}}(x)\mathcal{H} \begin{pmatrix} u_0 \\ u_1 \end{pmatrix}, \tag{9}$$

$$\mathcal{T}(x,t) = [1 - \overline{\mathcal{W}}(x)\mathcal{H}\bar{\ell}] \mathcal{W}_1(x-t), \quad \overline{\mathcal{W}}(x) = (\mathcal{W}_2(x), \mathcal{W}_3(x)), \quad \bar{\ell} = \begin{pmatrix} \ell_0 \\ \ell_1 \end{pmatrix},$$

$$\mathcal{W}_1(x) = \frac{1}{\beta}G_m^{\alpha_1}(x); \quad \mathcal{W}_2(x) = x + \nu_1 G_m^{\alpha_1+2}(x); \quad \mathcal{W}_3(x) = 1 + \nu_1 G_m^{\alpha_1+1}(x),$$

$$A = \bar{\ell} [\overline{\mathcal{W}}(x)] = \begin{pmatrix} \ell_0[\mathcal{W}_2(x)] & \ell_0[\mathcal{W}_3(x)] \\ \ell_1[\mathcal{W}_2(x)] & \ell_1[\mathcal{W}_3(x)] \end{pmatrix}, \quad \mathcal{H} = A^{-1} = \frac{1}{\det A} \begin{pmatrix} \ell_1[\mathcal{W}_3(x)] & -\ell_0[\mathcal{W}_3(x)] \\ -\ell_1[\mathcal{W}_2(x)] & \ell_0[\mathcal{W}_2(x)] \end{pmatrix},$$

is a regular solution to problem (1), (3), (4). The solution to problem (1), (3), (4) is unique if and only if the condition (8) is satisfied.

Proof. Let $u(x)$ be a regular solution to the problem (1), (3), (4). To find a solution to the problem (1), (3), (4) we use the solution of the Cauchy problem for the equation (1), which can be represented as [24, 25]:

$$u(x) = \int_0^1 f(t)\mathcal{W}_1(x-t)dt + \overline{\mathcal{W}}(x) \begin{pmatrix} C_1 \\ C_2 \end{pmatrix}. \tag{10}$$

Further, taking into account the introduced notation and equalities (7), we satisfy (10) the boundary conditions

$$\int_0^1 f(t)\bar{\ell}[\mathcal{W}_1(x-t)]dt + A \begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} u_0 \\ u_1 \end{pmatrix}.$$

From this we find

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathcal{H} \begin{pmatrix} u_0 \\ u_1 \end{pmatrix} - \int_0^1 f(t)\mathcal{H}\bar{\ell}[\mathcal{W}_1(x-t)]dt.$$

After elementary transformations, substituting the found value into (10), we obtain a representation of the solution to problem (1), (3), (4) in the form (8). This, in particular, implies the uniqueness of the solution.

Let us now check the fulfillment of the boundary conditions (3), (4).

$$\bar{\ell} \left[\int_0^1 f(t)\mathcal{T}(x,t)dt + \overline{\mathcal{W}}(x)\mathcal{H} \begin{pmatrix} u_0 \\ u_1 \end{pmatrix} \right] = I_1 + I_2,$$

where

$$I_1 = \bar{\ell} \left[\int_0^1 f(t)\mathcal{T}(x,t)dt \right] \quad \text{end} \quad I_2 = \bar{\ell} \left[\overline{\mathcal{W}}(x)\mathcal{H} \begin{pmatrix} u_0 \\ u_1 \end{pmatrix} \right].$$

Taking into account (7), we have

$$I_1 = \int_0^1 \bar{\ell} [\mathcal{T}(x,t)] f(t)dt = \int_0^1 (\bar{\ell}[\mathcal{W}_1(x,t)] - \bar{\ell}[\overline{\mathcal{W}}(x)]\mathcal{H}\bar{\ell}[\mathcal{W}_1(x,t)]) f(t)dt.$$

By virtue of the equality $\bar{\ell}[\bar{\mathcal{W}}(x)] = \mathcal{H}^{-1}$, we obtain that $I_1 = 0$. Similarly, for I_2 we have

$$I_2 = \bar{\ell}[\bar{\mathcal{W}}(x)]\mathcal{H} \begin{pmatrix} u_0 \\ u_1 \end{pmatrix} = \mathcal{H}^{-1}\mathcal{H} \begin{pmatrix} u_0 \\ u_1 \end{pmatrix} = \begin{pmatrix} u_0 \\ u_1 \end{pmatrix}.$$

Let us prove that the function $u(x)$, given by equality (9), is a solution to problem (3), (4) for equation (1).

$$\left[\sum_{j=1}^m \beta_j \partial_{0x}^{\alpha_j} + \lambda \right] y_i = f(x), \quad i = \overline{1, 3},$$

$$y_1 = \int_0^x f(t)\mathcal{W}_1(x-t)dt, \quad y_2 = \int_0^x f(t)\bar{\mathcal{W}}(x)\mathcal{H}\bar{\ell}\mathcal{W}_1(x-t)dt, \quad y_3 = \bar{\mathcal{W}}(x)\mathcal{H} \begin{pmatrix} u_0 \\ u_1 \end{pmatrix}.$$

Taking into account relations (5) – (8) and since the functions y_2 and y_3 is a linear combinations of the functions $\mathcal{W}_i(x), i = \overline{1, 3}$ we have

$$\left[\sum_{j=1}^m \beta_j \partial_{0x}^{\alpha_j} + \lambda \right] y_i = 0, \quad i = \overline{2, 3}.$$

Considering that (see [25])

$$\sum_{j=1}^m \beta_j \partial_{0x}^{\alpha_j} \int_0^x f(t)\mathcal{W}_1(x-t)dt = -\frac{\lambda}{\beta_1} \int_0^x f(t)G_m^\alpha(x-t)dt + f(x),$$

we get

$$\left[\sum_{j=1}^m \beta_j \partial_{0x}^{\alpha_j} + \lambda \right] y_1 = f(x).$$

It means that the solution satisfies equation (1).

Let us show that if condition (8) is satisfied, that is,

$$\ell_0[\mathcal{W}_2(x)]\ell_1[\mathcal{W}_3(x)] - \ell_0[\mathcal{W}_3(x)]\ell_1[\mathcal{W}_2(x)] = 0, \tag{11}$$

then the solution to the homogeneous problem is not unique.

Consider the function

$$\tilde{u}(x) = (k_1(x), k_2(x)) \begin{pmatrix} C_1 \\ C_2 \end{pmatrix},$$

where C_1 and C_2 are arbitrary constants,

$$k_1(x) = \bar{\mathcal{W}}(x) \begin{pmatrix} \ell_0[\mathcal{W}_2(x)] \\ -\ell_0[\mathcal{W}_3(x)] \end{pmatrix}, \quad k_2(x) = \bar{\mathcal{W}}(x) \begin{pmatrix} -\ell_1[\mathcal{W}_2(x)] \\ \ell_1[\mathcal{W}_3(x)] \end{pmatrix}.$$

Then it follows from (11) that the function $\tilde{u}(x)$ is a solution to a homogeneous problem

$$\sum_{j=1}^m \beta_j \partial_{0x}^{\alpha_j} \tilde{u}(x) + \lambda \tilde{u}(x) = 0, \quad \ell_0[\tilde{u}] = 0, \quad \ell_1[\tilde{u}] = 0.$$

References

- 1 Нахушев А.М. Дробное исчисление и его применение / А.М. Нахушев. — М.: Физматлит, 2003. — 272 с.
- 2 Килбас А.А. Теория и приложения дифференциальных уравнений дробного порядка (курс лекций). Методологическая школа-конференция «Математическая физика и нанотехнологии» / А.А. Килбас. — Самара, 2009. — 121 с.
- 3 Новоженова О.Г. Биография и научные труды Алексея Никифоровича Герасимова. О линейных операторах, упруго-вязкости, элвтерозе и дробных производных / О.Г. Новоженова. — М.: Перо, 2018. — 235 с.
- 4 Джрбашян М.М. Интегральные преобразования и представления функций в комплексной области / М.М. Джрбашян. — М.: Наука, 1966. — 672 с.
- 5 Самко С.Г. Интегралы и производные дробного порядка и некоторые их приложения / С.Г. Самко, А.А. Килбас, О.И. Маричев. — Минск: Наука и техника, 1987. — 688 с.
- 6 Нахушев А.М. О континуальных аналогах реологических уравнений состояния и логистическом законе изменения вязкоупругих свойств полимера / А.М. Нахушев, Р.Б. Тхакахов // Докл. Адыгской (Черкесской) Междунар. акад. наук. — 1995. — 1. — № 2. — С. 6–11.
- 7 Podlubny I. Fractional Differential Equations / I. Podlubny. — ACADEMIC PRESS, 1999. — 198. — 340 p.
- 8 Kilbas A. A. Theory and applications of fractional differential equations / A. A. Kilbas, H. M. Srivastava, J. J. Trujillo. — North-Holland Math. Stud., Elsevier, Amsterdam, 2006. — 204. — 540 p.
- 9 Oldham K.B. The fractional calculus / K. B. Oldham, J. Spanier. — N.-Y.; L.: Acad. press, 1974. — 234 p.
- 10 Учайкин В.В. Метод дробных производных / В.В. Учайкин. — Ульяновск: Изд-во «Артишок», 2008. — 512 с.
- 11 Miller K.S. An introduction to the fractional calculus and fractional differential equations / K.S. Miller, B. Ross. — New York: Jon Wiley & Sons. Inc., 1993. — 366 p.
- 12 Псху А. В. Уравнения в частных производных дробного порядка / А. В. Псху. — М.: Наука, 2005. — 199 с.
- 13 Barrett J. H. Differential equations of non-integer order / J. H. Barrett // Canadian J. Math. — 1954. — 6. — No. 4. — P. 529–541.
- 14 Джрбашян М. М. Дробные производные и задача Коши для дифференциальных уравнений дробного порядка / М. М. Джрбашян, А. Б. Нерсесян // Изв. АН Армянской ССР. Матем. — 1968. — 3. — № 1. — С. 3–28.
- 15 Джрбашян М. М. Краевая задача для дифференциального оператора дробного порядка типа Штурма-Лиувилля / М. М. Джрбашян // Изв. АН Армян ССР. — 1970. — 5. — № 2. — С. 71–96.
- 16 Pskhu A. V. Boundary Value Problem for a First-Order Partial Differential Equation with a Fractional Discretely Distributed Differentiation Operator / A. V. Pskhu // Differential Equations. — 2016. — 52:12. — P. 1610-1623.
- 17 Псху А. В. Уравнение дробной диффузии с оператором дискретно распределенного дифференцирования / А. В. Псху // Sibirean electr. Math. Reports. — 2016. — 13. — С. 1078–1098.
- 18 Caputo M. Diffusion with space memory modelled with distributed order space fractional differential equations / M. Caputo // Annals of Geophysics. — 2003. — 46. — No. 2. — P. 223–234.

- 19 Diethelm K. Numerical analysis for distributed-order differential equations / K. Diethelm, N.J. Ford // J. of Comp. and Appl. Math. — 2009. — 225. — P. 96–104.
- 20 Jiao Z. Distributed-Order Dynamic Systems: Stability, Simulation, Applications and Perspectives / Z. Jiao, Y. Qu. Chen, I. Podlubny. — London, 2012. — 103 p.
- 21 Ozturk I. On the theory of fractional differential equation / I. Ozturk // Reports of Adyghe (Circassian) International Academy of Sciences. — 1998. — 3. — No. 2. — P. 35–39.
- 22 Наймарк М.А. Линейные дифференциальные операторы / М.А. Наймарк. — М.: Наука, 1969. — 528 с.
- 23 Gadzova L. Kh. Dirichlet and Neumann problems for a fractional ordinary differential equation with constant coefficients / L.Kh. Gadzova // Differential Equations. — 2015. — 51:12. — P. 1556–1562.
- 24 Gadzova L. Kh. Boundary Value Problem for a Linear Ordinary Differential Equation with a Fractional Discretely Distributed Differentiation Operator / L.Kh. Gadzova // Differential Equations. — 2018. — 54:2. — P. 178–184.
- 25 Gadzova L. Kh. Nonlocal Boundary-Value Problem for a Linear Ordinary Differential Equation with Fractional Discretely Distributed Differentiation Operator / L.Kh. Gadzova // Mathematical Notes. — 2019. — 106. — No. 5–6. — P. 904–908.
- 26 Гадзова Л.Х. Обобщённая краевая задача для обыкновенного дифференциального уравнения дробного порядка / Л.Х. Гадзова // Челяб. физ.-мат. журн. — 2022. — 7. — № 1. — С. 20-29. <https://doi.org/10.47475/2500-0101-2022-17102>
- 27 Pskhu, A.V. Initial-value problem for a linear ordinary differential equation of noninteger order / A.V. Pskhu // Sbornik: Mathematics. — 2011. — 202. — No. 4. — P. 571–582.
- 28 Wright E.M. On the coefficients of power series having exponential singularities / E.M. Wright // J. London Math. Soc. — 1933. — 8. — No.29. — P. 71–79.

Л.Х. Гадзова

РҒА КБҒО Қолданбалы математика және автоматтандыру институты, Нальчик, Ресей

Бөлшек дискретті үлестірілген дифференциалдау операторы бар сызықтық қарапайым дифференциалдық теңдеу үшін жалпыланған шеттік есеп

Мақалада бөлшек дискретті үлестірілген дифференциалдау операторы бар сызықтық қарапайым дифференциалдық теңдеу үшін жалпыланған шеттік есеп құрастырылып және шешілген. Бөлшек туынды Герасимов-Капуто туындысы мағынасында түсініледі. Шеттік шарттар сызықтық функционалдар түрінде берілген, бұл сызықтық жергілікті және жергілікті емес жағдайлардың жеткілікті кең класын қамтуға мүмкіндік береді. Шешімнің мәні арнайы функциялар арқылы табылды. Зерттелетін есептің шешілу мүмкіндігінің қажетті және жеткілікті шарты, сондай-ақ шешілу шарты сөзсіз орындалатын шарттар алынды. Шешімнің бар болуы және бірегейлігі теоремасы дәлелденді.

Кілт сөздер: бөлшек дифференциалдау операторы, Капуто туындысы, шеттік есеп, функционал, Райт функциясы.

Л.Х. Гадзова

Институт прикладной математики и автоматизации КБНЦ РАН, Нальчик, Россия

Обобщенная краевая задача для линейного обыкновенного дифференциального уравнения с оператором дробного дискретно-распределенного дифференцирования

В статье сформулирована и решена обобщенная краевая задача для линейного обыкновенного дифференциального уравнения с оператором дробного дискретно-распределенного дифференцирования. Дробная производная понимается в смысле производной Герасимова–Капуто. Краевые условия задаются в форме линейных функционалов, это позволяет охватить достаточно широкий класс линейных локальных и нелокальных условий. В терминах специальных функций найдено представление решения. Получено необходимое и достаточное условие разрешимости исследуемой задачи, а также условия, при которых условие разрешимости заведомо выполняется. Доказана теорема существования и единственности решения.

Ключевые слова: оператор дробного дифференцирования, производная Капуто, краевая задача, функционал, функция Райта.

References

- 1 Nakhushiev, A.M. (2003). *Drobnoe ischislenie i ego primeneniye [Fractional calculus and its application]*. Moscow: Fizmatlit [in Russian].
- 2 Kilbas, A.A. (2009). *Teoriya i prilozheniya differentsialnykh uravnenii drobnogo poriadka (kurs lektsii) [Theory and applications of fractional differential equations (course of lectures)]*. Metodologicheskaya shkola-konferentsiya «Matematicheskaya fizika i nanotekhnologii» — "Mathematical Physics and Nanotechnology" Methodological School-Conference [in Russian].
- 3 Novozhenova, O.G. (2018). *Biografiya i nauchnye trudy Alekseia Nikiforovicha Gerasimova. O lineinykh operatorakh, uprugoviazkosti, elevteroze i drobnnykh proizvodnykh [Biography and scientific works of Alexey Nikiforovich Gerasimov. On linear operators, elastic viscosity, eleutherosis and fractional derivatives]*. Moscow: Pero [in Russian].
- 4 Dzhrbashian, M.M. (1966). *Integralnye preobrazovaniya i predstavleniya funktsii v kompleksnoi oblasti [Integral transformations and representations of functions in the complex domain]*. Moscow: Nauka [in Russian].
- 5 Samko, S. G., Kilbas, A. A., & Marichev, O. I. (1987). *Integraly i proizvodnye drobnogo poriadka i nekotorye ikh prilozheniya [Integrals and derivatives of fractional order and some of their applications]*. Minsk: Nauka i tekhnika [in Russian].
- 6 Nakhushiev, A. M., & Tkhakakhov, R. B. (1995). O kontinualnykh analogakh reologicheskikh uravnenii sostoianiya i logisticheskome zakone izmeneniya viazkouprugikh svoystv polimera [On continual analogs of rheological equations of state and the logistic law of change in the viscoelastic properties of a polymer] *Doklady Adygskoi (Cherkesskoi) Mezhdunarodnoi akademii nauk – Reports of Adyghe (Circassian) International Academy of Sciences. 1, 2, 6–11* [in Russian].
- 7 Podlubny, I. (1999). *Fractional Differential Equations*. ACADEMIC PRESS, New York.
- 8 Kilbas, A. A., Srivastava, H. M., & Trujillo, J. J. (2006). *Theory and applications of fractional differential equations*. North-Holland Math. Stud., Elsevier, Amsterdam.
- 9 Oldham, K. B., & Spanier J. (1974). *The fractional calculus*. N.-Y.; L.: Acad. press.
- 10 Uchaikin, V. V. (2008). *Metod drobnnykh proizvodnykh [The method of fractional derivatives]*. Ulianovsk: Izdatelstvo «Artishok» [in Russian].

- 11 Miller, K.S., & Ross, B. (1993). *An introduction to the fractional calculus and fractional differential equations*. New York.: Jon Wiley & Sons. Inc.
- 12 Pskhu, A.V. (2005). Uravneniia v chastnykh proizvodnykh drobnogo poriadka [Partial differential equations of fractional order]. Moscow: Nauka [in Russian].
- 13 Barrett, J.H. (1954). Differential equation of non-integer order. *Canad. J. Math.*, 6, 4, 529–541.
- 14 Dzhrbashyan, M.M., & Nersesyan, A.B. (1968). Drobnnye proizvodnye i zadacha Koshi dlia differentsialnykh uravnenii drobnogo poriadka [Fractional derivatives and the Cauchy problem for differential equations of fractional order]. *Izvestiia Akademii nauk Armianskoi SSR – Proceedings of the Academy of Sciences of the Armenian SSR*, 3, 1, 3–28 [in Russian].
- 15 Dzhrbashyan, M.M. (1970). Kraevaia zadacha dlia differentsialnogo operatora drobnogo poriadka tipa Shturma-Liuvillia [Boundary problem for fractional differential operator of Sturm-Liouville type]. *Izvestiia Akademii nauk Armianskoi SSR – Proceedings of the Academy of Sciences of the Armenian SSR*, 5, 2, 71–96 [in Russian].
- 16 Pskhu, A.V. (2016). Boundary value problem for a first-order partial differential equation with a fractional discretely distributed differentiation operator. *Differential Equations*, 52, 12, 1610–1623.
- 17 Pskhu, A.V. (2016). Uravnenie drobnoi diffuzii s operatorom diskretno raspredelenного differentsirovaniia [Fractional diffusion equation with discretely distributed differentiation operator]. *Siberian Electronic Mathematical Reports*, 13, 1078–1098.
- 18 Caputo, M. (2003). Diffusion with space memory modelled with distributed order space fractional differential equations. *Annals of Geophysics*, 46, 2, 223–234.
- 19 Diethelm, K., & Ford, N. J. (2009). Numerical analysis for distributed-order differential equations. *J. of Comp. and Appl. Math*, 225, 96–104.
- 20 Jiao, Z., Chen, Y. Qu., & Podlubny, I. (2012). *Distributed-Order Dynamic Systems: Stability, Simulation, Applications and Perspectives*. London.
- 21 Ozturk, I. (1998). On the theory of fractional differential equation. *Reports of Adyghe (Circassian) International Academy of Sciences*, 3, 2, 35–39.
- 22 Naimark, M.A. (1969). *Lineinye differentsialnye operatory [Linear differential operators]*. Moscow: Nauka [in Russian].
- 23 Gadzova, L. Kh. (2015). Dirichlet and Neumann problems for a fractional ordinary differential equation with constant coefficients. *Differential Equations*, 51, 12, 1556–1562.
- 24 Gadzova, L. Kh. (2018). Boundary Value Problem for a Linear Ordinary Differential Equation with a Fractional Discretely Distributed Differentiation Operator. *Differential Equations*, 54, 2, 178–184.
- 25 Gadzova, L. Kh. (2019). Nonlocal Boundary-Value Problem for a Linear Ordinary Differential Equation with Fractional Discretely Distributed Differentiation Operator. *Mathematical Notes*, 106, 5-6, 904–908.
- 26 Gadzova, L. Kh. (2022). Obobshchennaia kraevaia zadacha dlia obyknovennogo differentsialnogo uravneniia drobnogo poriadka [Generalized boundary value problem for an ordinary differential equation of fractional order]. *Cheliabinskii fiziko-matematicheskii zhurnal – Chelyabinsk Physical and Mathematical Journal*, 7(1), 20-29. [in Russian]. <https://doi.org/10.47475/2500-0101-2022-17102>
- 27 Pskhu, A.V. (2011). Initial-value problem for a linear ordinary differential equation of noninteger order *Sbornik: Mathematics*, 202, 4, 571–582.
- 28 Wright, E.M. (1933). On the coefficients of power series having exponential singularities *J. London Math. Soc.*, 8, 29, 71–79.