

Jiahao Li¹, Jiajin Li², Gorbachev Dmitrii³, Chengru Xu⁴, Huiping Yan⁵, Yifan Lu^{6*}

^{1, 5, 6}*School of Sports Medicine and Rehabilitation, Beijing Sport University, Beijing, China;*

²*Loughborough University, Loughborough, UK;*

³*Samara State Medical University, Samara, Russia;*

⁴*Jiangsu Health Vocational College, Jiangsu, China*

^{*}*Corresponding author's e-mail: luyifan@bsu.edu.cn*

¹ORCID 0000-0002-2383-6738

²ORCID 0000-0000-0000-0000

³ORCID 0000-0002-8044-9806

⁴ORCID 0009-0002-1550-7974

⁵ORCID 0009-0005-3177-3860

⁶ORCID 0000-0002-0661-3372

The study on the relationship between different exercise modalities and physiological effects

This study aimed to systematically review the exercise intensities associated with various physical activities, synthesize the physiological effects induced by different intensity levels, and establish a dose-response reference guide for exercise modalities, thereby providing a foundation for precision-based fitness regimens. Methods: A comprehensive literature search was conducted across PubMed, Cochrane, Embase, Web of Science, EBSCO, CNKI, VIP, and Wanfang databases to identify studies examining the effects of exercise modalities on physiological indicators. The methodological quality of included randomized controlled trials (RCTs) was assessed using the Cochrane 5.1 handbook criteria. Effect sizes were pooled, and subgroup analyses were performed using Review Manager 5.4. Results: A total of 42 RCT studies were included. Low-intensity exercise modalities included ordinary walking, stair climbing, and Pilates. Moderate-intensity exercise modalities included brisk walking, Baduanjin, jogging, square dancing, Tai Chi, swimming, cycling, and dancing. High-intensity exercises included fast walking, fast running, soccer, and yoga. Low-intensity exercises improved HDL-C (d=0.06), resting heart rate (d=-2.98), and diastolic blood pressure (d=-2.93). Moderate-intensity exercises improved TG (d=-0.21), TC (d=-0.32), HDL-C (d=0.09), resting heart rate (d=-4.22), systolic blood pressure (d=-4.92), diastolic blood pressure (d=-3.51), weight (d=-2.46), and vital capacity (d=271.03). High-intensity exercises improved blood glucose (d=-0.18), systolic blood pressure (d=-3.21), and diastolic blood pressure (d=-2.58). Traditional Chinese exercises improved HDL-C (d=0.19), blood glucose (d=-1.49), vital capacity (d=285.09), systolic blood pressure (d=-9.96), and diastolic blood pressure (d=-5.68). Common exercises improved TG (d=-0.18), TC (d=-0.18), HDL-C (d=0.08), vital capacity (d=223.62), resting heart rate (d=-3.51), systolic blood pressure (d=-4.90), diastolic blood pressure (d=-2.96), weight (d=-2.04), and BMI (d=-0.74). Conclusion: Moderate-intensity exercises (e.g., brisk walking, jogging, swimming, Tai Chi) yielded more comprehensive physiological improvements compared to other intensities. Traditional Chinese exercises exhibited superior efficacy in optimizing HDL-C, glucose metabolism, respiratory function, and blood pressure regulation.

Keywords: dose-response relationship, traditional Chinese exercises, exercise modalities, RCT.

Introduction

The health benefits of physical exercise are well-established, with extensive research demonstrating its positive effects on cardiopulmonary function, muscular strength, and metabolic health. Investigating the relationship between exercise modalities and physiological responses holds significant theoretical and practical implications for precision-based fitness interventions.

The latest international Compendium of Physical Activities (CPA) [1] catalogs the metabolic equivalent of task (MET) values for over 1,000 specific activities across 21 categories. Similarly, domestic studies have quantified energy expenditure in physical activities among Chinese populations [2]. However, these efforts primarily focus on energy expenditure calibration and lack exploration of the dose-response relationships between exercise intensity and physiological outcomes.

Furthermore, emerging evidence suggests that different exercise modalities, such as traditional Chinese exercises (e.g., Tai Chi, Baduanjin, Qigong) and conventional exercises (e.g., brisk walking, swimming, Pilates), elicit distinct physiological adaptations and health benefits. Thus, synthesizing the physiological

effects of traditional Chinese exercises alongside conventional exercises is critical for optimizing exercise prescriptions.

Methods and materials

2.1 Data Sources and Search Strategy

A systematic literature search was performed across multiple databases, including PubMed, Cochrane, Embase, Web of Science, EBSCO, CNKI, VIP, and Wanfang, to identify relevant studies. In PubMed and Cochrane, Medical Subject Headings (MeSH) terms were employed, while Emtree terms were used for Embase. The specific search strategies are detailed in Table 1.

Table 1

Summary of Search Strategies

	Search Term
P (population)	young people; adolescence; young; Middle Aged [Mesh]; middle age; aged [Mesh]; elderly
I (intervention)	Dancing[Mesh]; Dance; Ballet; Square Dance; Dance, Square; Hip-Hop Dance; Dance, Hip-Hop; Hip Hop Dance; Jazz Dance; Dance, Jazz; Tap Dance; Dance, Tap; Modern Dance; Dance, Modern; Salsa Dancing; Dancing, Salsa; Line Dancing; Dancing, Line; Tai Ji[Mesh]; Tai-ji; Tai Chi; Chi, Tai; Tai Ji Quan; Ji Quan, Tai; Quan, Tai Ji; Taiji; Taijiquan; T'ai Chi; Tai Chi Chuan; Martial Arts[Mesh]; Arts, Martial; Hap Ki Do; Judo; Karate; Jujitsu; Tae Kwon Do; Aikido; Wushu; Kung Fu; Gong Fu; Fu, Gong; Gongfu; Racquet Sports[Mesh]; Racquet Sport; Sport, Racquet; Sports, Racquet; Racket Sports; Racket Sport; Sport, Racket; Sports, Racket; Squash (Sport); Racquetball; Racketball; Racket Ball; Ball, Racket; Badminton; Lacrosse; Bicycling[Mesh]; Jogging[Mesh]; Joggings; Swimming[Mesh]; Basketball[Mesh]; Basketballs; Netbal; Soccer[Mesh]; Soccers; Yoga[Mesh]; Gymnastics[Mesh]; Calisthenics; Mountaineering[Mesh]; vigorously walks; walking; brisk walking; fitness walking; walk
C (comparison)	blank control
O (outcomes)	resting heart rate; hrrest; heart rate at rest; serum total cholesterol; total cholesterol; total serum cholesterol; Lipoproteins, HDL [Mesh]; HDL Lipoproteins; Heavy Lipoproteins; Lipoproteins, Heavy; High-Density Lipoproteins; High Density Lipoproteins; Lipoproteins, High-Density; alpha-Lipoproteins; alpha Lipoproteins; alpha-1 Lipoprotein; Cholesterol, LDL [Mesh]; Low Density Lipoprotein Cholesterol; beta-Lipoprotein Cholesterol; Cholesterol, beta-Lipoprotein; beta Lipoprotein Cholesterol; LDL Cholesterol; Cholesteryl Linoleate, LDL; LDL Cholesteryl Linoleate; Vital Capacity [Mesh]; Capacities, Vital; Capacity, Vital; Vital Capacities; Forced Vital Capacity; Capacities, Forced Vital; Capacity, Forced Vital; Forced Vital Capacities; Vital Capacities, Forced; Vital Capacity, Forced; Blood Pressure [Mesh]; Pressure, Blood; Diastolic Pressure; Pressure, Diastolic; Pulse Pressure; Pressure, Pulse; Systolic Pressure; Pressure, Systolic; Pressures, Systolic; Triglycerides[Mesh]; Triacylglycerol; Triacylglycerols; Metabolic Equivalent; Metabolic Equivalents; RPE;
S (study design)	randomized controlled trial; randomized; placebo

2.2 Inclusion and Exclusion Criteria

Eligible studies met the following criteria: (1) randomized controlled trials (RCTs); (2) participants categorized as adolescents, middle-aged, or older adults; (3) interventions involving common exercise modalities; and (4) outcome measures including at least one of the following: triglyceride (TG), total cholesterol (TC), high-density lipoprotein (HDL), low-density lipoprotein (LDL), resting heart rate, vital capacity (VC), systolic blood pressure (SBP), diastolic blood pressure (DBP), rating of perceived exertion (RPE), body weight, or body mass index (BMI). Studies were excluded if they were duplicate publications, literature reviews, letters to the editor, conference abstracts, or animal model research. Additionally, articles lacking full-text availability or original data were excluded.

2.3 Data Extraction

Two independent researchers extracted all data and compiled the results. Discrepancies were resolved through consultation with a third reviewer. Extracted data included: (1) primary author and publication year; (2) baseline characteristics (age and sample size of experimental and control groups); (3) exercise parameters (type, frequency, duration, and intensity); and (4) reported outcomes. The risk of bias in included studies was

assessed using the Cochrane Risk of Bias Tool, which evaluates seven domains: (1) random sequence generation, (2) allocation concealment, (3) blinding of participants and personnel, (4) blinding of outcome assessment, (5) incomplete outcome data, (6) selective reporting, and (7) other potential biases.

2.4 Data Analysis

Quantitative synthesis and subgroup analyses were performed using Review Manager software (version 5.4). Effect sizes were pooled using a random-effects model. Heterogeneity among studies was assessed using the I^2 statistic and Cochran's Q test, with $I^2 > 50\%$ or a Q test P-value ≤ 0.10 indicating significant heterogeneity. Publication bias was evaluated via funnel plot inspection. Sensitivity analyses were conducted by excluding trials with a high risk of bias to test the robustness of the pooled results.

Results and Discussion

2 Results

2.1 Search Results

A total of 4,846 potentially eligible articles were initially identified through database searches. After a systematic screening process—including removal of duplicates, title/abstract screening, and full-text assessment—42 studies met the predefined inclusion criteria and were included in the final analysis.




Figure 1. Flowchart of Literature Selection

2.2 Basic Characteristics of Included Literature

The basic characteristics of the included literature are shown in Table 2. The quality evaluation is presented in Figure 2.

Table 2

Basic Characteristics of Included Literature

Study	Sample Size	Gender	Age	Mode	Intensity	Inflammatory outcome
Guaxia Hu 2014 [3]	230	Male/Female	53~70	Baduanjin	100beats/min	LC, SBP, DBP
A RCOOPER 2000 [4]	176	Male/Female	46-49	walk	4~5mets	SBP, DBP, Body weight
Abdelbasset WK 2020 [5]	31	Male/Female	54.9±4.7	bicycle	50~70 %HRmax	TG, TC, LDL-C, HDL-C
Asikainen, T.M 2003 [6]	385	Female	55 (3.7)	walk	45 %VO2max	TC/LDL-C/HDL-C/Glucose/SBP/DBP
Baker, T.T 1986 [7]	34	Male	58.2	running	65 %~85 %HRmax	TC, LDL-C, HDL-C, Body weight
Buttelli 2021 [8]	61	Male	60-75	Pilates	1.5~2met	TG, TC, LDL-C, HDL-C, Glucose
F.M. Finucane 2010 [9]	100	Male/Female	67.4-76.3	Bicycle	50~70 %HRmax	HDL-C, Glucose, SBP, DBP, Body weight
Fulin Lu 2018 [10]	96	Female	57.2 ±10.1.	Dance	Heart rate=170-age (>60age) Heart rate =160-age (<60age)	TG, TC, LDL-C, HDL-C, Glucose, RHR, SBP, DBP, WHR, BMI
Haiping Zhang 2016 [11]	40	Female	62.85 ±1.66	walk	50 %~60 %VO2max	TG, TC, LDL-C, HDL-C, LC, VO2max
Hanxiao Zhu 2009 [12]	135	Male	61-65	walk	30 %~45 %F.C	LC, RHR, SBP, DBP, Body weight, BMI
JE Donnelly 2000 [13]	11	Female	45-60	walk	60~75 %HRmax	TC, HDL-C, Glucose, RHR, SBP, DBP
Koh, Y 2018 [14]	27	Male/Female	18-65	running	70 %HRmax	TG, TC, LDL-C, HDL-C, Body weight
Krustrup P 2013 [15]	66	Male	31~54	Soccer	85 % ± 7 %HRmax	LDL-C, HDL-C, Glucose, RHR
Krustrup, P 2009 [16]	47	Male	20-43	running/Soccer	82 %HRmax	TC, LDL-C, HDL-C, RHR, SBP, DBP, Body weight
Kuo, M. C. 2018 [17]	36	Male/Female	>65	walk	13~15 RPE	WHR
Lee SH 2019 [18]	20	Female	70 ± 4	taekwondo	30~60 %HRR	RHR, SBP, DBP, Body weight
LI He 2018 [19]	88	Male/Female	58±2	walk	45~50 %HRmax	RHR, SBP, DBP, Body weight
Lian, X. Q 2014 [20]	330	Male/Female	40~78	walk	3~6METs	TG, TC, LDL-C, HDL-C
Lorenzo A 2008 [21]	288	Male/Female	63.5±1	Dance/Yoga	70 %HRmax	TG, TC, Glucose
Mao, H. N 2006 [22]	62	Male/Female	45~72	Tai Chi	50~60 %VO2max	SBP, DBP
Marti, B 1990 [23]	61	Male	38.8 + 8.9	running	85 %HRmax	TG, TC, LDL-C, HDL-C, RHR, WHR
Mazurek, Krzysztof 2014 [24]	64	Male/Female	19.5±0.6	bicycle	65 %~75 % HRmax	TG, TC, LDL-C, HDL-C, WHR
Miao Sun 2018 [25]	60	Female	55-70	Dance	110~130beats/min	TG, TC, LDL-C, HDL-C, Glucose, LC, RHR, SBP, DBP, Body weight, WHR, BMI
Min Jeong 2018 [26]	18	Male/Female	20.8±1.9	walk/ Climb the stairs	50~60 %HRmax	RHR/SBP/DBP/ Body weight
Miyaki A 2012 [27]	22	Female	52~77	walk/bicycle	60~75 %HRmax	TG, TC, LDL-C, HDL-C, RHR, SBP, DBP, Body weight
Papp 2016 [28]	44	Male/Female	20~37	Yoga	RPE 14~17	RHR, SBP, DBP, Body weight, WHR
Philippe 2018 [29]	115	Female	44±11.5	Walk	4~10 RPE	SBP, DBP, Body weight
Rodrigues-Krause 2018 [30]	30	Female	60~70	Dance	60 % VO2peak	TG, TC, LDL-C, HDL-C, Glucose, Body weight
Shanwei Xue 2013 [31]	40	Male/Female	66.96±13.2	Baduanjin	170-age (>60age); 180-age (<60age)	LC, RHR
Songtao Wang 2005 [32]	99	Male	61-65	Walk	30 %~ 45 %F.C	TG, TC, LDL-C, HDL-C, Body weight, BMI
Tanaka 1997 [33]	18	Male/Female	44~50	Swimming	60 %HRmax	RHR, SBP, DBP
Tsung-Lin 2019 [34]	41	Male/Female	19.72±0.8	Walk	RPE 8	TG, HDL-C, RHR, SBP, DBP, Body weight
Vasconcellos 2016 [35]	30	Male/Female	12.8~15.4	Soccer	80.4 %~88.6 %HRmax	TG, TC, LDL-C, HDL-C, SBP, DBP, Body weight
Vasconcellos 2021 [36]	13	Male/Female	12.3~15.5	Soccer	84.5 %HRmax	TG, HDL-C, Glucose, SBP, DBP, Body weight
Woolf-May 1999 [37]	237	Male/Female	40~66	Walk	68.6 %~78.2 %HRmax	TC, LDL-C, HDL-C
Woolf-May 2000 [38]	97	Male/Female	40~69	Walk	75 %~80 %HRmax	TC, LDL-C, HDL-C
Xiao-Ling 2019 [39]	198	Male/Female	45.5~58.4	Tai Chi	70 %~80 %HRmax	TG, TC, LDL-C, HDL-C, SBP, DBP
Xiaoxia Li 2012 [40]	92	Male/Female	50~65	Walk	40 %~65 %F.C.	TG, TC, LDL-C, HDL-C
Xiaoying Han 2010 [41]	236	Male/Female	45~75	Tai Chi	70%~80%HRmax	SBP, DBP
Xijun Xiao 2005 [42]	120	Male/Female	≥40	Running	40 %~50 %HRR	TG, TC, HDL-C, SBP, DBP, Body weight, WHR, BMI
Yaping Bai 2019 [43]	133	Male/Female	45-75	Dance	60 %~80 %HRmax	TG, TC, LDL-C, HDL-C, SBP, DBP, BMI
zhang 2008 [44]	20	Female	51.2-63.6	Tai Chi	62 %~72.2 %HRmax	TG, TC, LDL-C, HDL-C, Glucose

LC : 肺活量, RHR, WHR : waist hip rate




Figure 2. Quality Evaluation Chart

2.3 Dose-Response Relationship of Exercise Intensity

2.3.1 Classification of Exercise Modalities by Intensity

The exercise interventions from the included studies were categorized into three intensity levels based on the American College of Sports Medicine (ACSM) guidelines [45] (Table 3).

Table 3

Classification of Exercise Modalities by Intensity

Intensity	Exercise Modalities
Low	Casual walking, Stair climbing, Pilates
Moderate	Walk, Ba duan jin, Jogging, Dance, Tai Chi, Swimming, Bicycle, Dance, Yoga, Taekwondo
High	Power walking, Running, Soccer, Yoga

2.3.2 Physiological Effects of Different Exercise Intensities

The physiological benefits of different exercise intensities are shown in Figure 3. Low-intensity exercise improved HDL-C ($d=0.06$, 95 % CI: 0.02–0.10; $P<0.05$), resting heart rate ($d=-2.98$, 95 % CI: -4.77 to -1.19; $P<0.05$), and diastolic blood pressure ($d=-2.93$, 95 % CI: -5.50 to -0.36; $P<0.05$). Moderate-intensity exercise improved TG ($d=-0.21$, 95 % CI: -3.5 to -0.07; $P<0.05$), TC ($d=-0.32$, 95 % CI: -0.52 to -0.12; $P<0.05$), HDL-C ($d=0.09$, 95 % CI: -0.02 to -0.15; $P<0.05$), resting heart rate ($d=-4.22$, 95 % CI: -5.56 to -2.88; $P<0.05$), systolic blood pressure ($d=-4.92$, 95 % CI: -8.59 to -1.24; $P<0.05$), diastolic blood pressure ($d=-3.51$, 95 % CI: -5.33 to -1.68; $P<0.05$), body weight ($d=-2.46$, 95 % CI: -3.79 to -1.16; $P<0.05$), and vital capacity ($d=271.03$, 95 % CI: 119.35–422.71; $P<0.05$). High-intensity exercise improved blood glucose ($d=-0.18$, 95 % CI: -0.27 to -0.08; $P<0.05$), SBP ($d=-3.21$, 95 % CI: -6.07 to -0.34; $P<0.05$), and DBP ($d=-2.58$, 95 % CI: -4.31 to -0.85; $P<0.05$).




Figure 3. Forest Plot of Physiological Effects by Exercise Intensity

2.4 Dose-Response Relationships of Different Exercise Types

2.4.1 Exercise Modalities Under Different Exercise Types

The specific exercise modalities corresponding to different exercise types are presented in Table 4 below.

Table 4

Exercise Types and Corresponding Modalities

Exercise Type	Exercise Modalities
Traditional Chinese Exercises	Tai Chi, Ba Duan Jin
Popular Exercises	Casual walking, Stair climbing, PilatesWalk, Jogging, Dance, Swimming, Bicycle, Dance, Yoga, TaekwondoPower walking, Running, Soccer, Yoga

2.4.2 Dose-Response Relationships of Different Exercise Types

As shown in Figure 4, traditional Chinese exercises significantly improved HDL-C ($d=0.19$, 95 %CI: 0.10–0.29, $P<0.05$), blood glucose ($d=-1.49$, 95 %CI: -2.83 to -0.15, $P<0.05$), vital capacity ($d=285.09$, 95 %CI: 84.05–486.13, $P<0.05$), systolic blood pressure ($d=-9.96$, 95 %CI: -15.85 to -4.08, $P<0.05$), and diastolic blood pressure ($d=-5.68$, 95 %CI: -7.22 to -4.13, $P<0.05$). Common exercises demonstrated benefits in improving TG ($d=-0.18$, 95 %CI: -0.27 to -0.09, $P<0.05$), TC ($d=-0.18$, 95 %CI: -0.34 to -0.01, $P<0.05$), HDL-C ($d=0.08$, 95 %CI: 0.01–0.15, $P<0.05$), vital capacity ($d=223.62$, 95 %CI: 40.97–406.28, $P<0.05$), resting heart rate ($d=-3.51$, 95 %CI: -4.92 to -2.10, $P<0.05$), systolic blood pressure ($d=-4.90$, 95 %CI: -7.05 to -2.75, $P<0.05$), diastolic blood pressure ($d=-2.96$, 95 %CI: -4.53 to -1.40, $P<0.05$), body weight ($d=-2.04$, 95 %CI: -3.36 to -0.72, $P<0.05$), and BMI ($d=-0.74$, 95 %CI: -1.33 to -0.15, $P<0.05$).




Figure 4. Forest Plot of Physiological Effects by Exercise Type

4 Discussion

4.1 Dose-Response Relationship of Exercise Intensity

The present study demonstrates that body weight improvement was observed only with moderate-intensity exercise, while no significant effects were found for low- or high-intensity interventions. Previous research has established a dose-response relationship between aerobic exercise volume and visceral fat reduction, with a minimum requirement of 10 METs·h/week for significant visceral fat loss [46–48]. The American College of Sports Medicine (ACSM) recommends moderate-intensity physical activity for at least 150 minutes/week for overweight/obese individuals to achieve weight or body fat reduction, supporting moderate intensity as the optimal exercise intensity for body weight management.

Regarding lipid metabolism, our findings show that low-intensity exercise improved HDL-C levels, moderate-intensity exercise benefited TG, TC, and HDL-C, while high-intensity exercise showed no significant lipid-modifying effects. The HDL-C improvement with low-intensity exercise may be mediated through

enhanced insulin sensitivity and reduced body fat content [49]. Sunami et al. [50] reported significant HDL-C elevation following 5 months of moderate-intensity cycling training. Ye Guohong et al. [51] proposed that prolonged moderate-intensity endurance exercise reduces plasma TG levels by increasing lipoprotein lipase activity, promoting fatty acid utilization in skeletal muscle and hepatic TG export. Moderate-intensity exercise may also improve TC levels by enhancing cholesterol ester transfer protein (CETP) activity, which facilitates cholesterol exchange between HDL-C and LDL-C [52]. Collectively, the lipid-lowering effects of exercise appear to diminish with increasing intensity, with small-to-moderate intensity aerobic exercise demonstrating superior lipid-modulating effects compared to high-intensity exercise [32].

High-intensity exercise significantly reduced blood glucose levels, likely through improved insulin sensitivity in muscle cells and accelerated glycogen synthesis in muscle and liver tissues [53]. Blood pressure improvements were observed across all three intensity levels, consistent with aerobic exercise's established benefits for vascular function, arterial stiffness reduction in hypertensive patients [54, 55], and decreased risk of acute cardiovascular events [56].

Both low- and moderate-intensity exercises reduced resting heart rate, potentially through enhanced parasympathetic nervous system activity [57]. Moderate-intensity exercise uniquely improved vital capacity, attributable to increased tidal volume during exercise and long-term adaptations in respiratory muscle function and lung-thorax elasticity [58].

4.2 Dose-Response Relationship of Different Exercise Modalities

Traditional Chinese exercises (e.g., Tai Chi, Baduanjin) emphasize fluid and continuous movements with mind-body integration, focusing on breath-movement coordination and mental concentration [59]. These exercises typically feature slow tempo and gentle motions but require high neuromuscular control [60]. In contrast, conventional exercises (e.g., running, swimming) prioritize cardiorespiratory endurance and muscular strength through higher-intensity movements with greater physical demands [61].

Regarding antioxidant and anti-inflammatory effects, traditional Chinese exercises reduce oxidative stress and inflammation through slow, rhythmic movements combined with deep breathing, potentially improving HDL-C and blood glucose levels more effectively [62]. Conventional exercises also exhibit antioxidant properties (e.g., running and swimming can increase HDL-C), though high-intensity exercise may transiently elevate oxidative stress [63].

For neuromodulation and cardiovascular health, traditional Chinese exercises improve vascular endothelial function by balancing the autonomic nervous system (reducing sympathetic activity while enhancing parasympathetic tone), thereby lowering blood pressure and improving cardiovascular outcomes [64]. Conventional exercises enhance cardiac function and vascular elasticity to improve blood pressure and cardiovascular health, though acute sympathetic activation may attenuate blood pressure reduction [65].

In terms of energy metabolism and body weight management, traditional Chinese exercises (with lower intensity) show limited direct effects on body weight and BMI [66], whereas conventional exercises effectively manage body weight and BMI through higher energy expenditure [67].

Conclusions

5.1 Low-intensity exercises include regular walking, stair climbing, and Pilates; Moderate-intensity exercises include brisk walking, Baduanjin, jogging, square dancing, Tai Chi, swimming, cycling, and dance; High-intensity exercises include power walking, running, soccer, and yoga.

5.2 Low-intensity exercise can improve HDL-C, resting heart rate, and diastolic blood pressure; moderate-intensity exercise can improve TG, TC, HDL-C, resting heart rate, systolic blood pressure, diastolic blood pressure, body weight, and vital capacity; high-intensity exercise can improve blood glucose, systolic blood pressure, and diastolic blood pressure.

5.3 Traditional Chinese exercises can improve HDL-C, vital capacity, blood glucose, systolic blood pressure, and diastolic blood pressure. Common exercises can improve TG, TC, HDL-C, vital capacity, resting heart rate, systolic blood pressure, diastolic blood pressure, body weight, and BMI.

Funding

This research was funded by the National key research and development program of China (2020YFC2002902).

References

- 1 Ainsworth, B.E., Haskell, W.L., Herrmann, S.D., et al. (2011). 2011 Compendium of Physical Activities: a second update of codes and MET values. *Medicine and science in sports and exercise*, 43(8), 1575–1581.
- 2 Junqiang, Q., Junchao, Y., Mingyue, L., et al. (2022). Compilation of Physical Activities of Healthy Chinese Adults: Reference Values for Energy Expenditure [J]. *Chinese Journal of Sports Medicine*, 41(05), 335–349.
- 3 Guangxia, H., & Keping, G. (2014). The Impact of Health-Qigong Eight-Section Exercises on the Quality of Life of Elderly People in a Community in Jilin Province. *Medicine and Society*, 27(5), 74–76.
- 4 Cooper, A.R., Moore, L.A., Mckenna, J., et al. (2000). What is the magnitude of blood pressure response to a programme of moderate intensity exercise? Randomised controlled trial among sedentary adults with unmedicated hypertension. *British journal of general practice*, 50(461), 958–962.
- 5 Abdelbasset, W.K., Elsayed, S.H., Nambi, G., et al. (2020). Effect of Moderate-Intensity Aerobic Exercise on Hepatic Fat Content and Visceral Lipids in Hepatic Patients with Diabesity: a Single-Blinded Randomised Controlled Trial. *Evidence-based complementary and alternative medicine*, 1923575.
- 6 Asikainen, T.M., Miilunpalo, S., Kukkonen-harjula, K., et al. (2003). Walking trials in postmenopausal women: effect of low doses of exercise and exercise fractionization on coronary risk factors. *Scandinavian journal of medicine & science in sports*, 13(5), 284–292.
- 7 Baker, T.T., Allen, D., Lei, K.Y., et al. (1986). Alterations in lipid and protein profiles of plasma lipoproteins in middle-aged men consequent to an aerobic exercise program. *Metabolism: clinical and experimental*, 35(11), 1037–1043.
- 8 Buttelli, A.C.K., Costa, R.R., Farinha, J.B., et al. (2021). Pilates training improves aerobic capacity, but not lipid or lipoprotein levels in elderly women with dyslipidemia: A controlled trial. *Journal of Bodywork and Movement Therapies*, 26, 227–232.
- 9 Finucane, F.M., Sharp, S.J., Purslow, L.R., et al. (2010). The effects of aerobic exercise on metabolic risk, insulin sensitivity and intrahepatic lipid in healthy older people from the Hertfordshire Cohort Study: a randomised controlled trial. *Diabetologia*, 53(4), 624–631.
- 10 Fulin, L., Shuyong, M., & Qi, Y. (2018). Effect of Square Dance Intervention on Type 2 Diabetes Mellitus in Middle-aged and Elderly Women. *International Journal of Geriatrics*, 39(3), 119–122.
- 11 Haiping, Z., Yixin, W., & Feng, Y. (2016). The impact of brisk walking on the cardio-pulmonary endurance and lipid-related indicators of elderly women. *Chinese Journal of Gerontology*, 36(14), 3514–3516.
- 12 Hanxiao, Z. (2009). Study on advisable exercise intensity for elder men. *Modern Preventive Medicine*, 36(1), 107–109, 116.
- 13 Donnelly, J.E., Jacobsen, D.J., Heelan, K.S., et al. (2000). The effects of 18 months of intermittent vs. continuous exercise on aerobic capacity, body weight and composition, and metabolic fitness in previously sedentary, moderately obese females. *International journal of obesity and related metabolic disorders*, 24(5), 566–572.
- 14 Koh, Y., Park, J., & Carter, R. (2018). Oxidized Low-Density Lipoprotein and Cell Adhesion Molecules Following Exercise Training. *International journal of sports medicine*, 39(2), 83–88.
- 15 Krustup, P., Randers, M.B., Andersen, L.J., et al. (2013). Soccer Improves Fitness and Attenuates Cardiovascular Risk Factors in Hypertensive Men. *Medicine and Science in Sports and Exercise*, 45(3), 553–560.
- 16 Krustup, P., Nielsen, J.J., Krustup, B.R., et al. (2009). Recreational soccer is an effective health-promoting activity for untrained men. *British journal of sports medicine*, 43(11), 825–831.
- 17 Kuo, M.C., Chen, C.M., & Jeng, C. (2018). A Randomized Controlled Trial of the Prescribed Stepper Walking Program in Preventing Frailty Among the Dwelling Elderly: application of Comprehensive Geriatric Assessment. *Topics in geriatric rehabilitation*, 34(3), 223–33.
- 18 Lee, S.H., Scott, S.D., Pekas, E.J., et al. (2019). Taekwondo training reduces blood catecholamine levels and arterial stiffness in postmenopausal women with stage-2 hypertension: randomized clinical trial. *Clinical and experimental hypertension*, 41(7), 675–681.
- 19 He, L., Wei, W.R., & Can, Z. (2018). Effects of 12-week brisk walking training on exercise blood pressure in elderly patients with essential hypertension: a pilot study. *Clinical and experimental hypertension*, 40(7), 673–679.
- 20 Lian, X.Q., Zhao, D., Zhu, M., et al. (2014). The influence of regular walking at different times of day on blood lipids and inflammatory markers in sedentary patients with coronary artery disease. *Preventive medicine*, 58, 64–69.
- 21 Gordon, L.A., Morrison, E.Y., Mcgrowder, D.A., et al. (2008). Effect of exercise therapy on lipid profile and oxidative stress indicators in patients with type 2 diabetes. *BMC complementary and alternative medicine*, 8, 21.
- 22 Mao, H.N., & Sha, P. (2006). Effect of Tai Chi exercise on blood pressure, plasma nitrogen monoxidum and endothelin in hypertensive patients. *Chinese journal of clinical rehabilitation*, 10(48), 65–67.
- 23 Marti, B., Suter, E., Riesen, W.F., et al. (1990). Effects of long-term, self-monitored exercise on the serum lipoprotein and apolipoprotein profile in middle-aged men. *Atherosclerosis*, 81(1), 19–31.
- 24 Mazurek, K., Krawczyk, K., Zmijewski, P., et al. (2014). Effects of aerobic interval training versus continuous moderate exercise programme on aerobic and anaerobic capacity, somatic features and blood lipid profile in c llegate females. *Annals of Agricultural and Environmental Medicine*, 21(4), 844–849.
- 25 Yan, S., Qingzhi, Z., Dan, Z., et al. (2018). The impact of square dancing exercise on the physical and mental health of middle-aged and elderly women. *Chinese Journal of Applied Physiology*, 34(03), 246–248.

- 26 Cho, M.J., Park, Y.R., Bunsawat, K., et al. (2018). Comparison of the effects of short-term stair climbing and walking exercise on vascular function in healthy young adults. *IJASS (International Journal of Applied Sports Sciences)*, 30(2), 125–33.
- 27 Miyaki, A., Maeda, S., Choi, Y., et al. (2012). Habitual aerobic exercise increases plasma pentraxin 3 levels in middle-aged and elderly women. *Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme*, 37(5), 907–911.
- 28 Papp, M.E., Lindfors, P., Nygren-Bonnier, M., et al. (2016). Effects of High-Intensity Hatha Yoga on Cardiovascular Fitness, Adipocytokines, and Apolipoproteins in Healthy Students: a Randomized Controlled Study. *Journal of alternative and complementary medicine (New York, NY)*, 22(1), 81–87.
- 29 Gradidge, P.J., & Golele, P.N. (2018). Walking as a feasible means of effecting positive changes in BMI, waist, and blood pressure in black South African women. *African historical studies*, 18(4), 917–21.
- 30 Rodrigues-Krause, J., Farinha, J.B., Ramis, T.R., et al. (2018). Effects of dancing compared to walking on cardiovascular risk and functional capacity of older women: a randomized controlled trial. *Experimental gerontology*, 114, 67–77.
- 31 Weishan, X. (2013). The effect of Health Qigong Ba Duan Jin on cardiopulmonary function of aged people. *Journal of Bohai University*, 34(4), 431–434.
- 32 Songtao, W., Anli, W., Zhengzhen, W., et al. (1997). A comparison of the effects of different-intensity brisk walking exercises on body composition and blood lipids in elderly men. *Chinese Journal of Sports Medicine*, 24(5), 599–601.
- 33 Tanaka, H., Bassett, D.R., Howley, E.T., et al. (1997). Swimming training lowers the resting blood pressure in individuals with hypertension. *Journal of hypertension*, 15(6), 651–657.
- 34 Chiang, T.L., Chen, C., Hsu, C.H., et al. (2019). Is the goal of 12,000 steps per day sufficient for improving body composition and metabolic syndrome? The necessity of combining exercise intensity: a randomized controlled trial. *BMC public health*, 19(1), 1215.
- 35 Vasconcellos, F., Seabra, A., Cunha, F., et al. (2016). Health markers in obese adolescents improved by a 12-week recreational soccer program: a randomised controlled trial. *Journal of Sports Sciences*, 34(6), 564–575.
- 36 Vasconcellos, F., Cunha, F.A., Gonet, D.T., et al. (2021). Does Recreational Soccer Change Metabolic Syndrome Status in Obese Adolescents? A Pilot Study. *Research Quarterly for Exercise and Sport*, 92(1), 91–99.
- 37 Woolf-May, K., Kearney, E.M., Owen, A., et al. (1999). The efficacy of accumulated short bouts versus single daily bouts of brisk walking in improving aerobic fitness and blood lipid profiles. *Health education research*, 14(6), 803–815.
- 38 Woolf-May, K., Jones, W., Kearney, E.M., et al. (2000). Factor XIIIa and triacylglycerol rich lipoproteins: responses to exercise intervention. *British journal of sports medicine*, 34(4), 289–292.
- 39 Shou, X.L., Wang, L., Jin, X.Q., et al. (2019). Effect of T'ai Chi Exercise on Hypertension in Young and Middle-Aged In-Service Staff. *Journal of alternative and complementary medicine (New York, NY)*, 25(1), 73–78.
- 40 Xiaoxia, L., Ming, J., Yong, W., et al. (2012). Effects of walking exercise on lipids of elderly obese people and its correlation with adiponectin gene polymorphisms. *Journal of Shandong Institute of Physical Education*, 28(06), 52–58.
- 41 Qiaoying, H., Xiufeng, H., Lei, L., et al. (2010). The effect of shadow boxing exercise on the long-term quality of life in middle-aged and elderly patients with primary hypertension. *Chinese Journal of Modern Nursing*, 16(14), 1617–1619.
- 42 Xijun, X. (2005). Effect of different exercise intensities to grade-1 hypertensives. *Chinese Journal of Rehabilitation Medicine*, 20(5), 349–352.
- 43 Yaping, B., Ying, L., Dongmei, W., et al. (2019). The impact of community square dancing on blood pressure and lipid levels of middle-aged and elderly patients with hypertension. *Chinese Journal of Hypertension*, 27(05), 474–478.
- 44 Zhang, Y., & Fu, F.H. (2008). Effects of 14-week Tai Ji Quan exercise on metabolic control in women with type 2 diabetes. *The American journal of Chinese medicine*, 36(4), 647–654.
- 45 Thompson, P.D., Arena, R., Riebe, D., et al. (2013). ACSM's new preparticipation health screening recommendations from ACSM's guidelines for exercise testing and prescription, ninth edition. *Current sports medicine reports*, 12(4), 215–217.
- 46 Miyatake, N., Nishikawa, H., Morishita, A., et al. (2002). Daily walking reduces visceral adipose tissue areas and improves insulin resistance in Japanese obese subjects. *Diabetes research and clinical practice*, 58(2), 101–107.
- 47 Halverstadt, A., Phares, D.A., Ferrell, R.E., et al. (2003). High-density lipoprotein-cholesterol, its subfractions, and responses to exercise training are dependent on endothelial lipase genotype. *Metabolism: clinical and experimental*, 52(11), 1505–1511.
- 48 Wilund, K.R., Ferrell, R.E., Phares, D.A., et al. (2002). Changes in high-density lipoprotein-cholesterol subfractions with exercise training may be dependent on cholesteryl ester transfer protein (CETP) genotype. *Metabolism: clinical and experimental*, 51(6), 774–778.
- 49 Kodama, S., Tanaka, S., Saito, K., et al. (2007). Effect of aerobic exercise training on serum levels of high-density lipoprotein cholesterol: a meta-analysis. *Arch Intern Med*, 167(10), 999–1008.
- 50 Sunami, Y., Motoyama, M., Kinoshita, F., et al. (1999). Effects of low-intensity aerobic training on the high-density lipoprotein cholesterol concentration in healthy elderly subjects. *Metabolism*, 48(8), 984–988.
- 51 Guohong, Y., Zhengyi, W., Jiansheng, X., et al. (2004). The Development of Exercise and Serum Lipids. *Journal of Beijing Sport University*, 27(7), 933–935.
- 52 Durstine, J.L., Grandjean, P.W., Davis, P.G., et al. (2001). Blood lipid and lipoprotein adaptations to exercise: a quantitative analysis. *Sports Med*, 31(15), 1033–1062.

- 53 Holloszy, J.O. (2005). Exercise-induced increase in muscle insulin sensitivity. *Journal of applied physiology (Bethesda, Md: 1985)*, 99(1), 338–343.
- 54 Woolley, B., Stoner, L., Lark, S., et al. (2015). Effect of early exercise engagement on arterial stiffness in patients diagnosed with a transient ischaemic attack. *Journal of human hypertension*, 29(2), 87–91.
- 55 Collier, S.R., Sandberg, K., Moody, A.M., et al. (2015). Reduction of plasma aldosterone and arterial stiffness in obese pre- and stage I hypertensive subjects after aerobic exercise. *Journal of human hypertension*, 29(1), 53–57.
- 56 Goldberg, M.J., Boutcher, S.H., & Boutcher, Y.N. (2012). The effect of 4 weeks of aerobic exercise on vascular and baroreflex function of young men with a family history of hypertension. *Journal of human hypertension*, 26(11), 644–649.
- 57 Nunan, D., Sandercock, G.R., & Brodie, D.A. (2010). A quantitative systematic review of normal values for short-term heart rate variability in healthy adults. *Pacing Clin Electrophysiol*, 33(11), 1407–1417.
- 58 Yuhong, X., & Yiming, B. (2017). The impact of moderate-intensity aerobic exercise on the cardio-pulmonary functions of minority college students in Guizhou. *Chinese Journal of School Health*, 38(09), 1424–1426.
- 59 Lan, C., Chen, S.Y., Lai, J.S., et al. (1999). The effect of Tai Chi on cardiorespiratory function in patients with coronary artery bypass surgery. *Medicine and science in sports and exercise*, 31(5), 634–638.
- 60 Hong, Y., & Li, J.X. (2007). Biomechanics of Tai Chi: a review. *Sports Biomech*, 6(3), 453–464.
- 61 Oja, P., & Titze, S. (2011). Physical activity recommendations for public health: development and policy context. *Epma j*, 2(3), 253–259.
- 62 Wang, C., Bannuru, R., Ramel, J., et al. (2010). Tai Chi on psychological well-being: systematic review and meta-analysis. *BMC Complement Altern Med*, 10, 23.
- 63 Fatouros, I.G., Jamurtas, A.Z., Villiotou, V., et al. (2004). Oxidative stress responses in older men during endurance training and detraining. *Medicine and science in sports and exercise*, 36(12), 2065–2072.
- 64 Taylor-Piliae, R.E., & Haskell, W.L. (2007). Tai Chi exercise and stroke rehabilitation. *Top Stroke Rehabil*, 14(4), 9–22.
- 65 Maron, B.J., Thompson, P.D., Ackerman, M.J., et al. (2007). Recommendations and considerations related to preparticipation screening for cardiovascular abnormalities in competitive athletes: 2007 update: a scientific statement from the American Heart Association Council on Nutrition, Physical Activity, and Metabolism: endorsed by the American College of Cardiology Foundation. *Circulation*, 115(12), 1643–1655.
- 66 Tsai, J.C., Wang, W.H., Chan, P., et al. (2003). The beneficial effects of Tai Chi Chuan on blood pressure and lipid profile and anxiety status in a randomized controlled trial. *Journal of alternative and complementary medicine (New York, NY)*, 9(5), 747–754.
- 67 Penedo, F.J., & Dahn, J.R. (2005). Exercise and well-being: a review of mental and physical health benefits associated with physical activity. *Curr Opin Psychiatry*, 18(2), 189–193.

Information about authors

Jiahao Li — School of Sports Medicine and Rehabilitation, Beijing Sport University, Beijing, China; ORCID ID: 0000-0002-2383-6738

Jiajin Li — Loughborough University, Loughborough, UK; ORCID ID: 0000-0000-0000-0000

Gorbachev Dmitrii — Doctor of Medical Sciences, Associate Professor, Samara state medical university, Samara, Russia; e-mail: d.o.gorbachev@samsmu.ru; ORCID ID: 0000-0002-8044-9806

Chengru Xu — Jiangsu Health Vocational College, Jiangsu, China; ORCID ID: 0009-0002-1550-7974

Huiping Yan — School of Sports Medicine and Rehabilitation, Beijing Sport University, Beijing, China; ORCID ID: 0009-0005-3177-3860

Yifan Lu (contact person) — Professor, School of Sports Medicine and Rehabilitation, Beijing Sport University, Beijing, China; e-mail: luyifan@bsu.edu.cn; ORCID ID: 0000-0002-0661-3372