

A.A. Bobodzhanov¹, B.T. Kalimbetov², V.F. Safonov¹

¹National Research University «Moscow Power Engineering Institute», Russia;
²Kh.A. Yassawi International Kazakh-Turkish University, Turkestan, Kazakhstan
(E-mail: bobojanova@yandex.ru)

Singularly perturbed control problems in the case of the stability of the spectrum of the matrix of an optimal system

The paper considers a singularly perturbed control problem with a quadratic quality functional. Such problems in their standard formulation under known spectrum restrictions (the points of the spectrum of the optimal system are not purely imaginary and are located symmetrically with respect to the imaginary axis) were previously considered using the Vasilyeva - Butuzov method of boundary functions. If at least one of the points of the spectrum for some values of the independent variable falls on the imaginary axis, the boundary functions method does not work. It is precisely this situation with the assumption of purely imaginary points of the spectrum that is investigated in this paper. In this case, you have to develop a different approach based on the ideas of the regularization method S.A. Lomov. It should also be noted that in the control problems considered earlier, the cost functional either did not depend on a small parameter at all, or allowed a smooth dependence on the parameter. In this paper, an irregular dependence on a small parameter is allowed, in particular, the presence in them of a rapidly changing damping function in the form of an exponential factor under the integral sign. In this case, the spectrum behavior of the optimal system depends on the damping coefficient, which (under certain conditions) can shift the spectrum in one direction or another in the complex plane. In this case, a situation may arise when some points of the spectrum for individual values (or even on a certain continuum set) of an independent variable can become purely imaginary. This situation is not amenable to investigation by the previously mentioned Vasilyeva - Butuzov method of boundary functions. However, it can be fully studied using the regularization method S.A. Lomov, the algorithm of which is applied to the considered control problem in the present paper. The presentation of this method begins with a brief description of the maximum principle of L.S. Pontryagin for the classical optimal control problem, which then, along with other ideas, is used to justify the results in the considered control problem.

Keywords: singularly perturbed, Pontryagin's maximum principle, regularization, asymptotic convergence.

Introduction

The presentation of the regularization method of S.A. Lomov [1] begins with a brief description of the Pontryagin's maximum principle for the classical optimal control problem, which is then applied to a linear singularly perturbed control problem with a quadratic quality functional (cost functional) in the case of a stable spectrum of the matrix of an optimal system.

Consider a linear control system

$$\frac{dx}{dt} = A(t)x + B(t)u + h(t), \quad x(0) = x^0; \quad (1)$$

$$J(u) = \frac{1}{2} \int_0^T (x^* Q(t)x + u^* R(t)u) dt \rightarrow \min, \quad (2)$$

where $x(t)$, $h(t)$ are n – dimensional; $u(t)$ is m – dimensional vector functions, x^0 is a constant n -dimensional vector; $A(t)$ is $(n \times n)$ –matrix; $B(t)$ is $(n \times m)$ – matrix; $Q(t)$ is a symmetric non-negative definite $(n \times n)$ –matrix, $R(t)$ is a symmetric positive definite $(m \times m)$ – matrix, $*$ is a transposition sign. It is required to transfer the system (1) from a given initial position $x(0) = x^0$ to a position $x(T)$ in a fixed time $T < +\infty$ ($x(T)$ is not fixed) so that the functional $J(u)$ takes the minimum value. A similar problem was considered in many sources devoted to the theory of optimal control. Our presentation follows the monograph [2]. We introduce a variable

$x_0 = x_0(t)$, satisfying the equation $\dot{x}_0 = f_0(x, u, t)$, $x(t_0) = 0$ (where $f_0(x, u, t) \equiv \frac{1}{2}(x^*Q(t)x + u^*R(t)u)$). Then the problem (1)–(2) will be rewritten as

$$\begin{aligned} \frac{dx_0}{dt} &= f_0(x, u, t), x(0) = 0; \\ \frac{dx}{dt} &= A(t)x + B(t)u + h(t), x(0) = x^0; \\ x_0(T) &\rightarrow \min, \quad 0 \leq t \leq T. \end{aligned} \quad (3)$$

Denote $f(x, u, t) = \{f_1, \dots, f_n\} \equiv A(t)x + B(t)u + h(t)$ and make Hamiltonian

$$\begin{aligned} \tilde{H}(\psi, x, u, t) &= \psi_0 f_0 + \sum_{j=1}^n \psi_j f_j \equiv \psi_0 f + (\psi_1, \dots, \psi_n) \begin{pmatrix} f_1 \\ \vdots \\ f_n \end{pmatrix} = \\ &= \frac{1}{2}\psi_0(x^*Q(t)x) + \frac{1}{2}\psi_0(u^*R(t)u) + \psi^*A(t)x + \psi^*B(t)u + \psi^*h(t), \end{aligned}$$

where $\psi^* \equiv (\psi_1, \dots, \psi_n)$. According to the maximum principle should be $\psi_0(t) = \text{const} \leq 0$. In the problem with a fixed time T and with a free end of the trajectory are equalities

$$\psi_0(T) = -1, \psi_1(T) = \psi_2(T) = \dots = \psi_n(T) = 0$$

(transversality conditions; see [2; 260]), therefore $\psi_0(t) \equiv -1$, and the function \tilde{H} takes the form

$$\begin{aligned} \tilde{H}(\psi, x, u, t) &= -[\frac{1}{2}(x^*Q(t)x) + \frac{1}{2}(u^*R(t)u) - \\ &- \psi^*A(t)x - \psi^*B(t)u - \psi^*h(t)] \equiv -[\frac{1}{2}(x^*Q(t)x) + \\ &+ \frac{1}{2}(u^*R(t)u) + p^*A(t)x + p^*B(t)u + p^*h(t)], \end{aligned}$$

where it is indicated: $p^* \equiv -\psi^*$. The function \tilde{H} must reach a maximum on the optimal control $u = u(t)$, which means, that the function $\hat{H}(p^*, x, u, t) \equiv \frac{1}{2}(x^*Q(t)x) + \frac{1}{2}(u^*R(t)u) + p^*A(t)x + p^*B(t)u + p^*h(t)$ should reach to $u = u(t)$ the minimum. We note now that $\frac{dp_j}{dt} = -\frac{d\psi_j}{dt}$ ($j = \overline{1, n}$), therefore, the auxiliary functions $p_j \equiv -\psi_j(t)$ satisfy the system of differential equations $\frac{dp_j}{dt} = \frac{\partial \hat{H}}{\partial x_j}$, $j = \overline{1, n}$ (the equation $\dot{p}_0 = \frac{\partial \hat{H}}{\partial x_0}$ is not written out, since it is trivially satisfied, because $\psi_0(t) \equiv -1$, $\frac{\partial \hat{H}}{\partial x_0} \equiv 0$). In terms of the function \hat{H} , this system can be written in the form

$$\frac{dp_j}{dt} = -\frac{\partial \hat{H}}{\partial x_j}, \quad j = \overline{1, n}. \quad (4)$$

Calculate $\frac{\partial \hat{H}}{\partial x_j}$. We have (consider that $Q(t)$ is a symmetric matrix)

$$\begin{aligned} \frac{\partial \hat{H}}{\partial x_j} &= \frac{\partial}{\partial x_j}(\frac{1}{2}x^*Q(t)x + p^*A(t)x) = \frac{1}{2}e_j^*Q(t)x + \\ &+ \frac{1}{2}x^*Q(t)e_j + p^*A(t)e_j = \frac{1}{2}(Q(t)x, e_j) + \\ &+ \frac{1}{2}(Q(t)e_j, x) + (A(t)e_j, p) = \\ &= \frac{1}{2}(x, Q(t)e_j) + \frac{1}{2}(Q(t)e_j, x) + (A(t)e_j, p) = (e_j, Q(t)x) + \\ &+ (e_j, A^*(t)p) = (e_j, Q(t)x + A^*(t)p), \end{aligned}$$

where $e_j = \left\{ 0, \dots, 0, \underset{(j)}{1}, 0, \dots, 0 \right\}$, $j = \overline{1, n}$. Therefore, the system (4) has the form

$$\begin{aligned} \frac{dp_j}{dt} &= -(e_j, Q(t)x + A^*(t)p), j = \overline{1, n} \Leftrightarrow \\ &\Leftrightarrow \frac{dp}{dt} = -Q(t)x - A^*(t)p. \end{aligned} \quad (5)$$

On the other hand, to search for the minimum points of the function $\hat{H}(p^*, x_0, x, u, t)$ relative to u , we must find its critical points (note that $R(t)$ is a symmetric matrix):

$$\begin{aligned} & \frac{\partial \hat{H}}{\partial u_j} = 0, j = \overline{1, n} \Leftrightarrow \\ & \Leftrightarrow \frac{1}{2} e_j^* R(t) u + \frac{1}{2} u^* R(t) e_j + p^* B(t) e_j = 0 \Leftrightarrow \\ & \quad \Leftrightarrow (e_j, R(t) u) = -(B(t) e_j, p) \Leftrightarrow \\ & \quad \Leftrightarrow (e_j, R(t) u) = -(e_j, B^*(t) p), j = \overline{1, n} \\ & \Leftrightarrow R(t) u = -B^*(t) p \Leftrightarrow u = -R^{-1}(t) B^*(t) p. \end{aligned}$$

So, the only one critical point $u = -R^{-1}(t) B^*(t) p$ is obtained. In order to check whether it will realize the minimum of the function \hat{H} for u , consider the matrix of the second derivatives:

$$\begin{aligned} & \frac{\partial^2 \hat{H}}{\partial u_j \partial u_k} = \frac{\partial}{\partial u_k} [(e_j, R(t) u) + (e_j, B^*(t) p)] = \\ & = \frac{\partial}{\partial u_k} (e_j, R(t) u) = (e_j, R(t) e_k) = R_{jk}(t), j, k = \overline{1, n}, \end{aligned}$$

where R_{jk} are the elements of the matrix $R(t)$. This shows that the matrix $(\frac{\partial^2 \hat{H}}{\partial u_j \partial u_k})$ does not depend on u . Since $R(t)$ is a positive definite matrix, then the point $u = -R^{-1}(t) B^*(t) p$ is indeed the minimum point of the function \hat{H} for u . So, if the optimal control of problem (1)–(2) exists, then it necessarily has the form $u = -R^{-1}(t) B^*(t) p$, where $p = p(t)$ satisfies system (5) and $x = x(t)$ is calculated from the system (3). In other words, the optimal system has the form

$$\begin{aligned} \frac{dx}{dt} &= A(t)x - B(t)R^{-1}(t)B^*(t)p + h(t), x(0) = x^0; \\ \frac{dp}{dt} &= -Q(t)x - A^*(t)p, p(T) = 0. \end{aligned} \quad (6)$$

It follows from [3] that the control $u = -R^{-1}(t) B^*(t) p$, where $p = p(t)$ satisfies system (6), is optimal, and the corresponding trajectory $x = x(t)$ is the optimal trajectory. The boundary-value problem (6) with continuous matrices $A(t), B(t), R(t), Q(t)$ and with continuous function $h(t)$ on a segment $[0, T]$ can be ambiguously solvable, and then the optimal control $u = -R^{-1}(t) B^*(t) p$ is calculated ambiguously. For the unique solvability of problem (6), it is necessary to require that the corresponding homogeneous boundary value problem

$$\begin{aligned} \frac{dx}{dt} &= A(t)x - B(t)R^{-1}(t)B^*(t)p, x(0) = 0; \\ \frac{dp}{dt} &= -Q(t)x - A^*(t)p, p(T) = 0. \end{aligned} \quad (6_0)$$

Has only a trivial solution $(x(t), p(t)) \equiv 0$. This will take place, for example, in the case when $h(t) \equiv 0$ and $p(t)$ is linear depends on phase coordinates: $p(t) = K(t)x(t)$. Indeed, in this case (as shown in [3; 318, 319]), the $(n \times n)$ – matrix $K(t) \neq 0$ is symmetric and satisfies the nonlinear Riccati matrix differential equation:

$$\begin{aligned} \dot{K} &= -K \cdot A(t) - A^*(t) \cdot K + \\ &+ K \cdot B(t)R^{-1}(t) \cdot B^*(t) \cdot K - Q(t), K(T) = 0(t \in [0, T]), \end{aligned}$$

and homogeneous problem (6₀) takes the form

$$\begin{aligned} \frac{dx}{dt} &= [A(t) - B(t)R^{-1}(t)B^*(t)K(t)] x, x(0) = 0; \\ p(t) &= K(t)x(t). \end{aligned}$$

This problem has only one solution $(x(t), p(t)) \equiv 0$.

2 Singularly perturbed control problems. Building an optimal system

We now consider a singularly perturbed control system

$$\begin{aligned} \varepsilon \dot{x} &= A(t)x(t, \varepsilon) + B(t)u(t, \varepsilon) + f(t), x(0, \varepsilon) = x^0, 0 \leq t \leq T; \\ J_\varepsilon(u) &= \frac{1}{2} \int_0^T (x^* Q(t) x + u^* R(t) u) \exp\left(\frac{1}{\varepsilon} \int_0^t \mu(\theta) d\theta\right) dt, \end{aligned} \quad (7)$$

where $\varepsilon > 0$ is a small parameter, $x(t, \varepsilon), f(t)$ are n – dimensional, $u(t, \varepsilon)$ is m – dimensional vector functions, x^0 is a constant n -dimensional vector; $A(t)$ is $(n \times n)$ –matrix; $B(t)$ is $(n \times m)$ –matrix; $Q(t)$ is a symmetric

non-negative definite $(n \times n)$ - matrix; $R(t)$ is a symmetric positive definite $(m \times m)$ - matrix, $\mu(t)$ is a scalar function, $t \in [0, T]$, $*$ is a transposition sign. It is required to transfer the system (7) from a given initial position $x(0) = x^0$ to a position $x(T)$ in a fixed time $T < +\infty$ ($x(T)$ is not fixed) so that the functional $J(u)$ takes the minimum value. In order to obtain asymptotic representations for $x(t, \varepsilon)$ and $u(t, \varepsilon)$ in the form of series in powers of ε , we require the following conditions to be satisfied:

1⁰. The elements of the matrices $A(t), B(t), Q(t)$ and $R(t)$, as well as the components of the vector $f(t)$ and the scalar function $\mu(t)$ belong to $C^\infty([0, T], \mathbb{R})$.

Applying the Pontryagin's maximum principle (see system (1) in the previous section) and taking into account that the role of $A(t)$ is played by the matrix $\frac{1}{\varepsilon}A(t)$, role $B(t)$ - matrix $\frac{1}{\varepsilon}B(t)$, role $Q(t)$ - matrix $Q(t) \exp\left(\frac{1}{\varepsilon} \int_T^t \mu(\theta) d\theta\right)$; role $R(t)$ - matrix $R(t) \exp\left(\frac{1}{\varepsilon} \int_T^t \mu(\theta) d\theta\right)$; role $h(t)$ - function $\frac{1}{\varepsilon}f(t)$, we get the following optimal system:

$$\begin{aligned} \varepsilon^2 \dot{x} &= \varepsilon A(t)x - B(t)R^{-1}(t)B^*(t) \exp\left(-\frac{1}{\varepsilon} \int_T^t \mu(\theta) d\theta\right) p + \varepsilon f(t), x(0, \varepsilon) = x^0, \\ \varepsilon \dot{p} &= -\varepsilon Q(t) \exp\left(\frac{1}{\varepsilon} \int_T^t \mu(\theta) d\theta\right) x - A^*(t)p, p(T, \varepsilon) = 0. \end{aligned}$$

Doing successive replacements

$$\varepsilon x = y, \quad \exp\left(-\frac{1}{\varepsilon} \int_T^t \mu(\theta) d\theta\right) p = q, \quad \{y, q\} = z, \quad (8)$$

and taking into account that

$$\varepsilon \cdot \frac{d}{dt} p(t) = \left(\varepsilon \cdot \frac{d}{dt} q(t) \right) e^{\frac{1}{\varepsilon} \int_T^t \mu(\theta) d\theta} + q(t) \mu(t) e^{\frac{1}{\varepsilon} \int_T^t \mu(\theta) d\theta},$$

we arrive at the following singularly perturbed boundary value problem with weak inhomogeneity:

$$\begin{aligned} \varepsilon \frac{d}{dt} y(t) &= A(t)y(t) - B(t) \cdot R^{-1}(t) \cdot B^*(t) q(t) + \varepsilon f(t); \\ \varepsilon \frac{d}{dt} q(t) &= -q(t) \mu(t) - Q(t)y(t) - A^*(t) \cdot q(t), \end{aligned}$$

or

$$\begin{aligned} \varepsilon \dot{z} &= S(t)z + \varepsilon h(t), \quad 0 \leq t \leq T; \\ Gz &\equiv Mz(0, \varepsilon) + Nz(T, \varepsilon) = \varepsilon \alpha, \end{aligned} \quad (9)$$

where $S(t)$ is $(2n \times 2n)$ -matrix with elements from the class $C^\infty[0, T]$:

$$S(t) = \begin{pmatrix} A & -BR^{-1}B^* \\ -Q & -(A^* + \mu I) \end{pmatrix},$$

where $M = \text{diag}(1, \dots, 1, 0, \dots, 0)$, $N = \text{diag}(0, \dots, 0, 1, \dots, 1)$, $h(t) = f(t), 0, \dots, 0$, $\alpha = x^0, 0, \dots, 0$ are $2n$ - dimensional vectors. In this case, the optimal control (see 1) will be

$$u(t) = -\frac{1}{\varepsilon} R^{-1}(t) B^*(t) q(t). \quad (10)$$

3 Regularization of the optimal system. Construction of solutions of iterative problems

Without detracting from the generality, we may assume that $T = 1$. Let $b_j(t), d_j(t)$ ($j = \overline{1, 2n}$) are eigenvectors of matrices $S(t)$ and $S^*(t)$ corresponding to eigenvalues $\lambda_j(t)$ and $\bar{\lambda}_j(t)$, respectively, and, moreover, $(b_j(t), d_j(t)) = \delta_{ij}$ is Kronecker's symbol. Suppose that besides the condition 1⁰, two more conditions are performed:

2⁰. The spectrum $\{\lambda_j(t)\}$ of the matrix $S(t)$ has the properties:

- $\lambda_j(t) \neq 0, j = \overline{1, 2n}, \forall t \in [0, 1]$;
- $\lambda_i(t) \neq \lambda_j(t), i \neq j, i, j = \overline{1, 2n}, \forall t \in [0, 1]$;
- $\text{Re } \lambda_j(t) < 0, j = \overline{1, n}, \text{Re } \lambda_j(t) \geq 0, j = \overline{n+1, 2n}, \forall t \in [0, 1]$;
- $\text{Re } \lambda_1(t) \leq \text{Re } \lambda_2(t) \leq \dots \leq \text{Re } \lambda_{2n}(t)$.

$$3^0. \det(b_{ij}(0))_{i,j=1}^n \cdot \det(b_{ij}(1))_{i,j=n+1}^{2n} \neq 0.$$

Here $b_j = \text{colon}(b_{1j}, \dots, b_{2nj})$, $j = \overline{1, 2n}$; the conditions 3^0 mean that in the matrix $\mathfrak{B}(t) = (b_{i,j}(t))_{i,j=\overline{1, 2n}}$ the left angle minor $\det(b_{ij}(t))_{i,j=1}^n$ of order n does not vanish at the point $t = 0$, and the right angle minor $\det(b_{ij}(t))_{i,j=n+1}^{2n}$ of order n does not vanish at the point $t = 1$.

The listed conditions $1^0 - 3^0$ are realizable. To verify this, consider the following scalar problem:

$$\begin{aligned} \varepsilon \left(\frac{d}{dt} x(t) \right) &= -x(t) + e^t u(t) + f(t), \quad x(0) = x^0 \quad (0 \leq t \leq 1); \\ J_\varepsilon(u) &= \frac{1}{2} \int_0^1 (x(t)^2 + u(t)^2) e^{\frac{t-1}{\varepsilon}} dt \rightarrow \min. \end{aligned}$$

By completing the replacement (8), we obtain the optimal system

$$\begin{aligned} \varepsilon \begin{pmatrix} \dot{y} \\ \dot{z} \end{pmatrix} &= \begin{bmatrix} -1 & -e^{2t} \\ -1 & 2 \end{bmatrix} \begin{pmatrix} y \\ z \end{pmatrix} + \varepsilon \begin{pmatrix} f(t); \\ 0 \end{pmatrix} \quad (0 \leq t \leq 1), \\ y(0) &= \varepsilon x^0, \quad q(1) = 0. \end{aligned}$$

The eigenvalues and eigenvectors of the matrix $S(t)$ are follows:

$$\begin{aligned} \lambda_1(t) &= \frac{1}{2} - \frac{1}{2} \sqrt{9 + 4e^{2t}} \leftrightarrow b_1(t) = \begin{pmatrix} \frac{2e^{2t}}{-3 + \sqrt{9 + 4e^{2t}}} \\ 1 \end{pmatrix}, \\ \lambda_2(t) &= \frac{1}{2} + \frac{1}{2} \sqrt{9 + 4e^{2t}} \leftrightarrow b_2(t) = \begin{pmatrix} \frac{-2e^{2t}}{3 + \sqrt{9 + 4e^{2t}}} \\ 1 \end{pmatrix}. \end{aligned}$$

Obviously, all the conditions $1^0 - 3^0$ for them are realized.

We draw attention to the fact that in the considerate problem may arise a situation when some points of the spectrum on separate values (or even on some continual set) of independent variable can become purely imaginary. This situation is not amenable to the study of the known methods of asymptotic integration (for example, using the Vasilyeva-Butuzov method of boundary functions [4–7]. However, it can be fully studied using the Lomov’s regularization method [1, 8–10]. Note, that the regularization method [1, 8] allows us to investigate a wide class of problems in the theory of singular perturbations [11–25].

For the regularization of problem (9), we apply the algorithm of the regularization method, developed for singularly perturbed problems in [1]. We introduce regularizing variables

$$\begin{aligned} \tau_k &= \frac{1}{\varepsilon} \int_{t_k}^t \lambda_k(s) ds \equiv \varphi_k \left(t, \frac{1}{\varepsilon} \right), \quad k = \overline{1, 2n}; \\ t_1 &= \dots = t_n = 0, \quad t_{n+1} = \dots = t_{2n} = 1 \end{aligned}$$

and consider a new function $\tilde{z} = \tilde{z}(t, \tau, \varepsilon)$, for which we set the following problem:

$$\begin{aligned} \varepsilon \frac{\partial \tilde{z}}{\partial t} + \sum_{j=1}^{2n} \lambda_j(t) \frac{\partial \tilde{z}}{\partial \tau_j} - S(t) \tilde{z} &= \varepsilon h(t); \\ G\tilde{z}(t, \tau, \varepsilon) &\equiv M\tilde{z}(M_0, \varepsilon) + N\tilde{z}(M_1, \varepsilon) = \varepsilon \alpha, \end{aligned} \tag{11}$$

where

$$M_0 \left(0, \varphi \left(0, \frac{1}{\varepsilon} \right) \right), M_1 \left(1, \varphi \left(1, \frac{1}{\varepsilon} \right) \right), \varphi \left(t, \frac{1}{\varepsilon} \right) \equiv \left(\varphi_1 \left(t, \frac{1}{\varepsilon} \right), \dots, \varphi_{2n} \left(t, \frac{1}{\varepsilon} \right) \right).$$

If $\tilde{z} = \tilde{z}(t, \tau, \sigma, \varepsilon)$ is a solution to problem (11), then the restriction $z(t, \varepsilon) \equiv \tilde{z}(t, \varphi(t, \frac{1}{\varepsilon}), \varepsilon)$ will obviously be a solution to problem (9). The resulting problem (11) is regular in ε , and therefore we are looking for its solution in the form of a series

$$\tilde{z} = \sum_{i=0}^{\infty} \varepsilon^i z_i(t, \tau) \tag{12}$$

by non-negative powers of the parameter ε . To determine the coefficients of this series, we obtain the following iteration systems:

$$L_0 z_0 \equiv \sum_{j=1}^{2n} \lambda_j(t) \frac{\partial z_0}{\partial \tau_j} - S(t) z_0 = 0, \quad Gz_0(t, \tau) \equiv Mz_0(M_0) + Nz_0(M_1) = 0, \tag{13_0}$$

$$L_0 z_1 = h(t) - \frac{\partial z_0}{\partial t}, G z_1 = \alpha; \quad (13_1)$$

$$L_0 z_i = -\frac{\partial z_{i-1}}{\partial t}, G z_i = 0, \quad i = 2, 3, \dots \quad (13_i)$$

In order to study the solvability of problems (13₀), (13₁), (13_i), consider the general iterative system

$$L_0 \xi = H(t, \tau, \sigma). \quad (14)$$

We will find a solution to system (14) in the space of functions

$$U = \left\{ \xi(t, \tau) : \xi = \sum_{k=1}^{2n} \xi_k(t) e^{\tau k} + \xi_0(t), \xi_k(t) \in C^\infty([0, 1]), \mathbb{C}^{2n}, k = \overline{0, 2n} \right\},$$

in which is the following scalar (for each $t \in [0, 1]$) product is introduced

$$\langle \xi(t, \tau), \zeta(t, \tau) \rangle = \sum_{k=0}^{2n} (\xi_k(t), \zeta_k(t)), \quad \xi = \xi(t, \tau), \zeta(t, \tau) \in U,$$

where $(,)$ is the usual scalar product in the complex space \mathbb{C}^{2n} . Suppose that the right-hand side of system (14) belongs to U , that is $H(t, \tau) = \sum_{k=1}^{2n} h_k(t) e^{\tau k} + H_0(t) \in U$. The following assertion holds true, which is proved in the same way as the analogous assertion in [1].

Theorem 1. If conditions 1⁰, 2⁰(a, b, c) are satisfied and $H(t, \tau) \in U$, then for the solvability of system (14) in space U , it is necessary and sufficient that

$$\langle H(t, \tau), d_k(t) e^{\tau k} \rangle \equiv 0, \quad \forall t \in [0, 1], \quad k = \overline{1, 2n}. \quad (15)$$

Remark 1. Under the conditions of Theorem 1 the system (14) has the following solution in the space U :

$$\xi(t, \tau) = \sum_{k=1}^{2n} \left(\alpha_k(t) b_k(t) + \sum_{s=1, s \neq k}^{2n} H_{ks}(t) b_s(t) \right) e^{\tau k} + \sum_{k=1}^n H_{0k}(t) b_k(t); \quad (16)$$

where $\alpha_k(t) \in C^\infty([0, 1], \mathbb{C}^1)$ are arbitrary functions, $H_{ks}(t) \equiv (\lambda_k(t) - \lambda_s(t))^{-1} (H_k(t), d_s(t))$;

$$H_{0k}(t) = -(H_0(t), d_k(t)) / \lambda_k(t), \quad k, s = \overline{1, 2n}.$$

Theorem 2. Let the conditions 1⁰–3⁰ be satisfied, $H(t, \tau) \in U$ satisfies the requirements (15). Then the system (14) under boundary conditions $G\xi = \xi^0$ and additional conditions

$$\left\langle -\frac{\partial \xi}{\partial t} + g(t), d_k(t) e^{\tau k} \right\rangle \equiv 0, \quad \overline{1, 2n}, \quad \forall t \in [0, 1], \quad (17)$$

where $g(t) \in C^\infty([0, 1], \mathbb{C}^{2n})$ is a known function, is uniquely solvable in U for $\varepsilon \in (0, \varepsilon_0]$, where $\varepsilon_0 > 0$ is sufficiently small.

Proof. We use the conditions of orthogonality (17) and obtain equations for the unknown functions $\alpha_k(t)$ in the representation (16):

$$\dot{\alpha}_k(t) + (\dot{b}_k(t), d_k(t)) \alpha_k = - \sum_{s=1, s \neq k}^{2n} H_{ks} (\dot{b}_s(t), d_k(t)), \quad k = \overline{0, 2n}. \quad (18)$$

From equations (18) we find

$$\alpha_k(t) = e^{-\int_0^t (\dot{b}_k, d_k) d\theta} \left[\alpha_k(0) - \sum_{s=1, s \neq k}^{2n} \int_0^t e^{\int_0^x (\dot{b}_k, d_k) d\theta} H_{ks}(x) (\dot{b}_s(x), d_k(x)) dx \right], \quad (19)$$

where $\alpha_k(0)$, $k = \overline{1, 2n}$, are while unknown numbers. Subordinate the vector function (16) to the boundary condition $G\xi = \xi^0 \Leftrightarrow M\xi(0, \varphi(0, \frac{1}{\varepsilon})) + N\xi(1, \varphi(1, \frac{1}{\varepsilon})) = \xi^0$. To simplify the calculations, we write the solution (16) in the form

$$\xi(t, \tau) = \Phi(t) \text{diag}(e^{\tau_1}, \dots, e^{\tau_{2n}}) \alpha(t) + \hat{\xi}(t, \tau),$$

where $\alpha(t) = \{\alpha_1(t), \dots, \alpha_{2n}(t)\}$, $\Phi(t) \equiv (b_1(t), \dots, b_{2n}(t))$ is the matrix of the eigenvectors of the matrix $T(t)$, and by $\hat{\xi}(t, \tau)$ is denoted the particular solution of system (14):

$$\hat{\xi}(t, \tau) = \sum_{k=1}^{2n} \left(\sum_{s=1, s \neq k}^{2n} H_{ks}(t) b_s(t) \right) e^{\tau_k} + \sum_{k=1}^n H_{0k}(t) b_k(t).$$

The condition $G\xi = \xi^0$ gives:

$$\begin{aligned} & M\Phi(0) \text{diag}\left(1, \dots, 1, e^{\varphi_{n+1}(0, \frac{1}{\varepsilon})}, \dots, e^{\varphi_{2n}(0, \frac{1}{\varepsilon})}\right) \alpha(0) + \\ & + N\Phi(1) \text{diag}\left(e^{\varphi_1(1, \frac{1}{\varepsilon})}, \dots, e^{\varphi_n(1, \frac{1}{\varepsilon})}, 1, \dots, 1\right) \alpha(1) = \\ & = l(\varepsilon), \quad l(\varepsilon) \equiv \xi^0 - M\hat{\xi}\left(0, \varphi\left(0, \frac{1}{\varepsilon}\right)\right) + N\hat{\xi}\left(1, \varphi\left(1, \frac{1}{\varepsilon}\right)\right). \end{aligned} \quad (20)$$

From equality (19) we find that

$$\alpha_k(1) = e^{-\int_0^1 (\dot{b}_k, d_k) d\theta} \left[\alpha_k(0) - \sum_{s=1, s \neq k}^{2n} \int_0^1 \int_0^x (\dot{b}_k, d_k) d\theta H_{ks}(x) (\dot{b}_s(x), d_k(x)) dx \right], \quad k = \overline{1, 2n},$$

so

$$\alpha(1) = \text{diag}\left(e^{-\int_0^1 (\dot{b}_1, d_1) d\theta}, \dots, e^{-\int_0^1 (\dot{b}_{2n}, d_{2n}) d\theta}\right) \alpha(0) + \beta,$$

where β is a constant vector having the form

$$\begin{aligned} \beta = -\{ & \sum_{s=1, s \neq 1}^{2n} \int_0^1 e^{\int_0^x (\dot{b}_1, d_1) d\theta} H_{1s}(x) (\dot{b}_s(x), d_1(x)) dx, \dots; \\ & \sum_{s=1, s \neq 1}^{2n} \int_0^1 e^{\int_0^x (\dot{b}_{2n}, d_{2n}) d\theta} H_{1s}(x) (\dot{b}_s(x), d_{2n}(x)) dx\}. \end{aligned}$$

Substituting $\alpha(1)$ in (20), we will have

$$\begin{aligned} & [M\Phi(0) \text{diag}\left(1, \dots, 1, e^{\varphi_{n+1}(0, \frac{1}{\varepsilon})}, \dots, e^{\varphi_{2n}(0, \frac{1}{\varepsilon})}\right) + \\ & + N\Phi(1) \text{diag}\left(e^{\varphi_1(1, \frac{1}{\varepsilon}) - \int_0^1 (\dot{b}_1, d_1) d\theta}, \dots, e^{\varphi_n(1, \frac{1}{\varepsilon}) - \int_0^1 (\dot{b}_n, d_n) d\theta}, e^{-\int_0^1 (\dot{b}_{n+1}, d_{n+1}) d\theta}, \dots, e^{-\int_0^1 (\dot{b}_{2n}, d_{2n}) d\theta}\right)] \times \\ & \times \alpha(0) = l(\varepsilon) - N\beta. \end{aligned} \quad (21)$$

We divide the matrix $\Phi(t)$ into blocks of size $n \times n$: $\Phi(t) = \begin{pmatrix} \Phi_{11}(t) & \Phi_{12}(t) \\ \Phi_{21}(t) & \Phi_{22}(t) \end{pmatrix}$ and we introduce the notation:

$$\begin{aligned} \Lambda_1(t) &= \text{diag}(\lambda_1(t), \dots, \lambda_n(t)), \quad \Lambda_2(t) = \text{diag}(\lambda_{n+1}(t), \dots, \lambda_{2n}(t)); \\ G_1 &= \text{diag}\left(-\int_0^1 (\dot{b}_1, d_1) d\theta, \dots, -\int_0^1 (\dot{b}_n, d_n) d\theta\right); \\ G_2 &= \text{diag}\left(-\int_0^1 (\dot{b}_{n+1}, d_{n+1}) d\theta, \dots, -\int_0^1 (\dot{b}_{2n}, d_{2n}) d\theta\right). \end{aligned}$$

Then the determinant $\Delta(\varepsilon)$ of the system (21) can be written as

$$\Delta(\varepsilon) = \begin{vmatrix} \Phi_{11}(0) & \Phi_{12}(0) e^{\frac{1}{\varepsilon} \int_0^1 \Lambda_2(\theta) d\theta} \\ \Phi_{21}(1) e^{\frac{1}{\varepsilon} \int_0^1 \Lambda_1(\theta) d\theta + G_1} & \Phi_{22}(1) e^{G_2} \end{vmatrix}.$$

To calculate it, we apply the block Gauss method (see [26; 44, 45]). Multiply the first «column» of this determinant by $-\Phi_{11}^{-1}(0)\Phi_{12}(0)e^{\frac{1}{\varepsilon}\int_0^1\Lambda_2(\theta)d\theta}$ the matrix and adding the result to the second «column», we get that

$$\Delta(\varepsilon) = \begin{vmatrix} \Phi_{11}(0) & 0 \\ \Phi_{21}(1)e^{\frac{1}{\varepsilon}\int_0^1\Lambda_1(\theta)d\theta+G_1} & K \end{vmatrix} = \det \Phi_{11}(0) \cdot \det K,$$

where $K = \Phi_{22}(1)e^{G_2} - \Phi_{11}^{-1}(0)\Phi_{12}(0)e^{\frac{1}{\varepsilon}\int_0^1\Lambda_2(\theta)d\theta}\Phi_{21}(1)e^{\frac{1}{\varepsilon}\int_0^1\Lambda_1(\theta)d\theta+G_1}$. Considering conditions 2⁰c), we conclude that $\Delta(\varepsilon) \rightarrow \Delta(0) = \det \Phi_{11}(0) \cdot \det[\Phi_{22}(1)e^{G_2} (\varepsilon \rightarrow +0)]$. By virtue of the condition 3⁰ the determinant $\Delta(0) \neq 0$, and therefore $\Delta(\varepsilon) \neq 0$ for $\varepsilon \in (0, \varepsilon_0]$ and $\varepsilon_0 > 0$ is sufficiently small. This means that system (21) has a unique solution $\alpha(0) = \alpha^0(\varepsilon)$. Substituting this solution into (19), we find uniquely the functions $\alpha_k(t) = \alpha_k(t, \varepsilon)$, and, therefore, we calculate the solution (16) of system (14) in the space U in a unique way. The theorem is proved.

4 Correct solvability of a singularly perturbed two-point boundary value problem and estimation of the remainder term

Applying Theorems 1, 2 to iterative problems (13_i), we find in a unique way solutions of these problems in space U and construct series (12). Arguing by analogy with the monograph [1; 107–126], we prove the following result.

Lemma 1. Let the conditions 1⁰ – 3⁰ be fulfilled. Then the function $z_{\varepsilon N}(t) = S_N(t, \varphi(t, \frac{1}{\varepsilon}), \varepsilon) \equiv \sum_{i=0}^N \varepsilon^i z_i(t, \frac{\varphi(t)}{\varepsilon})$ satisfies the problem

$$\varepsilon \frac{dz_{\varepsilon N}}{dt} = S(t)z_{\varepsilon N}(t) + \varepsilon h(t) + \varepsilon^{N+1}R_N(t, \varepsilon), \quad Gz_{\varepsilon N}(t) = \varepsilon \alpha,$$

where $\|R_N(t, \varepsilon)\|_{C[0,1]} \leq \bar{R}_N = \text{const}$ for $\varepsilon \in (0, \varepsilon_0]$ ($\varepsilon_0 > 0$ is sufficiently small).

To substantiate the asymptotic convergence of the formal solution $z_{\varepsilon N}(t) = S_N(t, \varphi(t, \frac{1}{\varepsilon}), \varepsilon)$ of the problem (9) to its exact solution $z(t, \varepsilon)$, we must prove the solvability of an arbitrary boundary value problem

$$\varepsilon \frac{dz}{dt} = S(t)z + h(t), \quad Gz(t, \varepsilon) = z^0, \quad t \in [0, T], \quad (22)$$

under the above conditions 1⁰ – 3⁰ and give an estimate of its solution through $\|z^0\|$ and $\|h(t)\|$. Under conditions of stability of the spectrum $\{\lambda_j(t)\}$ of an operator $S(t)$, such an estimate was carried out using the Green function (see, for example, [1]), the construction of which essentially uses a rather strict condition 2⁰d). Here we propose another procedure based on the study of a special integral equation equivalent to system (22). The implementation of this procedure (its main ideas are presented in [27]) makes it possible to avoid constructing the Green function, and also to get rid of condition 2⁰d). We formulate some of the statements from [27], which will be used later.

Consider a more general, than (22), boundary value problem

$$\frac{dw}{dt} = S(t)w + h(t), \quad lw \equiv M_1w(0) + N_1w(T) = w^0, \quad t \in [0, T], \quad (23)$$

not containing a parameter. We assume that M_1 and N_1 are arbitrary $(m \times m)$ -matrices, $w = \{w_1, \dots, w_m\}$, $S(t)$ is a well-known $(m \times m)$ -matrix, $h(t) = \{h_1, \dots, h_m\}$ is a well-known vector function, $w^0 = \{w_1^0, \dots, w_m^0\}$ is a well-known constant vector. We formulate conditions under which the boundary value problem (23) is solvable.

Let $\Phi(t)$ be the fundamental matrix of solutions of the corresponding homogeneous system $\dot{w} = S(t)w$ with columns $c_j(t)$, i.e. $\Phi(t) = (c_1(t), \dots, c_m(t))$. Form the matrices

$$\begin{aligned} \Phi_1(t) &= (c_1(t), \dots, c_k(t); 0, \dots, 0); \\ \Phi_2(t) &= (0, \dots, 0; c_{k+1}(t), \dots, c_m(t)), \end{aligned}$$

where $k \in \{1, \dots, m\}$ is an arbitrary number. The following properties of these matrices are obvious:

- a) $\Phi_1(t) + \Phi_2(t) = \Phi(t)$; b) $\dot{\Phi}_j(t) \equiv S(t)\Phi_j(t)$, $j = 1, 2$; c) $\dot{\Phi}_1(t) + \dot{\Phi}_2(t) = \dot{\Phi}(t)$.

Lemma 2 [27]. Let the matrix $S(t) \in C^\infty([0, T], \mathbb{C}^{m \times m})$, $h(t) \in C^\infty([0, T], \mathbb{C}^m)$. Then the general solution of the system $\dot{w} = S(t)w + h(t)$ on the segment $[0, T]$ can be written as

$$w(t, c_0) = \Phi(t)c_0 + \Phi_1(t) \int_0^t \Phi^{-1}(\theta)h(\theta)d\theta + \Phi_2(t) \int_T^t \Phi^{-1}(\theta)h(\theta)d\theta, \quad (24)$$

where $c_0 \in \mathbb{C}^m$ is an arbitrary constant vector.

Proof. Since $\Phi(t)c_0$ it is a general solution of the corresponding homogeneous system, it is necessary to show that the vector function

$$\tilde{w}(t) = \Phi_1(t) \int_0^t \Phi^{-1}(\theta)h(\theta)d\theta + \Phi_2(t) \int_T^t \Phi^{-1}(\theta)h(\theta)d\theta \quad (25)$$

is a particular solution to a non-homogeneous system of equations $\dot{w} = S(t)w + h(t)$. Using the properties a) – c) of the matrices $\Phi_j(t)$, we will have

$$\begin{aligned} \frac{d\tilde{w}}{dt} &= \dot{\Phi}_1(t) \int_0^t \Phi^{-1}(\theta)h(\theta)d\theta + \Phi_1(t)\Phi^{-1}(t)h(t) + \\ &+ \dot{\Phi}_2(t) \int_T^t \Phi^{-1}(\theta)h(\theta)d\theta + \Phi_2(t)\Phi^{-1}(t)h(t) = \\ &= S(t)\Phi_1(t) \int_0^t \Phi^{-1}(\theta)h(\theta)d\theta + S(t)\Phi_2(t) \int_T^t \Phi^{-1}(\theta)h(\theta)d\theta + \\ &+ [\Phi_1(t) + \Phi_2(t)]\Phi^{-1}h(t) = S(t)[\Phi_1(t) \int_0^t \Phi^{-1}(\theta)h(\theta)d\theta + \\ &+ \Phi_2(t) \int_T^t \Phi^{-1}(\theta)h(\theta)d\theta] + h(t) = A(t)\tilde{w}(t) + h(t), \end{aligned}$$

that is, function (25) is indeed a solution of an non-homogeneous system $\dot{w} = S(t)w + h(t)$. The lemma is proved.

Lemma 3 [27]. Let the conditions of Lemma 2 be fulfilled. Then for the unique solvability of the boundary value problem (23) it is necessary and sufficient that

$$\det[M_1\Phi(0) + N_1\Phi(T)] \neq 0. \quad (26)$$

If condition (26) is satisfied, then the solution of the problem (23) is given by formula (24), where the vector c_0 has the form

$$\begin{aligned} c_0 &= [M_1\Phi(0)c_0 + M_1\Phi(T)]^{-1}(w^0 - M_1\Phi_2(0) \int_T^0 \Phi^{-1}(\theta)h(\theta)d\theta - \\ &- N_1\Phi_1(T) \int_0^T \Phi^{-1}(\theta)h(\theta)d\theta). \end{aligned} \quad (27)$$

Proof. Subject (24) to the boundary condition $lw = w^0$. We will have

$$\begin{aligned} M_1\Phi(0)c_0 + M_1\Phi_2(0) \int_T^0 \Phi^{-1}(\theta)h(\theta)d\theta + N_1\Phi(T)c_0 + \\ + N_1\Phi_1(T) \int_0^T \Phi^{-1}(\theta)h(\theta)d\theta = w^0; \end{aligned}$$

or

$$[M_1\Phi(0) + M_2\Phi(T)]c_0 = w^0 - M_1\Phi_2(0) \int_T^0 \Phi^{-1}(\theta)h(\theta)d\theta - \\ - N_1\Phi_1(t) \int_0^T \Phi^{-1}(\theta)h(\theta)d\theta.$$

For the unique solvability of this system, it is necessary and sufficient to satisfy condition (26). At the same time, c_0 has the form (27). The lemma is proved.

We now turn to the study of the solvability of the boundary value problem (23). According to [1; 109, 110], under the assumption that $S(t) \in C^1([0, T], \mathbb{C}^{2n \times 2n})$ transformation

$$z = [B(t)(I + \varepsilon B_1(t))]\eta(t, \varepsilon),$$

where $B^{-1}(t)T(t)B(t) = \Lambda(t) = \text{diag}(\lambda_1(t), \dots, \lambda_{2n}(t))$;

$$(B_1(t))_{ij} = \begin{cases} 0, & i = j, \\ \frac{1}{\lambda_i(t) - \lambda_j(t)} (B^{-1}(t)B'(t))_{ij}, & i \neq j \quad (i, j = \overline{1, 2n}), \end{cases}$$

leads system (22) to the system

$$\varepsilon \frac{d\eta}{dt} = \Lambda_0(t, \varepsilon)\eta + \varepsilon^2 C(t, \varepsilon)\eta + H_1(t, \varepsilon); \\ P_1(\varepsilon)\eta(0, \varepsilon) + P_2(\varepsilon)\eta(T, \varepsilon) = z^0, \quad (28)$$

where indicated:

$$P_1(\varepsilon) = M[B(0)(I + \varepsilon B_1(0))], \quad P_2(\varepsilon) = N[B(T)(I + \varepsilon B_1(T))]; \\ \Lambda_0(t, \varepsilon) = \text{diag}(\lambda_1(t) + \varepsilon\mu_1(t), \dots, \lambda_{2n}(t) + \varepsilon\mu_{2n}(t)); \\ H_1(t) = [B(t)(I + \varepsilon B_1(t))]^{-1}h(t)$$

and $C(t, \varepsilon)$ is a known matrix. Let $\Phi(t, \varepsilon)$ be the fundamental solution system of a shortened system

$$\varepsilon \frac{dz}{dt} = \Lambda_0(t, \varepsilon)z. \quad (29)$$

Obviously, it can be taken in the following form:

$$\Phi(t, \varepsilon) = \begin{pmatrix} e^{\frac{1}{\varepsilon} \int_0^t \bar{\Lambda}(\theta, \varepsilon) d\theta} & 0 \\ 0 & e^{\frac{1}{\varepsilon} \int_T^t \bar{\Lambda}(\theta, \varepsilon) d\theta} \end{pmatrix}, \quad (30)$$

where

$$\bar{\Lambda}(t) = \text{diag}(\lambda_1(t) + \varepsilon\mu_1(t), \dots, \lambda_n(t) + \varepsilon\mu_n(t)); \\ \bar{\bar{\Lambda}}(t) = \text{diag}(\lambda_{n+1}(t) + \varepsilon\mu_{n+1}(t), \dots, \lambda_{2n}(t) + \varepsilon\mu_{2n}(t)).$$

Using (30) and Lemma 3, we reverse the system (28) and obtain an equivalent system of integral equations

$$\eta(t, \varepsilon) = \Phi(t, \varepsilon)c_0(\eta) + \\ + \Phi_1(t, \varepsilon) \int_0^t \varepsilon \Phi^{-1}(s, \varepsilon) C(s, \varepsilon) \eta(s, \varepsilon) ds + \\ + \frac{\Phi_1(t, \varepsilon)}{\varepsilon} \int_0^t \Phi^{-1}(s, \varepsilon) H_1(s, \varepsilon) ds + \\ + \Phi_2(t, \varepsilon) \int_T^t \varepsilon \Phi^{-1}(s, \varepsilon) C(s, \varepsilon) \eta(s, \varepsilon) ds + \\ + \frac{\Phi_2(t, \varepsilon)}{\varepsilon} \int_T^t \Phi^{-1}(s, \varepsilon) H_1(s, \varepsilon) ds, \quad (31)$$

where

$$\begin{aligned}
 c_0(\eta) &= [P_1(\varepsilon)\Phi(0, \varepsilon) + P_2(\varepsilon)\Phi(T, \varepsilon)]^{-1} \times \\
 &\times \left[z^0 - P_1(\varepsilon)\Phi_2(0, \varepsilon) \int_T^0 \varepsilon\Phi^{-1}(s, \varepsilon)C(s, \varepsilon)\eta(s, \varepsilon)ds - \right. \\
 &\quad - P_2(\varepsilon)\Phi_1(T, \varepsilon) \int_0^T \varepsilon\Phi^{-1}(s, \varepsilon)C(s, \varepsilon)\eta(s, \varepsilon)ds - \\
 &\quad - P_1(\varepsilon) \frac{\Phi_2(0, \varepsilon)}{\varepsilon} \int_T^0 \Phi^{-1}(s, \varepsilon)H_1(s, \varepsilon)ds - \\
 &\quad \left. - P_2(\varepsilon) \frac{\Phi_1(T, \varepsilon)}{\varepsilon} \int_0^T \Phi^{-1}(s, \varepsilon)H_1(s, \varepsilon)ds \right]
 \end{aligned} \tag{32}$$

and by $\Phi_j(t, \varepsilon)$ we denoted $(2n \times 2n)$ -matrices

$$\Phi_1(t, \varepsilon) = \text{diag}\left(e^{\frac{1}{\varepsilon} \int_0^t \bar{\Lambda}(\theta, \varepsilon) d\theta}, 0\right); \quad \Phi_2(t, \varepsilon) = \text{diag}\left(0, e^{\frac{1}{\varepsilon} \int_T^t \bar{\Lambda}(\theta, \varepsilon) d\theta}\right).$$

Substituting (32) into (31), we write the integral system in operator form:

$$\eta = A\eta. \tag{33}$$

Obviously, the operator A acts from space $C([0, T], \mathbb{C}^{2n})$ to itself. In assessing the norm $\|A\eta_1 - A\eta_2\|$, we use the fact that the product $\Phi_1(t, \varepsilon)\Phi^{-1}(s, \varepsilon)$ is uniformly bounded of ε (for sufficiently small $\varepsilon > 0$) for s and t satisfying the inequalities $0 \leq s \leq t \leq T$, and the product $\Phi_2(t, \varepsilon)\Phi^{-1}(s, \varepsilon)$ is uniformly bounded (for sufficiently small $\varepsilon > 0$) for s and t satisfying the inequalities $0 \leq t \leq s \leq T$. Let us show this. We have at $0 \leq s \leq t \leq T$:

$$\begin{aligned}
 &\|\Phi_1(t, \varepsilon)\Phi^{-1}(s, \varepsilon)\| \equiv \\
 &\equiv \|\exp\{\frac{1}{\varepsilon} \int_s^t \bar{\Lambda}(\theta) d\theta\}\| = \|\exp\{\frac{1}{\varepsilon} \int_s^t \text{Re}\bar{\Lambda}(\theta) d\theta\}\| \leq \\
 &\leq \|\text{diag}(\exp\{\frac{1}{\varepsilon} \int_s^t \text{Re}\lambda_1 d\theta\}, \dots, \exp\{\frac{1}{\varepsilon} \int_s^t \text{Re}\lambda_n d\theta\})\| \times \\
 &\times \|\text{diag}(\exp\{\int_s^t \text{Re}\mu_1 d\theta\}, \dots, \exp\{\int_s^t \text{Re}\mu_n d\theta\})\| \leq \nu_1 = \text{const};
 \end{aligned}$$

because $\text{Re}\lambda_i(\theta) \leq 0$ when $0 \leq s \leq \theta \leq t \leq T, i = \overline{1, n}$. When $0 \leq t \leq s \leq T$ we have

$$\begin{aligned}
 &\|\Phi_2(t, \varepsilon)\Phi^{-1}(s, \varepsilon)\| \equiv \\
 &\equiv \|\exp\{\frac{1}{\varepsilon} \int_s^t \bar{\Lambda} d\theta\}\| = \|\exp\{-\frac{1}{\varepsilon} \int_t^s \text{Re}\bar{\Lambda} d\theta\}\| \leq \\
 &\leq \|\text{diag}(\exp\{-\frac{1}{\varepsilon} \int_t^s \text{Re}\lambda_{n+1} d\theta\}, \dots, (\exp\{-\frac{1}{\varepsilon} \int_t^s \text{Re}\lambda_{2n} d\theta\})\| \times \\
 &\times \|\text{diag}(\exp\{\int_s^t \text{Re}\mu_{n+1} d\theta\}, \dots, \exp\{\int_s^t \text{Re}\mu_{2n} d\theta\})\| \leq \\
 &\leq \nu_2 = \text{const},
 \end{aligned}$$

since $\text{Re}\lambda_j(\theta) \geq 0$ when $0 \leq t \leq \theta \leq s \leq T, j = \overline{n+1, 2n}$. In this case, constants ν_1 and ν_2 does not depend on ε when $\varepsilon \in (0, \varepsilon_0]$, where $\varepsilon_0 > 0$ is sufficiently small.

We now turn to the estimate $\|A\eta_1 - A\eta_2\| \equiv \|A\eta_1 - A\eta_2\|_{C[0,T]}$. Using the boundedness of matrices $\Phi_1(t, \varepsilon) \times \Phi^{-1}(t, \varepsilon)$ and $\Phi_2(t, \varepsilon) \cdot \Phi^{-1}(t, \varepsilon)$, we will have

$$\begin{aligned} \|A\eta_1 - A\eta_2\| &\leq \|\Phi(t, \varepsilon)(c_0(\eta_1) - c_0(\eta_2))\| + \\ &+ \varepsilon \|\Phi_1(t, \varepsilon) \int_0^t \Phi^{-1}(s, \varepsilon) C(s, \varepsilon) (\eta_1(s, \varepsilon) - \eta_2(s, \varepsilon)) ds\| + \\ &+ \varepsilon \|\Phi_2(t, \varepsilon) \int_T^t \Phi^{-1}(s, \varepsilon) C(s, \varepsilon) (\eta_1(s, \varepsilon) - \eta_2(s, \varepsilon)) ds\| \leq \\ &\leq \nu_0 \|c_0(\eta_1) - c_0(\eta_2)\| + \varepsilon \nu_3 \|\eta_1 - \eta_2\| + \varepsilon \nu_4 \|\eta_1 - \eta_2\|. \end{aligned}$$

On the other hand,

$$\begin{aligned} \|c_0(\eta_1) - c_0(\eta_2)\| &\leq \|(P_1(\varepsilon)\Phi(0, \varepsilon) + P_2(\varepsilon)\Phi(T, \varepsilon))^{-1}\| \times \\ &\times (\|\varepsilon P_1(\varepsilon)\Phi_2(0, \varepsilon) \int_0^T \Phi^{-1}(s, \varepsilon) C(s, \varepsilon) (\eta_1(s, \varepsilon) - \eta_2(s, \varepsilon)) ds\| + \\ &+ \|\varepsilon P_2(\varepsilon)\Phi_1(T, \varepsilon) \int_0^T \Phi^{-1}(s, \varepsilon) C(s, \varepsilon) (\eta_1(s, \varepsilon) - \eta_2(s, \varepsilon)) ds\|) \leq \\ &\leq \varepsilon \nu_5 \|\eta_1 - \eta_2\|. \end{aligned}$$

Substituting this into the previous inequality, we get $\|A\eta_1 - A\eta_2\| \leq \varepsilon \nu_6 \|\eta_1 - \eta_2\|$, where ν_6 do not depend on ε at $\varepsilon \in (0, \varepsilon_0]$. From here it follows that the operator A is contraction operator in space $C([0, T], \mathbb{C}^{2n})$, and therefore, the equation (33) is uniquely solvable in $C([0, T], \mathbb{C}^{2n})$.

From (31) and (32), passing to the norms in $C([0, T], \mathbb{C}^{2n})$, we get

$$\|\eta(t, \varepsilon)\| \leq \nu_0 \|c_0(\eta)\| + \varepsilon \nu_7 \|\eta\| + \frac{\nu_8}{\varepsilon} \|H_1\|;$$

$$\|c_0(\eta)\| \leq \nu_9 \|z^0\| + \varepsilon \nu_{10} \|\eta\| + \frac{\nu_{11}}{\varepsilon} \|H_1\|,$$

which means, that

$$\|\eta\| \leq \frac{1}{1 - \varepsilon(\nu_7 + \nu_{10}\nu_0)} (\nu_0\nu_9 \|z^0\| + \frac{\nu_8 + \nu_0\nu_{11}}{\varepsilon} \|H_1\|).$$

Take $\varepsilon > 0$ such that

$$1 - \varepsilon(\nu_7 + \nu_{10}\nu_0) \geq \frac{1}{2}.$$

Then

$$\|\eta\| \leq 2(\nu_0\nu_9 \|z^0\| + \frac{\nu_8 + \nu_0\nu_{11}}{\varepsilon} \|H_1\|).$$

Since $z = [B(t)I + \varepsilon B_1(t)]\eta(t, \varepsilon)$, then the initial boundary problem (22) is uniquely solvable in $C^1([0, T], \mathbb{C}^{2n})$ and for its solution $z(t, \varepsilon)$ we have the estimate

$$\|z(t, \varepsilon)\|_{C[0,T]} \leq K_1 \|z^0\| + \frac{K_2}{\varepsilon} \|h\|_{C[0,T]}. \quad (34)$$

However, all our calculations are correct only when

$$|\det[P_1(\varepsilon)\Phi(0, \varepsilon) + P_2(\varepsilon)\Phi(T, \varepsilon)]| \geq \delta_1 = \text{const} > 0 \quad (35)$$

at $\varepsilon \in (0, \varepsilon_0]$. Insofar as $P_1(\varepsilon) = MB(0) + \varepsilon MB_1(0)$, $P_2(\varepsilon) = NB(T) + \varepsilon NB_1(T)$, then inequality (35) for $\varepsilon \in (0, \varepsilon_0]$ follows from the fact that

$$|\det[MB(0)\Phi(0, \varepsilon) + NB(T)\Phi(T, \varepsilon)]| \geq \delta_2 = \text{const} > 0$$

for $\varepsilon \in (0, \varepsilon_0]$. This fact follows from the conditions $3^0a) - 3^0c)$, what can be proved in the same way as the inequality $|\Delta(\varepsilon)| \geq \delta_0 = \text{const}$ (see system (21) and the following calculations).

The following result is proved.

Theorem 3. Suppose that $S(t) \in C^1([0, T], \mathbb{C}^{2n \times 2n})$, $h(t) \in C([0, T], \mathbb{C}^{2n})$ and the conditions $1^0, 2^0a) - 2^0c), 3^0$ are satisfied. Then for sufficiently small $\varepsilon \in (0, \varepsilon_0]$ the boundary value problem (22) has a unique solution $z(t, \varepsilon)$ and estimate (34) is valid for it.

Using the estimate (34) for the remainder term $r_N(t, \varepsilon) = z(t, \varepsilon) - z_{\varepsilon N}(t)$, as well as Lemma 1, we can easily prove the following statement.

Theorem 4. Let the conditions $1^0, 2^0a) - 2^0c), 3^0$ are fulfilled. Then the boundary value problem (9) for sufficiently small $\varepsilon > 0$ ($0 < \varepsilon \leq \varepsilon_0$) is uniquely solvable in the class $C^1([0, T], \mathbb{C}^{2n})$ and the estimate

$$\|z(t, \varepsilon) - z_{\varepsilon N}(t)\|_{C[0, T]} \leq C_N \varepsilon^{N+1}$$

holds true (here $C_N > 0$ is a constant independent of $\varepsilon \in (0, \varepsilon_0]$, $z_{\varepsilon N}(t) = S_N(t, \varphi(t, \frac{1}{\varepsilon}), \varepsilon)$ the formal asymptotic solution constructed above).

So, we have obtained the asymptotic solution of the problem (9) in the form of a series (12), taken at the restriction $\tau = \varphi(t, \frac{1}{\varepsilon})$. Using equations (8) and (10), we construct the asymptotics of the optimal control $u(t, \varepsilon)$ and the optimal trajectory $x(t, \varepsilon)$.

Remark 2. Due to the uniqueness of the solution of the system (13₀) in space U , it will have only a trivial solution $z_0 = z_0(t, \tau) \equiv 0$, therefore the asymptotic series for the optimal control (10) and the optimal trajectory $x(t, \varepsilon) = \frac{1}{\varepsilon} y(t, \varepsilon)$ will not contain coefficients with a negative degree of the parameter ε .

Remark 3. Condition $2^0d)$ is related to the construction of the Green function for the boundary value problem (9) and its application in estimating the remainder term (see [1; 108–126]). Using our technique, one can do without constructing the Green function and then condition $2^0d)$ can be removed.

Acknowledgement

This work is supported by the grant № AP05133858 «Contrast structures in singularly perturbed equations and their application in the theory of phase transitions» (2018–2020) by the Committee of Science, Ministry of Education and Science of the Republic of Kazakhstan.

References

- 1 Ломов С.А. Введение в общую теорию сингулярных возмущений / С.А. Ломов. — М.: Наука, 1981.
- 2 Ройтенберг Я.Н. Автоматическое управление / Я.Н. Ройтенберг. — М.: Наука, 1971.
- 3 Болтянский И.Г. Достаточные условия оптимальности и обоснования метода динамического программирования / И.Г. Болтянский // Изв. АН СССР. Сер. Матем. — Т. 28, № 3. — 1964. — С. 481–514.
- 4 Глизер В.Я. Метод пограничного слоя при решении некоторых задач аналитического проектирования регулятора / В.Я. Глизер, В.Я. Глизер, М.Г. Дмитриев // Тез. докл. Всесоюз. конф. по асимпт. методам. — Фрунзе: Илим, 1975. — С. 112–115.
- 5 Васильева А.Б. Об асимптотике решения типа контрастной структуры / А.Б. Васильева, В.Ф. Бутузов // Матем. заметки. — 1987. — Т. — 42. — №. 6. — С. 831–841.
- 6 Бутузов В.Ф. Асимптотическая теория контрастных структур / В.Ф. Бутузов, А.Б. Васильева, Н.Н. Нефедов // Автомат. и телемех. — 1977. — 7. — С. 3–32.
- 7 Васильева А.Б. Контрастные структуры в сингулярно возмущенных задачах / А.Б. Васильева, В.Ф. Бутузов, Н.Н. Нефедов // Фундамент. и приклад. матем. — 1998. — Т. 4(3). — С. 799–851.
- 8 Ломов С.А. Основы математической теории пограничного слоя / С.А. Ломов, И.С. Ломов. — М.: Изд-во МГУ, 2011.
- 9 Ломов С.А. Теория сингулярных возмущений в случае спектральных особенностей предельного оператора / С.А. Ломов, А.Г. Елисеев // Матем. сб. — 1986. — Т. 131 (173), № 4 (26). — С. 544–557.
- 10 Бободжанов А.А. Регуляризация оптимальных сингулярно возмущенных систем с быстро меняющейся демпфирующей функцией / А.А. Бободжанов, Н.А. Рашепкина, В.Ф. Сафонов // Вестн. МЭИ. — 1998. — № 6. — С. 18–26.
- 11 Иманбаев Н.С. Алгоритм метода регуляризации для сингулярно возмущенной задачи с нестабильным значением ядра интегрального оператора / Н.С. Иманбаев, Б.Т. Калимбетов, М.А. Темирбеков // Вестн. Караганд. ун-та. Сер. математика. — 2013. — № 4(72). — С. 64–70.

- 12 Калимбетов Б.Т. Математическое описание внутреннего пограничного слоя для нелинейной интегродифференциальной системы / Б.Т. Калимбетов, Б.И. Ескараева, М.А. Темирбеков // Вестн. Караганд. ун-та. Сер. математика. — 2014. — № 3(75). — С. 77–87.
- 13 Ескараева Б.И. Дискретный пограничный слой в случае нулевых точек спектра для систем интегродифференциальных уравнений / Б.И. Ескараева, Б.Т. Калимбетов, М.А. Темирбеков // Вестн. Караганд. ун-та. Сер. математика. — № 3(75). — 2014. — С. 88–95.
- 14 Kalimbetov B.T. Internal boundary layer for integral-differential equations with zero spectrum of the limit operator and rapidly changing kernel / B.T. Kalimbetov, B.I. Yeskarayeva, A.S. Tolep // Applied Mathematical Sciences. — 9. — 141–144. — 2015. — P. 7149–7165.
- 15 Kalimbetov B.T. Asymptotic solution of singular perturbed problems with an instable spectrum of the limiting operator / B.T. Kalimbetov, M.A. Temirbekov, Zh.O. Habibullaev // Abstract and Applied Analysis. — 2012. Article ID 120192.
- 16 Kalimbetov B.T. Regularized asymptotical solutions of integro-differential systems with spectral singularities / B.T. Kalimbetov, N.S. Imanbaev, D.A. Sapakov, L.T. Tashimov // Advances in Difference Equations. — 2013. — Vol. 109. — DOI: 10.1186/1687-1847-2013-109.
- 17 Safonov V.F. A regularization method for systems with instable spectral of the kernel of the integral operator / V.F. Safonov, B.T. Kalimbetov // Differential Equations. — 31. — 4. — 1995. — P. 647–656.
- 18 Safonov V.F. Regularized asymptotic solutions of the initial problem of systems of integro-partial differential equations / V.F. Safonov, A.A. Bobodzhanov // Mathematical Notes. — 102. — 1. — 2017. — P. 22–30.
- 19 Safonov V.F. Regularized asymptotics of solutions to integro-differential partial differential equations with rapidly varying kernels / V.F. Safonov, A.A. Bobodzhanov // Ufimsk. Mat. Zh. — 10. — 2. — 2018. — P. 3–12.
- 20 Bobodzhanov A.A. A generalization of the regularization method to the singularly perturbed integro-differential equations with partial derivatives / A.A. Bobodzhanov, V.F. Safonov // Russian Math. (Iz. vuz). — 62. — 3. — 2018. — P. 6–17.
- 21 Bobodzhanov A.A. A problem with inverse time for a singularly perturbed integro-differential equation with diagonal degeneration of the kernel of high order / A.A. Bobodzhanov, V.F. Safonov // Izv. Math. — 80. — 2. — 2016. — P. 285–298.
- 22 Bobodzhanov A.A. Asymptotic solutions of Fredholm integro-differential equations with rapidly changing kernels and irreversible limit operator / A.A. Bobodzhanov, V.F. Safonov // Russian Math. (Iz. VUZ). — 59. — 10. — 2015. — P. 1–15.
- 23 Bobodzhanov A.A. The method of normal forms for singularly perturbed systems of Fredholm integro-differential equations with rapidly varying kernels / A.A. Bobodzhanov, V.F. Safonov // Sb. Math. — 204. — 7. — 2013. — P. 979–1002.
- 24 Safonov V.F. Volterra integral equations with rapidly varying kernels and their asymptotic integration / V.F. Safonov, A.A. Bobodzhanov // Sb. Math. — 192. — 8. — 2001. — P. 1139–1164.
- 25 Safonov V.F. «Splashes» in Fredholm Integro-Differential Equations with Rapidly Varying Kernels / V.F. Safonov, A.A. Bobodzhanov // Math. Notes. — 85. — 2. — 2009. — P. 153–167.
- 26 Gantmakher F.R. Theory of matrices / F.R. Gantmakher. — Moscow: GITTL, 1953.
- 27 Konyaev Yu.A. On a method of studying some problems of the perturbation theory / Yu.A. Konyaev // Mathem. Coll. — 1993. — No. 12. — P. 133–144.

А.А. Бободжанов, Б.Т. Калимбетов, В.Ф. Сафонов

Оңтайлы жүйенің матрицасының спектрі ұдайы болған жағдайдағы басқарудың сингуляр ауытқымалы есебі

Мақалада сапаның квадрат функционалы бар сингуляр ауытқымалы есебі қарастырылды. Мұндай есептер бұрын спектрдің белгілі шектеулерінде (оңтайлы жүйенің спектр нүктелері таза жорамал емес және жорамал оське қатысты симметриялы орналасса) Васильева-Бутузовтың шекаралық функциялар әдісі арқылы қарастырылған. Егер спектрдің нүктелерінің ең болмағанда біреуі тәуелсіз айналымының кейбір мәндерінде жорамал оське түссе, шекаралық функциялар әдісі жұмыс істемейді.

Жұмыста спектрдің таза жорамал нүктелері бар жағдай зерттелді. Бұл жағдайда С.А. Ломовтың регуляризациялау әдісінің идеяларына негізделген басқа әдісті дамытуға тура келеді. Сондай-ақ бұрын қаралған басқару есептерінде шығындар функционалы кіші параметрге мүлдем тәуелді емес немесе кіші параметрге тегіс тәуелді болатыны байқалған. Бұл мақалада шығынның кіші параметрге регуляр емес тәуелділігі, яғни интеграл таңбасы астында экспоненциалды көбейткіш түрінде жылдам өзгертін демпфирлеу функциясының бар болуы жағдайы қарастырылған. Осы жағдайда оңтайлы жүйе спектрінің беталысы демпфирлеу коэффициентіне байланысты болады, ол (белгілі бір жағдайларда) спектрді комплекс жазықтықтың қандай да бір жағына жылжытуы мүмкін. Онда спектрдің кейбір нүктелері жекеленген мәндерде (немесе тіпті кейбір континуалды жиындарда) тәуелсіз айнымалы таза жорамал болған жағдай туындауы мүмкін. Мұндай жағдайды жоғарыда аталған Васильева-Бутузовтың шекаралық функциялары әдісімен зерттеуге болмайды. Алайда қарастырылып отырған басқару есебін зерттеуге қатысты С.А. Ломовтың регуляризациялау әдісінің алгоритмін қолдану осы жұмыста толық қарастырылған. Бұл әдісті баяндау оңтайлы басқарудың классикалық есебі үшін Л.С. Понтрягиннің максимум принципі қысқаша сипаттаудан басталып, одан соң басқа идеялармен қатар, қарастырылып отырған басқару есебінің нәтижелерін негіздеу үшін қолданылды.

Кілт сөздер: сингуляр ауытқу, Понтрягиннің максимум принципі, регуляризация, асимптотикалық жинақтылық.

А.А. Бободжанов, Б.Т. Калимбетов, В.Ф. Сафонов

Сингулярно возмущенные задачи управления в случае стабильности спектра матрицы оптимальной системы

В статье рассмотрена сингулярно возмущенная задача управления с квадратичным функционалом качества. Такие задачи в их стандартной постановке при известных ограничениях на спектр (точки спектра оптимальной системы не являются чисто мнимыми и расположены симметрично относительно мнимой оси) были рассмотрены ранее с помощью метода пограничных функций Васильевой-Бутузова. Если же хотя бы одна из точек спектра при некоторых значениях независимой переменной попадает на мнимую ось, метод погранфункций не работает. Именно такая ситуация с допущением чисто мнимых точек спектра исследована в настоящей работе. В этом случае приходится развивать другой подход, основанный на идеях метода регуляризации С.А. Ломова. Следует заметить также, что в рассмотренных ранее задачах управления функционал затрат либо вообще не зависел от малого параметра, либо допускал гладкую зависимость от параметра. В данной работе допущена нерегулярная зависимость от малого параметра, в частности, наличие в них быстро изменяющейся функции демпфирования в виде экспоненциально множителя под знаком интеграла. В этом случае поведение спектра оптимальной системы зависит от коэффициента демпфирования, который (при определенных условиях) может смещать спектр в ту или иную сторону в комплексной плоскости. При этом может возникнуть ситуация, когда некоторые точки спектра при отдельных значениях (или даже на некотором континуальном множестве) независимой переменной могут становиться чисто мнимыми. Эта ситуация не поддается исследованию упомянутым ранее методом пограничных функций Васильевой-Бутузова. Однако его можно полностью изучить с помощью метода регуляризации С.А. Ломова, алгоритм которого применительно к рассматриваемой задаче управления развивается в настоящей работе. Изложение этого метода начинается с краткого описания принципа максимума Л.С. Понтрягина для классической задачи оптимального управления, который затем, наряду с другими идеями, применяется для обоснования результатов в рассматриваемой задаче управления.

Ключевые слова: сингулярное возмущение, принцип максимума Понтрягина, регуляризация, асимптотическая сходимость.

References

- 1 Lomov, S.A. (1981). *Vvedenie v obshchuiu teoriyu sinhuliarnykh vozmushchenii [Introduction to the general theory of singular perturbations]*. Moscow: Nauka [in Russian].
- 2 Reutenberg, J.N. (1971). *Avtomaticheskoe upravlenie [Automatic control]*. Moscow: Nauka [in Russian].

- 3 Boltyansky, I.G. (1964). Dostatochnye usloviia optimalnosti i obosnovaniia metoda dinamicheskogo programmirovaniia [Sufficient conditions for optimality and justification of the dynamic programming method]. *Izvestiia Akademii nauk SSSR. Serii Matematika – News of Academy of Sciences of the USSR, Mathematical series*, 28, 3, 481–514 [in Russian].
- 4 Glizer, V.Ya., & Dmitriev, M.G. (1975). Metod pohranichnogo sloia pri reshenii nekotorykh zadach analiticheskogo proektirovaniia rehulyatora [The boundary layer method in solving some problems of analytical design of the regulator]. *Tezisy dokladov Vsesoiuznoi konferentsii po asimptoticheskim metodam – Abstracts of the All-Union conf. by asymp. methods, Frunze: Ilim*, 112–115 [in Russian].
- 5 Vasilyeva, A.B., & Butuzov, V.F. (1987). Ob asimptotike resheniia tipa kontrastnoi struktury [On the asymptotics of a solution of the type of contrast structure]. *Matematicheskie zametki – Mathem. notes*, 42, 6, 831–841 [in Russian].
- 6 Butuzov, V.F., Vasilyeva, A.B., & Nefedov, N.N. (1977). Asimptoticheskaia teoriia kontrastnykh struktur [Asymptotic theory of contrast structures]. *Avtomatika i Telemekhanika – Automat. and Telemech.*, 7, 3–32 [in Russian].
- 7 Vasilyeva, A.B., Butuzov, V.F., & Nefedov, N.N. (1998). Kontrastnye struktury v sinhuliarno vozmushchennykh zadachakh [Contrast structures in singularly perturbed problems]. *Fundamentalnaia i prikladnaia matematika – Fundament. and Appl. Math.*, 4(3), 799–851 [in Russian].
- 8 Lomov, S.A., & Lomov, I.S. (2011). *Osnovy matematicheskoi teorii pohranichnogo sloia [Foundations of mathematical theory of boundary layer]*. Moscow: Izdatelstvo Moskovskogo gosudarstvennogo universiteta [in Russian].
- 9 Lomov, S.A., & Eliseev, A.G. (1986). Teoriia sinhuliarnykh vozmushchenii v sluchae spektralnykh osobennosti predelnogo operatora [The theory of singular perturbations in the case of spectral singularities of the limit operator]. *Matematicheskii sbornik – Math. collect.*, 131(173), 4(26), 544–557 [in Russian].
- 10 Bobodzhanov, A.A., Rashepkina, N.A., & Safonov, V.F. (1998). Rehuliarizatsiia optimalnykh sinhuliarno vozmushchennykh sistem s bystro meniaiushcheisia dempfiroiushchei funktsiiei [Regularization of optimal singularly perturbed systems with a rapidly changing damping function]. *Vestnik MEI – Vestnik MPEI*, 6, 18–26 [in Russian].
- 11 Imanbaev, N.S., Kalimbetov, B.T., & Temirbekov, M.A. (2013). Alhoritm metoda rehuliarizatsii dlia sinhuliarno vozmushchennoi zadachi s nestabilnym znacheniem yadra intehralnogo operatora [Algorithm the regularization method for a singularly perturbed problem with unstable value of kernel of the integral operator]. *Vestnik Karahandinskogo universiteta. Matematicheskaiia serii – Vestnik of Karaganda University, Mathematics series*, 4(72), 64–70 [in Russian].
- 12 Kalimbetov, B.T., Yeskaraeva, B.I., & Temirbekov, M.A. (2014). Matematicheskoe opisaniie vnutrenneho pohranichnogo sloia dlia nelineinoy intehro-differentsialnoi sistemy [Mathematical description of the inner boundary layer for a nonlinear integro-differential system]. *Vestnik Karahandinskogo universiteta. Serii matematika – Vestnik of Karaganda University. Mathematics series*, 3(75), 77–87 [in Russian].
- 13 Yeskaraeva, B.I., Kalimbetov, B.T., & Temirbekov, M.A. (2014). Diskretnyi pohranichnyi sloi v sluchae nulevykh toчек spektra dlia sistem intehro-differentsialnykh uravnenii [Discrete boundary layer in the case of zero points of the spectrum for systems of integro-differential equations]. *Vestnik Karahandinskogo universiteta. Serii matematika – Vestnik of Karaganda University, Mathematics series*, 75, 3, 88–95 [in Russian].
- 14 Kalimbetov, B.T., Yeskarayeva, B.I., & Tolep A.S. (2015). Internal boundary layer for integral-differential equations with zero spectrum of the limit operator and rapidly changing kernel. *Journal Applied Mathematical Sciences*, 9, 141–144, 7149–7165.
- 15 Kalimbetov, B.T., Temirbekov, M.A., & Habibullaev, Zh.O. (2012). Asymptotic solution of singular perturbed problems with an instable spectrum of the limiting operator. *Journal Abstract and Applied Analysis*, ID 120192.
- 16 Kalimbetov, B.T., Imanbaev, N.S., Sapakov, D.A., & Tashimov, L.T. (2013). Regularized asymptotical solutions of integro-differential systems with spectral singularities. *Journal Advances in Difference Equations*, 109, doi: 10.1186/1687-1847-2013-109.
- 17 Safonov, V.F., & Kalimbetov, B.T. (1995). A regularization method for systems with unstable spectral value of the kernel of the integral operator. *Journal Differential Equations*, 31(4), 647–656.

- 18 Safonov, V.F., & Bobodzhanov, A.A. (2017). Regularized asymptotic solutions of the initial problem of systems of integro-partial differential equations. *Mathematical Notes*, 102, 1, 22-30.
- 19 Safonov, V.F., & Bobodzhanov, A.A. (2018). Regularized asymptotics of solutions to integro-differential partial differential equations with rapidly varying kernels. *Ufmsk. Mat. Zh.*, 10, 2. 3-12.
- 20 Bobodzhanov, A.A., & Safonov, V.F. (2018). A generalization of the regularization method to the singularly perturbed integro-differential equations with partial derivatives. *Journal Russian Math. (Izvest. VUZ)*, 62, 3, 6-17.
- 21 Bobodzhanov, A.A., & Safonov, V.F. (2016). A problem with inverse time for a singularly perturbed integro-differential equation with diagonal - of the kernel of high order. *Journal Izvest. Math*, 80, 2, 285-298.
- 22 Bobodzhanov, A.A., & Safonov, V.F. (2015). Asymptotic solutions of Fredholm integro-differential equations with rapidly changing kernels and irreversible limit operator. *Journal Russian Math. (Izvest. vuz)*, 59, 10, 1-15.
- 23 Bobodzhanov, A.A., & Safonov, V.F. (2013). The method of normal forms for singularly perturbed systems of Fredholm integro-differential equations with rapidly varying kernels. *Sibir. Math. J.*, 204, 7, 979-1002.
- 24 Safonov, V.F., & Bobodzhanov, A.A. (2001). Volterra integral equations with rapidly varying kernels and their asymptotic integration. *Sibir. Math. J.*, 192, 8, 1139-1164.
- 25 Safonov, V.F., & Bobodzhanov, A.A. (2009). «Splashes» in Fredholm integro-differential equations with rapidly varying kernels. *Math. Notes*, 85, 2. 153-167.
- 26 Gantmakher, F.R. (1953). *Teoriia matrits [Theory of matrices]*. Moscow: HITTL [in Russian].
- 27 Konyaev, Yu.A. (1993). On a method of studying some problems of the perturbation theory. *Mathematicheskii sbornik - Mathem. collect.*, 12, 133-144 [in Russian].