

N.S. Imanbaev^{1,2}, M.N. Ospanov³¹South Kazakhstan State Pedagogical University, Shymkent, Kazakhstan;²Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan;³L.N. Gumilyov Eurasian National University, Nur-Sultan, Kazakhstan

(E-mail: imanbaevnur@mail.ru, myrzan66@mail.ru)

On eigenvalues of third order composite type equations with regular boundary value conditions

In the paper the question about distribution of eigenvalues of third-order composite type equations with regular, more precisely, with periodic boundary value conditions is studied. After, applying the Fourier method, the original problem splits into two problems on eigenvalues of third-order ordinary differential operators with periodic boundary value conditions in $L_2(0, 1)$. Characteristic determinants are calculated and zeros of entire analytic functions are found, and their location on the complex plane is determined. Existence of an infinite number of eigenvalues of a third order composite type operator is proved. Distance between the neighboring eigenvalues of the third order composite type operator of each series, which lie on rays, perpendicular to sides of a conjugate indicator diagram, that is, a regular hexagon on the complex plane, is determined. Moreover, it is determined that zero is not an eigenvalue of a third order composite type operator, in other words, zero is a regular point of the operator that belongs to resolvent set of the original operator. Adjoint operator with periodic boundary value conditions is constructed.

Keywords: composite type equations, regular, periodic boundary value conditions, rectangular domain, Fourier method, characteristic determinant, entire analytic functions, eigenvalue, zeros of entire functions.

Introduction and Formulation of the problem

Series of spectral boundary value problems for composite type equations

$$\frac{\partial}{\partial x}(u_{xx} + u_{yy}) + \lambda u = 0$$

has been investigated in [1, 2]. Solution of initial-boundary value problems for partial differential equations by the Fourier method is almost always reduced to the problem on determining eigenvalues and eigen-functions of some differential operators [3–6].

This paper is devoted to finding the eigenvalues of one boundary value problem for the equation

$$Lu \equiv u_{xxx} + u_{yyy} + \lambda u = 0, \quad (1)$$

where λ is a spectral parameter and complex number, which is also composite type equation [1].

In a rectangular domain D we consider the problem on eigenvalues of the equation (1), satisfying the following boundary conditions:

$$u|_{\partial D} = 0, \quad u_x(0, y) = u_x(1, y), \quad u_y(x, 0) = u_y(x, 1), \quad (2)$$

where $D = \{x, y : 0 < x < 1, 0 < y < 1\}$.

Solution of the problem

Looking for a solution of the problem (1), (2) by the Fourier method as follows:

$$u(x, y) = X(x) \cdot Y(y),$$

we come to the following spectral problems in the space $L_2(0, 1)$ for ordinary differential operators

$$L_0 X \equiv X''' + \mu X = 0, \quad X(0) = X(1) = 0, \quad X'(0) = X'(1), \quad (3)$$

$$L_1 Y \equiv Y''' + \nu Y = 0, \quad Y(0) = Y(1) = 0, \quad Y'(0) = Y'(1), \quad (4)$$

moreover, $\lambda = \mu + \nu$. Boundary value problems in (3),(4) are regular by G.D. Birkhoff [7]. In the monograph of M.A. Naimark [8; 67] a subclass of regular boundary conditions is distinguished, and it is noted that for an odd order of the equation, all strongly regular conditions are regular.

We solve the problem (3) (problem (4) is solved analogically). General solution of the equation (3) has the form

$$X(x) = C_1 e^{2ax} + \left(C_2 \cdot \cos \sqrt{3}ax + C_3 \cdot \sin \sqrt{3}ax \right) \cdot e^{-ax}, \quad (5)$$

where C_1, C_2, C_3 are arbitrary constants,

$$a = \frac{\sqrt[3]{-\mu}}{2} \neq 0. \quad (6)$$

Putting (5) into the boundary value condition (3), we will have the linear system concerning to the coefficients C_j :

$$\begin{cases} C_1 + C_2 = 0, \\ C_1 \cdot e^{2a} + C_2 \cdot e^{-a} \cdot \cos \sqrt{3}a + C_3 \cdot e^{-a} \cdot \sin \sqrt{3}a = 0, \\ C_1 \cdot (2a - 2a \cdot e^{2a}) + C_2 \cdot (-a + \sqrt{3}a \cdot e^{-a} \cdot \sin \sqrt{3}a + a \cdot e^{-a} \cdot \cos \sqrt{3}a) + \\ + C_3 \cdot (\sqrt{3}a - \sqrt{3}a \cdot e^{-a} \cdot \cos \sqrt{3}a + a \cdot e^{-a} \cdot \sin \sqrt{3}a) = 0. \end{cases}$$

Its determinant will be a characteristic determinant for the problem (3):

$$\Delta(a) = \begin{vmatrix} 1 & 1 & 0 \\ e^{2a} & e^{-a} \cos \sqrt{3}a & e^{-a} \sin \sqrt{3}a \\ 2a - 2ae^{2a} & ae^{-a} (\sqrt{3} \sin \sqrt{3}a + \cos \sqrt{3}a) - a & \sqrt{3}a - ae^{-a} (\sqrt{3} \cos \sqrt{3}a + \sin \sqrt{3}a) \end{vmatrix}. \quad (7)$$

From where by standard calculations and transformations the determinant (7) is reduced to the form:

$$\begin{aligned} \Delta(a) = & (\sqrt{3} + 3i) e^{(1+i\sqrt{3})a} + (\sqrt{3} - 3i) e^{(1-i\sqrt{3})a} + (\sqrt{3} + 3i) e^{-(1+i\sqrt{3})a} + \\ & + (\sqrt{3} - 3i) e^{-(1-i\sqrt{3})a} - 2\sqrt{3}e^{2a} - 2\sqrt{3}e^{-2a}. \end{aligned} \quad (8)$$

We formulate the obtained result as the following theorem.

Theorem 1. Characteristic determinant of the spectral problem (3) is represented as a form of quasi-polynomial (7) and is the entire analytic function of the variable a .

Connection of quasi-polynomials zeros with spectral problems is reflected in [9–13].

Sometimes entire analytical functions coincide with quasi-polynomials, zeros of which are investigated in [8, 14–18].

The papers [19, 20] are devoted to study of zeros of entire functions with an integral representation, related to spectral problems of a third-order differential operator with nonlocal boundary value conditions.

In [21, 22] the characteristic determinant of spectral problem for the Sturm-Liouville operator with perturbed regular boundary value conditions, which is an entire analytic function of the spectral parameter, is calculated. Also in this paper, they study stability problems of basis property of root functions systems of the original operator.

Zeros of the entire analytical function $\Delta(a)$ in (8) are eigenvalues of the operator L_0 . Therefore, further we consider the question about distribution of eigenvalues of the entire function $\Delta(a)$ on the complex plane a .

Taking into account results of the monographs [9, 10, 16], the conjugate indicator diagram of the function $\Delta(a)$ will be a regular hexagon on the complex plane a . Sides of the hexagon consist of the following segments:

$$\begin{aligned} & \left[1 - i\sqrt{3}; \overline{-1 - i\sqrt{3}} \right], \left[-1 + i\sqrt{3}; \overline{1 + i\sqrt{3}} \right], \left[-2; \overline{-1 - i\sqrt{3}} \right], \\ & \left[-1 + i\sqrt{3}; \overline{-2} \right], \left[1 + i\sqrt{3}; \overline{2} \right], \left[2; \overline{1 - i\sqrt{3}} \right], \end{aligned}$$

where lines mean complex conjugation, and they are commensurable numbers, that is, the length of each segment is equal to $d = 2$, and therefore they form the regular hexagon. From the origin we draw rays that are perpendicular to the sides of the regular hexagon.

Rays, which are perpendicular to the indicator diagram, are called critical. According to [10] the critical rays on the plane a are exactly six, that is

$$\arg \sqrt[3]{a} = \frac{\pi}{6} + \frac{\pi n}{3}, \quad n = 0, 1, 2, 3, 4, 5.$$

Along the ray, perpendicular to the segment passing through the points $\bar{2}$; $\overline{1 - i\sqrt{3}}$, there are zeros of the quasi-polynomial $(\sqrt{3} + 3i) \cdot e^{(1+i\sqrt{3})a} - 2\sqrt{3} \cdot e^{2a}$ from (8), which are majorizing exponents. Moreover, along this ray other exponents from (8) do not contribute.

We find zeros of the quasi-polynomial:

$$\begin{aligned} (\sqrt{3} + 3i) \cdot e^{(1+i\sqrt{3})a} - 2\sqrt{3} \cdot e^{2a} &= 0, \\ (\sqrt{3} + 3i) \cdot e^{(1+i\sqrt{3})a} &= 2\sqrt{3} \cdot e^{2ik\pi}, \\ a_{k1} &= \frac{2ik\pi}{-1 + i\sqrt{3}} + \frac{\ln \left| \frac{2\sqrt{3}}{\sqrt{3}+3i} \right| + i \operatorname{Arg} \left(\frac{2\sqrt{3}}{\sqrt{3}+3i} \right)}{-1 + i\sqrt{3}}, \quad k = 1, 2, 3, \dots, \end{aligned}$$

which are zeroes of the first series, where $\ln \left| \frac{2\sqrt{3}}{\sqrt{3}+3i} \right| + i \operatorname{Arg} \left(\frac{2\sqrt{3}}{\sqrt{3}+3i} \right) = \text{const}$.

Otherwise, from (7) it follows that eigenvalues of the first series of the operator L_0 will be

$$\mu_{k1} = - \left(\frac{4ik\pi}{-1 + i\sqrt{3}} + \frac{\text{const}}{-1 + i\sqrt{3}} \right)^3, \quad k = 1, 2, 3, \dots$$

A similar procedure is carried out on the other sides of the hexagon, and along the other perpendicular rays we have the corresponding series of quasi-polynomials zeros from (8):

– segment $\left[-1 - i\sqrt{3}; 1 - i\sqrt{3} \right]$, 2-nd series of zeros

$$\begin{aligned} a_{k2} &= \frac{ik\pi}{1 + i\sqrt{3}} + \frac{\text{const}}{2(1 + i\sqrt{3})}, \quad k = 1, 2, 3, \dots \\ \mu_{k2} &= - \left(\frac{2ik\pi}{1 + i\sqrt{3}} + \frac{\text{const}}{1 + i\sqrt{3}} \right)^3, \quad k = 1, 2, 3, \dots; \end{aligned}$$

– segment $\left[-1 + i\sqrt{3}; 1 + i\sqrt{3} \right]$, 3-rd series of zeros

$$\begin{aligned} a_{k3} &= ik\pi + \text{const}, \quad k = 1, 2, 3, \dots, \\ \mu_{k3} &= -(2ik\pi + \text{const})^3, \quad k = 1, 2, 3, \dots; \end{aligned}$$

– segment $\left[-2; -1 - i\sqrt{3} \right]$, 4-th series of zeros

$$\begin{aligned} a_{k4} &= \frac{2ik\pi}{1 + i\sqrt{3}} + \frac{\text{const}}{1 + i\sqrt{3}}, \quad k = 1, 2, 3, \dots, \\ \mu_{k4} &= - \left(\frac{2ik\pi}{1 + i\sqrt{3}} + \frac{\text{const}}{1 + i\sqrt{3}} \right)^3, \quad k = 1, 2, 3, \dots; \end{aligned}$$

– segment $\left[\bar{2}; \overline{1 + i\sqrt{3}} \right]$, 5-th series of zeros

$$\begin{aligned} a_{k5} &= - \frac{2ik\pi}{1 + i\sqrt{3}} - \frac{\text{const}}{1 + i\sqrt{3}}, \quad k = 1, 2, 3, \dots, \\ \mu_{k5} &= \left(\frac{2ik\pi}{1 + i\sqrt{3}} + \frac{\text{const}}{1 + i\sqrt{3}} \right)^3, \quad k = 1, 2, 3, \dots \end{aligned}$$

– segment $\left[\overline{-1 + i\sqrt{3}}; \overline{-2} \right]$, 6-th series of zeros

$$a_{k6} = \frac{2ik\pi}{1 - i\sqrt{3}} + \frac{const}{1 - i\sqrt{3}}, \quad k = 1, 2, 3, \dots,$$

$$\mu_{k6} = -\left(\frac{4ik\pi}{1 - i\sqrt{3}} + \frac{const}{1 - i\sqrt{3}} \right)^3, \quad k = 1, 2, 3, \dots$$

Thus,

Proposition 1.

1. There exists an infinite number of operator eigenvalue of the operator L_0 .
2. Distance between neighboring eigenvalues of the operator L_0 of the each series is equal to $\frac{2\pi}{|d|}$.
3. Eigenvalues of each series of the operator L_0 lie on the rays, perpendicular to the segment containing the numbers

$$\left(\overline{1 - i\sqrt{3}} \overline{-1 - i\sqrt{3}} \right), \left(\overline{-1 + i\sqrt{3}} \overline{1 + i\sqrt{3}} \right), \left(\overline{-2} \overline{-1 - i\sqrt{3}} \right), \left(\overline{-1 + i\sqrt{3}} \overline{-2} \right),$$

$$\left(\overline{1 + i\sqrt{3}} \overline{2} \right), \left(\overline{2} \overline{1 - i\sqrt{3}} \right).$$

Similarly, by repeating the whole process of researching the problem (3), we solve the problem (4), and we obtain eigenvalues of the operator L_1 :

$$\nu_{l1} = -\left(\frac{4il\pi}{-1 + i\sqrt{3}} + \frac{const}{-1 + i\sqrt{3}} \right)^3, \quad l = 1, 2, 3, \dots$$

$$\nu_{l2} = -\left(\frac{2il\pi}{1 + i\sqrt{3}} + \frac{const}{1 + i\sqrt{3}} \right)^3, \quad l = 1, 2, 3, \dots$$

$$\nu_{l3} = -(2il\pi + const)^3, \quad l = 1, 2, 3, \dots$$

$$\nu_{l4} = -\left(\frac{2il\pi}{1 + i\sqrt{3}} + \frac{const}{1 + i\sqrt{3}} \right)^3, \quad l = 1, 2, 3, \dots$$

$$\nu_{l5} = \left(\frac{2il\pi}{1 + i\sqrt{3}} + \frac{const}{1 + i\sqrt{3}} \right)^3, \quad l = 1, 2, 3, \dots$$

$$\nu_{l6} = -\left(\frac{4il\pi}{1 - i\sqrt{3}} + \frac{const}{1 - i\sqrt{3}} \right)^3, \quad l = 1, 2, 3, \dots$$

Analogically, all points of Proposition 1 are true for the operator L_1 .

So, we have proved the following:

Theorem 2. Suppose that all conditions of Theorem 1 and Proposition 1 hold for the operators L_0 and L_1 . Then eigenvalues of the operator L are $\lambda_{klj} = \pm(\mu_{kj} + \nu_{lj})$, where $k = 1, 2, 3, \dots, l = 1, 2, 3, \dots, j = \overline{1-6}$ mean the each series.

Remark. In the case $a = \frac{\sqrt[3]{-\mu}}{2} = 0$, representing general solution (3) as $X(x) = ax^2 + bx + c$ and satisfying the boundary value conditions in (3), we have $X(x) = 0$, that is, $\mu_0 = 0$ is not eigenvalue of the operator L_0 . Similarly, $\nu_0 = 0$ is a regular point of the operator L_1 . Thus, $\lambda_0 = 0$ is not eigenvalue of the operator L .

Conjugate problems

$L_0 X \equiv l_0(X) = X'''(x)$. Applying the method of integration by parts, we get the Lagrange formula:

$$\int_0^1 l_0(X) \overline{v(x)} dx + \int_0^1 X(x) \overline{l_0^*(v)} dx = X''(1) \overline{v(1)} - X''(0) \overline{v(0)} -$$

$$- \left[\overline{v'(0)} - \overline{v'(1)} \right] \cdot X'(0) + X(1) \cdot \overline{v''(1)} - X(0) \cdot \overline{v''(0)}.$$

Here $l_0^*(v)$ is the conjugate differential expression:

$$l_0^*(v) = -v'''(x), \quad 0 < x < 1. \tag{9}$$

Consequently, the operator оператор L_0^* , conjugate to the operator L_0 , is given by the differential expression (9) and the boundary value conditions

$$v(1) = v(0) = 0, \quad v'(0) - v'(1) = 0. \quad (10)$$

Analogically, for the operator L_1 conjugate operator is

$$L_1 Y \equiv l_1(Y) = Y''''(y), \quad L_1^* : l_1^*(v) = -v''''(y), \quad 0 < y < 1$$

with the boundary value conditions (10). From this, it follows that in the domain D conjugate problem to the problem (1), (2) will be

$$L^*V = V_{xxx} + V_{yyy} - \lambda V = 0,$$

satisfying the boundary value conditions

$$V|_{\partial D} = 0, \quad V_x(1, y) = V_x(0, y), \quad V_y(x, 0) = V_y(x, 1).$$

This study was supported by Committee of Science of the Ministry of Education and Sciences RK (Grant No. AP05132587).

References

- 1 Джураев Т.Д. О классификации и приведении к каноническому виду уравнений с частными производными третьего порядка / Т.Д. Джураев, Я. Попелек // Дифференциальные уравнения. — 1991. — Т. 27, № 10. — С. 1734–1745.
- 2 Джураев Т.Д. Вычисление собственных значений и собственных функций некоторых дифференциальных операторов третьего и четвертого порядков / Т.Д. Джураев, Б.В. Логинов, И.А. Малюгина // Дифференциальные уравнения математической физики и их приложения. — Ташкент: Фан, 1989. — С. 24–36.
- 3 Ильин В.А. Спектральная теория дифференциальных операторов. Разложение по фундаментальным функциям самосопряженных эллиптических операторов / В.А. Ильин. — М.: Наука, 1991. — 167 с.
- 4 Моисеев Е.И. Уравнения смешанного типа со спектральным параметром / Е.И. Моисеев. — М.: Изд-во МГУ, 1988. — 150 с.
- 5 Кальменов Т.Ш. О спектре задачи Трикоми для уравнения Лаврентьева — Бицадзе / Т.Ш. Кальменов // Дифференциальные уравнения. — 1977. — Т. 13, — № 8. — С.1418–1425.
- 6 Садыбеков М.А. Спектральные свойства одного класса краевых задач для парабола-гиперболического уравнения / М.А. Садыбеков, Г.Д. Тойжанова // Дифференциальные уравнения. — 1992. — Т. 28, № 1. — С. 176–179.
- 7 Birkhoff G.D. On the asymptotic character of the certain linear differential equations containing a parameter / G.D. Birkhoff // Transactions of the American Mathematical Society. — 1908. — Vol. 9, No. 2. — P. 219–231.
- 8 Наймарк М.А. Линейные дифференциальные операторы / М.А. Наймарк. — М.: Наука, 1969. — 496 с.
- 9 Беллман Р. Дифференциально-разностные уравнения / Р. Беллман, К. Кук. — М.: Наука, 1967. — 548 с.
- 10 Леонтьев А.Ф. Целые функции и ряды экспонент / А.Ф. Леонтьев. — М.: Наука, 1983. — 176 с.
- 11 Hald O.H. Discontinuous Inverse Eigen Value Problems / O.H. Hald // Communications on Pure Applied Mathematics. — 1984. — Vol. 37. — P. 539–577.
- 12 Левин Б.Я. Распределение корней целых функций / Б.Я. Левин. — М.: Наука, 1956. — 632 с.
- 13 Malamud M.M. On the completeness of root vectors of first order systems/ M.M. Malamud, L.L. Oridoroga // Doklady Mathematics. — 2010. — Vol. 82, No. 3. — P. 809–905.

- 14 Садовничий В.А. О нулях целых функций одного класса / В.А. Садовничий, В.А. Любишкин, Ю. Беллаббаси // Тр. семинара им. И.Г. Петровского. — 1982. — 8. — С. 211–217.
- 15 Шкалик А.А. О базисности собственных функций обыкновенных дифференциальных операторов с интегральными краевыми условиями / А.А. Шкалик // Вест. Москов. ун-та. Сер. 1. Математика. Механика. — 1982. — № 6. — С. 12–21.
- 16 Кангужин Б.Е. Дифференциальные операторы на отрезке. Распределение собственных значений / Б.Е. Кангужин, М.А. Садыбеков. — Шымкент: Ғылым, 1996. — 270 с.
- 17 Imanbaev N.S. Distribution of Eigen values of a Third-Order Differential Operator with Strongly Regular non local Boundary Conditions / N.S. Imanbaev // AIP Conference Proceedings. — 2018. — Vol. 1997. — P. 020027-1 – 020027-5.
- 18 Imanbaev N.S. On zeros of a Quasi-Polynomial of Exponential Type Connected with a Regular Third order Differential Operator / N.S. Imanbaev // Математ. журн. — 2018. — Т. 18, № 2. — С. 124–132.
- 19 Иманбаев Н.С. О нулях целых функций, имеющих интегральное представление / Н.С. Иманбаев, Б.Е. Кангужин // Изв. НАН РК. Сер. физ.-мат. — 1995. — № 3. — С. 47–52.
- 20 Imanbaev N.S. On zeros the characteristic determinant of the spectral problem for a third-order differential operator on a segment with nonlocal boundary conditions / N.S. Imanbaev, B.E. Kanguzhin, B.T. Kalimbetov // Advances in Difference Equations. — 2013. — Doi:10.1186/1687-1847-20113-110.
- 21 Sadybekov M.A. Characteristic determinant of a boundary value problem, which does not have the basis property / M.A. Sadybekov, N.S. Imanbaev // Eurasian Mathematical Journal. — 2017. — 8, 2. — P. 40–46.
- 22 Иманбаев Н.С. Характеристический определитель спектральной задачи для оператора Штурма-Лиувилля с интегральным возмущением краевого условия антипериодического типа / Н.С. Иманбаев // Вестн. Караганд. ун-та. Сер. Математика. — 2016. — № 2(82). — С. 68–72.

Н.С. Иманбаев, М.Н. Оспанов

Регулярлық шеттік шарттармен берілген үшінші ретті құрамдас типті теңдеудің меншікті мәндері жайлы

Мақалада регулярлық, дәлірек айтқанда, периодтық шеттік шарттармен берілген үшінші ретті құрамдас типті теңдеудің меншікті мәндерінің орналасуы туралы мәселе зерттелді. Бастапқы есепті Фурье әдісімен шешуді қолданғаннан кейін, $L_2(0, 1)$ кеңістігінде периодтық шарттармен берілген үшінші ретті жай дифференциалдық теңдеудің меншікті мәндерін зерттеуге арналған екі есепке тармақталған. Осы есептердің сипаттамалық анықтауыштарының бүтін аналитикалық функциялар болатындығы дәлелденіп, олардың нөлдері табылып, кешенді жазықтықтағы орындары анықталған. Бастапқы оператордың меншікті мәндерінің саналымды, шексіз екендігі көрсетілген. Түйіндес индикаторлық диаграммасы құрылып, әр сериядағы меншікті мәндердің перпендикуляр сәулелердің бойында арақашықтықтары анықталған. Спектралдық параметрдің нөлдік мәні оператордың меншікті мәні болмайтындығы көрсетілген. Түйіндес операторы құрылған.

Кілт сөздер: құрамдас типті теңдеу, регулярлық периодтық шеттік шарттар, төртбұрыш аймақ, Фурье әдісі, характеристикалық анықтауыш, бүтін аналитикалық функция, меншікті мәндер, бүтін функцияның нөлдері.

О собственных значениях уравнений третьего порядка составного типа с регулярными краевыми условиями

В статье исследован вопрос распределения собственных значений уравнений третьего порядка составного типа с регулярными, точнее, с периодическими краевыми условиями. После применения метода Фурье исходная задача распадается на две задачи, т.е. на собственные значения обыкновенных дифференциальных операторов третьего порядка с периодическими краевыми условиями в $L_2(0, 1)$. Вычислены характеристические определители и найдены нули целых аналитических функций и определено их расположение на комплексной плоскости. Доказано существование бесконечного числа собственных значений оператора третьего порядка составного типа. Определено расстояние между соседними собственными значениями оператора третьего порядка составного типа каждой серии, которое лежит на лучах перпендикулярно сторонам сопряженной индикаторной диаграммы, то есть правильного шестиугольника на комплексной плоскости. Доказано, что нуль не является собственным значением оператора третьего порядка составного типа, иначе говоря, нуль является регулярной точкой оператора, которая принадлежит резольвентному множеству исходного оператора. Построен сопряженный оператор с периодическими краевыми условиями.

Ключевые слова: уравнения составного типа, регулярные, периодические краевые условия, прямоугольная область, метод Фурье, характеристический определитель, целые аналитические функции, собственные значения, нули целых функций.

References

- 1 Djuraev, T.D. & Popelek, Ya. (1991). O klassifikatsii i privedenii k kanonicheskomu vidu uravnenii s chastnymi proizvodnymi tretieho poriadka [On classification and reduction to the canonical form of third-order partial differential equations]. *Differentsialnye uravnenia – Differential Equations*, 27, 10, 1734–1745 [in Russian].
- 2 Djuraev, T.D., Loginov, B.V. & Malugina, I.A. (1989). Vychislenie sobstvennykh znachenii i sobstvennykh funktsii nekotorykh differentsialnykh operatorov tretieho i chetvertogo poriadkov [Calculation of eigenvalues and eigen-functions of some differential operators of the third and fourth orders]. *Differentsialnye uravnenia matematicheskoi fiziki i ikh prilozhenia – Differential equations of mathematical physics and their applications*, 24–36. Tashkent: Fan [in Russian].
- 3 Iliin, V.A. (1991). *Spektralnaya teoriya differentsialnykh operatorov. Razlozhenie po fundamentalnym funktsiam samosopriazhennykh ellipticheskikh operatorov* [Spectral theory of differential operators. Expansion by fundamental functions of self-adjoint elliptic operators]. Moscow: Nauka [in Russian].
- 4 Moiseev, E.I. (1988). *Uravnenia smeshannogo tipa so spektralnym parametram* [Mixed type equations with spectral parameters]. Moscow: Isdatelstvo MHU [in Russian].
- 5 Kalmenov, T.Sh. (1977). O spektre zadachi Triкоми dlia uravnenia Lavrentieva-Bitsadze [On spectrum of the Tricomi problem for the Lavrent'ev-Bitsadze equation]. *Differentsialnye uravnenia – Differential Equations*, 13, 8, 1418–1425 [in Russian].
- 6 Sadybekov, M.A. & Toizhanova, G.D. (1992). Spektralnye svoistva odnogo klassa kraevykh zadach dlia parabolno-giperbolicheskogo uravnenia [Spectral properties of a class of boundary value problems for a parabolic-hyperbolic equation]. *Differentsialnye uravnenia – Differential Equations*, 28, 1, 176–179 [in Russian].
- 7 Birkhoff, G.D. (1908). On asymptotic character of the certain linear differential equations containing a parameter. *Transactions of the American Mathematical Society*, 9, 2, 219–231.
- 8 Naimark, M.A. (1969). *Lineinye differentsialnye operatory* [Linear differential operators]. Moscow: Nauka [in Russian].
- 9 Bellman, R. & Kuk, K. (1967). *Differentsialno-raznostnye uravnenia* [Differential-difference Equations]. Moscow: Nauka [in Russian].
- 10 Leontiev, A.F. (1983). *Tselye funktsii i riady eksponent* [Entire Functions and Exponential Series]. Moscow: Nauka [in Russian].

- 11 Hald, O.H. (1984). Discontinuous Inverse Eigen Value Problems. *Communications on Pure Applied Mathematics*, 37, 539–577.
- 12 Levin, B.Ia. (1956). *Raspredelenie kornei tselykh funktsii [Distribution of roots of entire functions]*. Moscow: Nauka [in Russian].
- 13 Malamud, M.M. & Oridoroga, L.L. (2010). On completeness of root vectors of first order systems. *Doklady Akademii nauk. Matematika. Doklady Mathematics*, 82, 3, 809–905.
- 14 Sadovnichii, V.A., Liubishkin, V.A. & Belabbasi, U. (1982). O nuliakh tselykh funktsii odnogo klassa [On zeros of entire functions of one class]. *Trudy seminara imeni I.G. Petrovskoho – Proceedings of the I.G. Petrovsky seminar*, 8, 211–217 [in Russian].
- 15 Shkalikov, A.A. (1982). O bazisnosti sobstvennykh funktsii obyknovennykh differentsialnykh operatorov s intehralnymi kraevymi usloviami [On basis property of eigen-functions of ordinary differential operators with integral boundary value conditions]. *Vestnik Moskovskoho universiteta. Seria 1. Matematika. Mekhanika – Moscow University Mathematics. Mechanics Bulletin*, 6, 12–21 [in Russian].
- 16 Kanguzhin, B.E. & Sadybekov, M.A. (1996). *Differentsialnye operatory na otrezke. Raspredelenie sobstvennykh znachenii. [Differential operators on a segment. Distribution of eigenvalues]*. Shymkent: Hylym [in Russian].
- 17 Imanbaev, N.S. (2018). Distribution of eigenvalues of a third-order differential operator with strongly regular non local boundary conditions. *AIP Conference Proceedings*, 1997: 020027-1 – 020027-5.
- 18 Imanbaev, N.S. (2018). On zeros of an exponential type quasi-polynomial connected with a regular third order differential operator. *Matematicheskii zhurnal – Mathematical Journal*, 18, 2, 124–132.
- 19 Imanbaev, N.S. & Kanguzhin, B.E. (1995). O nuliakh tselykh funktsii, imeiushikh intehralnoe predstavlenie [On zeros of entire functions, having an integral representation]. *Izvestia NAN RK. Seria fiziko-matematicheskaya – News of NAS RK. Physics. Mathematics Series*, 3, 47–52 [in Russian].
- 20 Imanbaev, N.S., Kanguzhin, B.E. & Kalimbetov, B.T. (2013). On zeros of characteristic determinant of spectral problem for a third-order differential operator on a segment with nonlocal boundary conditions. *Advances in Difference Equations*, doi:10.1186/1687-1847-20113-110.
- 21 Sadybekov, M.A. & Imanbaev, N.S. (2017). Characteristic determinant of a boundary value problem, which does not have basis property. *Eurasian Mathematical Journal*, 8, 2, 40–46.
- 22 Imanbaev, N.S. (2016). Harakteristicheskii opredelitel spektralnoi zadachi dlia operatora Shturma-Liuvillia s intehralnym vozmusheniem kraevogo uslovia antiperiodicheskogo tipa [Characteristic determinant of the spectral problem for the Sturm-Liouville operator with the perturbed boundary value conditions]. *Vestnik Karahandinskoho universiteta. Seria Matematika – Bulletin of the Karaganda University. Mathematics*, 2 (82), 68–72 [in Russian].