

N.S. Imanbaev<sup>1,2,\*</sup>, Ye. Kurmysh<sup>1</sup><sup>1</sup>South Kazakhstan State Pedagogical University, Shymkent, Kazakhstan;<sup>2</sup>Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan

(E-mail: imanbaevnur@mail.ru, kurmysh2212@mail.ru)

## On zeros of an entire function coinciding with exponential type quasi-polynomials, associated with a regular third-order differential operator on an interval

In this paper, we consider the question on study of zeros of an entire function of one class, which coincides with quasi-polynomials of exponential type. Eigenvalue problems for some classes of differential operators on a segment are reduced to a similar problem. In particular, the studied problem is led by the eigenvalue problem for a linear differential equation of the third order with regular boundary value conditions in the space  $W_2^3(0, 1)$ . The studied entire function is adequately characteristic determinant of the spectral problem for a third-order linear differential operator with periodic boundary value conditions. An algorithm to construct a conjugate indicator diagram of an entire function of one class is indicated, which coincides with exponential type quasi-polynomials with comparable exponents according to the monograph by A.F. Leontyev. Existence of a countable number of zeros of the studied entire function in each series is proved, which are simultaneously eigenvalues of the above-mentioned third-order differential operator with regular boundary value conditions. We determine distance between adjacent zeros of each series, which lies on the rays perpendicular to sides of the conjugate indicator diagram, that is a regular hexagon on the complex plane. In this case, zero is not an eigenvalue of the considered operator, that is, zero is a regular point of the operator. Fundamental difference of this work is finding the corresponding eigenfunctions of the operator. System of eigenfunctions of the operator corresponding in each series is found. Adjoint operator is constructed.

*Keywords:* entire function, zeros, quasi-polynomials, indicator diagram, series, operator, regular periodic boundary value conditions, eigenvalues, system of eigenfunctions.

### Introduction and Formulation of the problem

We consider the question on distribution of zeros of an entire function of the following form:

$$\Delta(\lambda) = \sqrt[3]{\lambda}((k_2 - k_3)e^{k_1 \sqrt[3]{\lambda}} + (k_1 - k_2)e^{(k_2+k_1) \sqrt[3]{\lambda}} + (k_3 - k_1)e^{k_2 \sqrt[3]{\lambda}} + (k_3 - k_1)e^{(k_3+k_1) \sqrt[3]{\lambda}} + (k_1 - k_2)e^{k_3 \sqrt[3]{\lambda}} + (k_2 - k_3)e^{(k_2+k_3) \sqrt[3]{\lambda}}),$$

where  $k_1 = 1$ ,  $k_2 = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$ ,  $k_3 = -\frac{1}{2} - i\frac{\sqrt{3}}{2}$ .

Eigenvalue problems for some classes of differential operators on a segment are reduced to a similar problem. In particular, the following problem on eigenvalues in the space  $W_2^3(0, 1)$  leads to the studied question:

$$L_0 u \equiv l(u) = u'''(x) = -\lambda u(x), \quad 0 < x < 1, \quad (1)$$

$$U_1(u) = u(0) = 0, \quad U_2(u) = u(1) = 0, \quad U_3(u) = u'(0) = u'(1), \quad (2)$$

where  $U_1(u)$ ,  $U_2(u)$ ,  $U_3(u)$  are linear forms, which are regular, according to J.D. Birkhoff [1, 2]. An important result established by Birkhoff was to estimate resolvent of a regular differential operator and to establish asymptotics of the spectrum. In the monograph by M.A. Naimark [3; 67], a subclass of regular boundary conditions, so-called strongly regular boundary conditions, was singled out, where it was noted that for an odd order of the equation all regular conditions are strongly regular.

Connection between zeros of quasi-polynomials and spectral problems was reflected in [3–15]. Zeros of entire functions having an integral representation were studied in [16–23].

\*Corresponding author.

E-mail: imanbaevnur@mail.ru

Main Results

We consider the question on distribution of zeros of the entire function  $\Delta_1(\lambda) = \frac{\Delta(\lambda)}{\sqrt[3]{\lambda}}$  on the complex plane  $\lambda$ .

$$\Delta_1(\lambda) = (k_2 - k_3)e^{k_1 \sqrt[3]{\lambda}} + (k_1 - k_2)e^{(k_2+k_1) \sqrt[3]{\lambda}} + (k_3 - k_1)e^{k_2 \sqrt[3]{\lambda}} + (k_3 - k_1)e^{(k_3+k_1) \sqrt[3]{\lambda}} + (k_1 - k_2)e^{k_3 \sqrt[3]{\lambda}} + (k_2 - k_3)e^{(k_2+k_3) \sqrt[3]{\lambda}} = 0. \tag{3}$$

In [11, 14] the following was proved:

Proposition 1.

1. There are infinitely many zeros of an entire function  $\Delta_1(\lambda)$ ;
2. Distance between two adjacent zeros of the same series ( $j - const$ ) is exactly  $\frac{2\pi}{|d|}$ ;
3. Zeros of each series lie on the rays perpendicular to the segment, that is, perpendicular to sides of the hexagon containing

$$(\overline{k_1, (k_3 + k_1)}); (\overline{k_3, k_3 + k_1}); (\overline{k_2 + k_3, k_3}); (\overline{k_2, k_2 + k_3}); (\overline{k_2, k_2 + k_1}); (\overline{k_2 + k_1, k_1}).$$

The rays which are perpendicular to the indicator diagram are called critical. According to the result of the monograph [6], there are exactly six critical rays on the plane  $\lambda$ , that is  $arg \sqrt[3]{\lambda} = \frac{\pi}{6} + \frac{\pi n}{3}, n = 0, 1, 2, 4, 5$ ;

In [11, 14] the zeros of the entire function  $\Delta(\lambda)$ :

$$\lambda_{jk} = \frac{(\ln|z_j| + i(Arg(z_j) + 2\pi k))^3}{d^3}, \quad k \in \{0, \pm 1, \pm 2, \dots\}; \quad j = \overline{1, m} \tag{4}$$

were found, and conjugate indicator diagram-hexagon was constructed on the complex plane  $\lambda$ .

Taking  $k_1 = 1, k_2 = -\frac{1}{2} + i\frac{\sqrt{3}}{2}, k_3 = -\frac{1}{2} - i\frac{\sqrt{3}}{2}$ , into account, due to the formula (4) and Proposition 1, we have that along the ray perpendicular to the segment passing through the points  $\overline{2}; 1 - i\sqrt{3}$  there are zeros of the quasi-polynomial  $(\sqrt{3} + 3i) \cdot e^{(1+i\sqrt{3})\lambda} - 2\sqrt{3} \cdot e^{2\lambda}$ , they are majorizing exponents. In this case, other exponents from (3) do not contribute along this ray. Let's find zeros of the quasi-polynomial:

$$(\sqrt{3} + 3i) \cdot e^{(1+i\sqrt{3})\lambda} - 2\sqrt{3} \cdot e^{2\lambda} = 0$$

$$(\sqrt{3} + 3i) \cdot e^{(1+i\sqrt{3})\lambda} = 2\sqrt{3} \cdot e^{2\lambda}$$

$$\lambda_{k1} = \frac{2ik\pi}{-1 + i\sqrt{3}} + \frac{\ln|\frac{2\sqrt{3}}{\sqrt{3}+3i}| + iarg(\frac{2\sqrt{3}}{\sqrt{3}+3i})}{-3 + i\sqrt{3}}, \quad k = 1, 2, 3, \dots$$

Which are zeroes of the first series, where  $\ln|\frac{2\sqrt{3}}{\sqrt{3}+3i}| + iArg(\frac{2\sqrt{3}}{\sqrt{3}+3i}) = const$ . Similar procedure is performed on the other sides of the hexagon, and along other perpendicular rays we have the corresponding series of zeros of the quasi-polynomials from (3):

- segment  $[-1 - i\sqrt{3}; 1 - i\sqrt{3}]$ , 2-nd series of zeroes  $\lambda_{k2} = \frac{ik\pi}{1+i\sqrt{3}} + \frac{const}{2(1+i\sqrt{3})}, \quad k = 1, 2, \dots, (1 + i\sqrt{3})$
- segment  $[-1 + i\sqrt{3}; 1 + i\sqrt{3}]$ , 3-rd series of zeroes  $\lambda_{k3} = ik\pi + const, \quad k = 1, 2, \dots,$
- segment  $[-2; -1 - i\sqrt{3}]$ , 4-th series of zeroes  $\lambda_{k4} = \frac{2ik\pi}{1+i\sqrt{3}} + \frac{const}{1+i\sqrt{3}}, \quad k = 1, 2, \dots,$
- segment  $[\overline{2}; 1 + i\sqrt{3}]$ , 5-th series of zeroes  $\lambda_{k5} = -\frac{2ik\pi}{1+i\sqrt{3}} - \frac{const}{1+i\sqrt{3}}, \quad k = 1, 2, \dots,$
- segment  $[-1 + i\sqrt{3}; -2]$ , 6-th series of zeroes  $\lambda_{k6} = \frac{2ik\pi}{1+i\sqrt{3}} + \frac{const}{2(1+i\sqrt{3})}, \quad k = 1, 2, \dots,$

The zeros that were found are adequately eigenvalues of the operator  $L_0$  [11].

Fundamental difference of this section from [11, 14, 22, 23] is the determination of eigenfunctions of the operator  $L_0$ . The following theorem takes place.

Theorem. Let the entire function  $\Delta_1(\lambda) = \frac{\Delta(\lambda)}{\sqrt[3]{\lambda}}$  in (3), according to [11, 14], be a characteristic polynomial of the spectral problem (1), (2) and all points of Proposition 1. be satisfied, as well as zeros of the characteristic polynomial (4) be the corresponding eigenvalues of the operator  $L_0$ . Then the system of eigenfunctions of the operator  $L_0$  of each series:

$$\begin{aligned}
 u_{k1}(x) = & C_1 e^{-2\frac{k\sqrt{3}}{2}x} \cdot e^{2\frac{\pi}{4\sqrt{3}}x} \left[ \cos 2 \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x - i \sin 2 \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x \right] + \left\{ C_2 \left[ \cos \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x \cdot \cos \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{ch} \sqrt{3} \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x - \sin \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x \cdot \sin \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x - i \left( \cos \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{ch} \sqrt{3} \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x - \frac{\pi}{6} \right) x \sin \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x + \cos \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x \cdot \sin \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x - \frac{\pi}{6} \right) x \right] + C_3 \left[ \cos \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x \cdot \sin \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{ch} \sqrt{3} \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x - \sin \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x \cdot \cos \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x - i \left( \sin \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x \cdot \sin \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{ch} \sqrt{3} \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x + \cos \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x \cdot \cos \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x \right] \right\} e^{-\frac{k\sqrt{3}}{2}x} \cdot e^{\frac{\pi}{4\sqrt{3}}x};
 \end{aligned}$$

$$\begin{aligned}
 u_{k2}(x) = & C_1 e^{k\sqrt{3}x} \cdot e^{-\frac{\pi}{4}x} \left[ \cos \left( \frac{k\pi}{2} - \frac{\pi}{12} \right) x + i \sin \left( \frac{k\pi}{2} - \frac{\pi}{12} \right) x \right] + \left\{ C_2 \left[ \cos \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x \cdot \cos \sqrt{3} \left( \frac{k\pi\sqrt{3}}{2} - \frac{\pi}{8} x \right) \cdot \operatorname{ch} \sqrt{3} \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x + \sin \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x \cdot \sin \sqrt{3} \left( \frac{k\pi\sqrt{3}}{2} - \frac{\pi}{8} x \right) \cdot \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x - \frac{\pi}{8} \right) x \cdot \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x - i \left( \cos \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x \cdot \sin \sqrt{3} \left( \frac{k\pi\sqrt{3}}{2} - \frac{\pi}{8} x \right) x \cdot \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x - \frac{\pi}{24} \right) x - \sin \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x \cdot \cos \sqrt{3} \left( \frac{k\pi\sqrt{3}}{2} - \frac{\pi}{8} x \right) x \cdot \operatorname{ch} \sqrt{3} \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x \right] + C_3 \left[ \cos \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x \cdot \sin \sqrt{3} \left( \frac{k\pi\sqrt{3}}{2} - \frac{\pi}{8} x \right) \cdot \operatorname{ch} \sqrt{3} \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x + \sin \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x \cdot \cos \sqrt{3} \left( \frac{k\pi\sqrt{3}}{2} - \frac{\pi}{8} x \right) x \cdot \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x - i \left( \cos \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x \cos \sqrt{3} \left( \frac{k\pi\sqrt{3}}{2} - \frac{\pi}{8} x \right) x \cdot \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x + \sin \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x \cdot \sin \sqrt{3} \left( \frac{k\pi\sqrt{3}}{2} - \frac{\pi}{8} x \right) \cdot \operatorname{ch} \sqrt{3} \left( \frac{k\pi}{4} - \frac{\pi}{24} \right) x - \frac{\pi}{24} \right) x \right] \right\} e^{\frac{k\pi\sqrt{3}}{2}x} \cdot e^{-\frac{\pi}{8}x};
 \end{aligned}$$

$$\begin{aligned}
 u_{k3}(x) = & C_1 \left( \cos 2 \left( k\pi - \frac{\pi}{3} \right) x + i \sin 2 \left( k\pi - \frac{\pi}{3} \right) x \right) + \left[ C_2 \operatorname{ch} \sqrt{3} \left( k\pi - \frac{\pi}{3} \right) x \cdot \cos \left( k\pi - \frac{\pi}{3} \right) x + C_3 \operatorname{sh} \sqrt{3} \left( k\pi - \frac{\pi}{3} \right) x \cdot \sin \left( k\pi - \frac{\pi}{3} \right) x + i \left( C_3 \operatorname{sh} \sqrt{3} \left( k\pi - \frac{\pi}{3} \right) x \cdot \right. \right.
 \end{aligned}$$

$$\begin{aligned}
 & \cdot \cos \left( k\pi - \frac{\pi}{3} \right) x - C_2 \operatorname{ch} \sqrt{3} \left( k\pi - \frac{\pi}{3} \right) x \cdot \sin \left( k\pi - \frac{\pi}{3} \right) x \Big]; \\
 u_{k4}(x) = & C_1 e^{2\frac{k\pi\sqrt{3}}{2}x} \cdot e^{-2\frac{\pi}{12}x} \left[ \cos 2 \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x + i \sin 2 \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x \right] + \left\{ C_2 \left[ \cos \left( \frac{k\pi}{2} + \right. \right. \right. \\
 & \left. \left. \left. + \frac{\pi\sqrt{3}}{12} \right) x \cdot \cos \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{ch} \sqrt{3} \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x - \sin \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x \cdot \right. \right. \\
 & \left. \left. \sin \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x - i \left( \sin \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x \cdot \cos \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \right. \right. \right. \right. \\
 & \left. \left. \left. + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{ch} \sqrt{3} \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x + \cos \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x \cdot \sin \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \right. \right. \\
 & \left. \left. \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x \right] + C_3 \left[ \cos \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x \cdot \sin \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{ch} \sqrt{3} \left( \frac{k\pi}{2} - \right. \right. \right. \\
 & \left. \left. \left. - \frac{\pi}{6} \right) x - \sin \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x \cdot \cos \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x - i \left( \sin \left( \frac{k\pi}{2} - \right. \right. \right. \\
 & \left. \left. \left. - \frac{\pi}{6} \right) x \cdot \sin \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{ch} \sqrt{3} \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x + \cos \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x \cdot \cos \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \right. \right. \right. \\
 & \left. \left. \left. + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{2} - \frac{\pi}{6} \right) x \right] \right\} e^{-\frac{k\pi\sqrt{3}}{2}x} \cdot e^{\frac{\pi}{12}x}; \\
 u_{k5}(x) = & C_1 e^{-2\frac{k\pi\sqrt{3}}{2}x} \cdot e^{2\frac{\pi}{12}x} \left[ \cos 2 \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x + i \sin 2 \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x \right] + \\
 & + \left\{ C_2 \left[ \cos \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x \cdot \cos \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{ch} \sqrt{3} \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x + \right. \right. \\
 & + \sin \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x \sin \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x + i \left( \sin \left( \frac{k\pi}{2} + \right. \right. \\
 & \left. \left. + \frac{\pi\sqrt{3}}{12} \right) x \cdot \cos \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{ch} \sqrt{3} \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x - \right. \\
 & \left. - \cos \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x \right) \sin \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x \right] - C_3 \left[ \cos \left( \frac{k\pi}{2} + \right. \right. \\
 & \left. \left. + \frac{\pi\sqrt{3}}{12} \right) x \cdot \sin \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{ch} \sqrt{3} \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x + \sin \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x \cdot \right. \\
 & \left. \cos \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{sh} \sqrt{3} \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x + i \left( \sin \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x \sin \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \right. \right. \right. \\
 & \left. \left. \left. + \frac{\pi}{4\sqrt{3}} \right) x \cdot \operatorname{ch} \sqrt{3} \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x - \cos \left( \frac{k\pi}{2} + \frac{\pi\sqrt{3}}{12} \right) x \cdot \cos \sqrt{3} \left( -\frac{k\sqrt{3}}{2} + \frac{\pi}{4\sqrt{3}} \right) x \cdot \right. \right.
 \end{aligned}$$



## Acknowledgments

The first author was supported by a grant № AP 09260752 the Science Committee of the Ministry of Education and Science of the Republic of Kazakhstan.

## References

- 1 Birkhoff G.D. On the asymptotic character of the solutions of certain linear differential equations containing a parameter / G.D. Birkhoff // Transactions of the American Mathematical Society. — 1908. — 9. — No. 2. — P. 219–231.
- 2 Birkhoff G.D. Boundary value and expansion problems of ordinary linear differential equations / G.D. Birkhoff // Transactions of the American Mathematical Society. — 1908. — 9. — No. 4. — P. 373–395.
- 3 Наймарк М.А. Линейные дифференциальные операторы / М.А. Наймарк. — М.: Наука, 1969. — 496 с.
- 4 Titchmarsh E.C. The zeros of certain integral functions / E.C. Titchmarsh // Proc. London Math. Soc. — 1926. — 25. — No. 4. — P. 283–302.
- 5 Беллман Р. Дифференциально-разностные уравнения / Р. Беллман, К. Кук. — М., 1967. — 548 с.
- 6 Леонтьев А.Ф. Целые функции и ряды экспонент / А.Ф. Леонтьев. — М., 1983. — 176 с.
- 7 Hald O.H. Discontinuous Inverse Eigen Value Problems / O.H. Hald // Communications on Pure Applied Mathematics. — 1984. — XXXVII. — P. 539–577.
- 8 Лидский В.Б. Регуляризованные суммы корней одного класса целых функций / В.Б. Лидский, В.А. Садовничий // Функциональный анализ. — 1967. — 1. — № 2. — С. 52–59.
- 9 Седлецкий А.М. Когда все нули целой функций экспоненциального типа лежат в криволинейной полуплоскости (необходимое условие) / А.М. Седлецкий // Мат. сб. — 1995. — 186. — № 9. — С. 125–134.
- 10 Шкаликов А.А. О базисности собственных функций обыкновенных дифференциальных операторов с интегральными краевыми условиями / А.А. Шкаликов // Вестн. МГУ. Сер. Мат. мех. — 1982. — № 6. — С. 12–21.
- 11 Imanbaev N.S. Distribution of Eigen values of a Third-Order Differential Operator with Strongly Regular non local Boundary Conditions / N.S. Imanbaev // AIP Conference Proceedings. — 2018. — 1997. — 020027. — P. 020027-1 – 020027-5.
- 12 Садовничий В.А. О регуляризованных суммах корней целой функции одного класса / В.А. Садовничий, В.А. Любишкин, Ю. Белаббаси // Докл. АН СССР. — 1980. — 254. — № 6. — С. 1346–1348.
- 13 Садовничий В.А. О нулях целых функций одного класса / В.А. Садовничий, В.А. Любишкин, Ю. Белаббаси // Тр. семин. им. И.Г. Петровского. — 1982. — Вып. 8. — С. 211–217.
- 14 Imanbaev N.S. On zeros of a quasi-polynomial of exponential type connected with a regular third order differential operator / N.S. Imanbaev // Mathematical Journal. — 2018. — 18. — No. 2(68). — P. 124–132.
- 15 Левин Б.Я. Распределение корней целых функций / Б.Я. Левин. — М., 1956. — 632 с.
- 16 Седлецкий А.М. О нулях преобразования Фурье финитной меры / А.М. Седлецкий // Мат. заметки. — 1993. — 53. — № 1. — С. 111–119.
- 17 Кангужин Б.Е. Дифференциальные операторы на отрезке. Распределение собственных значений / Б.Е. Кангужин, М.А. Садыбеков. — Шымкент: Гылым, 1996. — 270 с.
- 18 Иманбаев Н.С. О нулях целых функций, имеющих интегральное представление / Н.С. Иманбаев, Б.Е. Кангужин // Изв. НАН Республики Казахстан. Сер. физ.-мат. — 1995. — № 3. — С. 47–52.
- 19 Imanbaev N.S. On zeros the characteristic determinant of the spectral problem for a third-order differential operator on a segment with nonlocal boundary conditions / N.S. Imanbaev, B.E. Kanguzhin, B.E. Kalimbetov // Advances in Difference Equations. — 2013. doi: 10.1186/1687-1847-20113-110.
- 20 Lunyov A.A. On the completeness and Riesz basis property of root subspaces of boundary value problems for first order systems and applications / A.A. Lunyov, M.M. Malamud // Journal of Spectral Theory. — 2015. — 5. — P. 17–70. doi 10.4171/jst/90
- 21 Sadybekov M.A. A Regular Differential Operator with Perturbed Boundary Condition / M.A. Sadybekov, N.S. Imanbaev // Mathematical Notes. — 2017. — 101. — № 5. — P. 878–887.

- 22 Imanbaev N.S. On eigenvalues of third order composite type equations with regular boundary value conditions / N.S. Imanbaev, M.N. Ospanov // Bulletin of the Karaganda University. Mathematics series. — 2019. — № 4(96). — P. 44–51. doi 10.31489/2019M4/44-51
- 23 Иманбаев Н.С. Распределение собственных значений уравнений третьего порядка составного типа с регулярными краевыми условиями / Н.С. Иманбаев, М.Н. Оспанов // Бюлл. Ин-та мат. — Ташкент, 2019. — № 5. — С. 28–33. doi: <http://mib.mathinst.UZ>
- 24 Makin A.S. On a nonlocal perturbation of a periodic eigenvalue problem / A.S. Makin // Differential Equations. — 2006. — 42. — № 4. — P. 599–602.
- 25 Kritskov L.V. Nonlocal spectral problem for a second-order differential equation with an involution / L.V. Kritskov, M.A. Sadybekov, A.M. Sarsenbi // Bulletin of the Karaganda University. Mathematics series. Special issue. — 2018. — № 3(91). — P. 53–61.
- 26 Baranetsky Ya.O. Nonlocal Multipoint Problem with Multiple Spectrum for an Ordinary (2n) Order Differential Equation / Ya.O. Baranetsky, P.I. Kalenyuk // Journal of Mathematical Sciences. — 2020. — 246. — P. 152–169.
- 27 Imanbaev N.S. On stability of the basis property of root vectors system of the Sturm-Liouville operator with an integral perturbation of conditions in nonstrongly regular Samarskii-Ionkin type problems / N.S. Imanbaev // International Journal of Differential Equations. — 2015. — Article ID 641481. — P. 1–6. doi: <http://dx.doi.org/10.1155/2015/641481>
- 28 Sadybekov M.A. Characteristic Determinant of a Boundary Value Problem, which does not have the basis Property / M.A. Sadybekov, N.S. Imanbaev // Eurasian Mathematical Journal. — 2017. — 8. — No. 2. — P. 40–46.
- 29 Sadybekov M.A. On the integral perturbation of the boundary condition of one problem that does not have a basic property / M.A. Sadybekov, N.S. Imanbaev // Kazakh Mathematical Journal. — 2019. — No. 19(3). — P. 55–65.
- 30 Imanbaev N.S. On a Problem that Does not Have Basis Property of Root Vectors, Associated with a Perturbed Regular Operator of Multiple Differentiation / N.S. Imanbaev // Journal of Siberian Federal University. Mathematics & Physics. — 2020. — 13. — No. 5. — P. 568–573.

Н.С. Иманбаев<sup>1,2</sup>, Е. Құрмыш<sup>1</sup>

<sup>1</sup>Оңтүстік Қазақстан мемлекеттік педагогикалық университеті, Шымкент, Қазақстан;

<sup>2</sup>Математика және математикалық модельдеу институты, Алматы, Қазақстан

## Кесіндідегі үшінші ретті регулярлы дифференциалдық оператормен байланысқан, экспоненциалды типтегі квазикөпмүшеліктермен сәйкес келетін бүтін функцияның нөлдері жайлы

Мақалада көрсеткіштері өлшемді экспоненциалды типтегі квазикөпмүшеліктермен сәйкес келетін бір кластағы бүтін функциялардың нөлдерін зерттеу мәселесі қарастырылды. Мұндағы қарастырылатын мәселе, көп жағдайларда, кейбір кластардағы кесіндідегі дифференциалдық операторлардың меншікті мәндерін зерттеуге берілген есептерден туындайды. Дәлірек айтқанда, қарастырылатын мәселеге  $W_2^3(0, 1)$  кеңістігіндегі регулярлы шеттік шарттармен берілген үшінші ретті сызықтық дифференциалдық теңдеудің меншікті мәндерін зерттеуге арналған есепке алып келеді. Зерттелетін бүтін функция, тікелей периодтық шеттік шарттармен берілген сызықтық дифференциалдық үшінші ретті оператор үшін аталған спектралдық есептің характеристикалық анықтаушы болып табылады. А.Ф. Леонтьевтің монографиясындағы нәтижесінің негізінде, қарастырылып отырған бір кластағы өлшемді көрсеткіштері бар экспоненциалды типтегі квазикөпмүшеліктермен сәйкес келетін бүтін функцияның түйіндес индикаторлық диаграммасын құрудың алгоритмі көрсетілген. Бүтін функцияның әрбір сериядағы саналымды нөлдерінің бар болуы дәлелденген және олардың кесіндідегі регулярлы периодтық шеттік шарттармен берілген сызықтық үшінші ретті дифференциалдық оператордың меншікті

мәндері екендігі сипатталған. Бүтін функцияның әр сериядағы көршілес жатқан нөлдерінің арақашықтығы анықталған және әр серия комплексті жазықтықтағы түйіндес индикаторлық диаграмманың, яғни дұрыс алтыбұрыштың қабырғаларына перпендикуляр, координаталар бас нүктесінен шығатын сәулелер болатындығы көрсетілген. Алайда, нөл нүктесі жоғарыда айтылған қарастырылатын оператордың меншікті мәні болмайтындығы, яғни нөл оператордың регулярлы нүктесі екендігі сипатталған. Бұл жұмыстағы алынған нәтиженің ерекшелігі, оператордың әр сериядағы меншікті мәндеріне сәйкес меншікті функциялар жүйесінің табылуында. Сондай-ақ, осы жұмыстың зерттеу нысанына айналып отырған оператордың түйіндес операторы құрылған.

*Кілт сөздер:* бүтін функцияның нөлдері, квазикөпмүшеліктер, индикаторлық диаграмма, серия, оператор, регулярлы периодтық шеттік шарттар, меншікті мәндер, меншікті функциялардың жүйесі.

Н.С. Иманбаев<sup>1,2</sup>, Е. Курмыш<sup>1</sup>

<sup>1</sup> Южно-Казахстанский государственный педагогический университет, Шымкент, Казахстан;

<sup>2</sup> Институт математики и математического моделирования, Алматы, Казахстан

## О нулях целой функции, совпадающей с квазиполиномами экспоненциального типа, связанной с регулярным дифференциальным оператором третьего порядка на отрезке

В статье рассмотрен вопрос распределения нулей целой функции одного класса, которые являются квазиполиномами экспоненциального типа. К подобной проблеме редуцированы задачи на собственные значения для некоторых классов дифференциальных операторов на отрезке. В частности, к изучаемому вопросу приводит задача на собственные значения линейного дифференциального уравнения третьего порядка с регулярными краевыми условиями в пространстве  $W_2^3(0, 1)$ . Исследуемая целая функция адекватно является характеристическим определителем спектральной задачи для линейного дифференциального оператора третьего порядка с периодическими краевыми условиями. Построена сопряженная индикаторная диаграмма целой функции экспоненциального типа соизмеримыми показателями. Доказано существование счетного числа нулей исследуемой целой функции в каждой серии, которые являются одновременно собственными значениями рассматриваемого дифференциального оператора третьего порядка с периодическими краевыми условиями. Определено расстояние между соседними нулями каждой серии, лежащее на лучах, перпендикулярных сторонам сопряженной индикаторной диаграммы, то есть правильного шестиугольника на комплексной плоскости. При этом ноль не является собственным значением рассматриваемого оператора. Принципиальным отличием настоящей работы является нахождение соответствующих собственных функций рассматриваемого оператора. Построен сопряженный оператор.

*Ключевые слова:* целая функция, нули, квазиполиномы, индикаторная диаграмма, серия, оператор, регулярные периодические краевые условия, собственные значения, система собственных функций.

### References

- 1 Birkhoff, G.D. (1908). On asymptotic character of solutions of certain linear differential equations containing a parameter. *Transactions of the American Mathematical Society*, 9, 2, 219–231.
- 2 Birkhoff, G.D. (1908). Boundary value and expansion problems of ordinary linear differential equations. *Transactions of the American Mathematical Society*, 9, 4, 373–395.
- 3 Naimark, M.A. (1969). *Lineinye differentsialnye operatory [Linear Differential Operators]*. Moscow: Nauka [in Russian].
- 4 Titchmarsh, E.C. (1926). Zeros of certain integral functions. *Proc. London Math. Soc.* 25, 4, 283–302.
- 5 Bellman, R., & Cook, K. (1967). *Differentsialno-raznostnye uravneniia [Differential-Difference Equations]*. Moscow [in Russian].
- 6 Leontiev, A.F. (1983). *Tselye funktsii i riady eksponent [Entire functions and Exponential Series]*. Moscow [in Russian].

- 7 Hald, O.H. (1984). Discontinuous Inverse Eigenvalue Problems. *Communications on Pure Applied Mathematics*, XXXVII, 539–577.
- 8 Lidskiy, V.B., & Sadovnichy, V.A. (1967). Regularizovannye summy kornei odnogo klassa tselykh funktsii [Regularized sums of roots of one class of entire functions]. *Functional analysis*, 1, 2, 52–59 [in Russian].
- 9 Sedletsky, A.M. (1995). Kogda vse nuli tseloi funktsii eksponentsialnogo tipa lezhat v krivolineinoi poluploskosti (neobkhodimoe uslovie) [When all zeros of an entire function of exponential type lie in a curvilinear half-plane (necessary condition)]. *Mathematical Collection*, 186, 9, 125–134 [in Russian].
- 10 Shkalikov, A.A. (1982). O bazisnosti sobstvennykh funktsii obyknovennykh differentsialnykh operatorov s integralnymi kraevymi usloviiami [On the basis property of eigenfunctions of ordinary differential operators with integral boundary value conditions]. *Vestnik Moskovskogo gosudarstvennogo universiteta. Seriya Math. mech.* 6, 12–21 [in Russian].
- 11 Imanbaev, N.S. (2018) Distribution of Eigen values of a Third-Order Differential Operator with Strongly Regular nonlocal Boundary Conditions. *AIP Conference Proceedings*, 1997: 020027, 020027-1 – 020027-5.
- 12 Sadovnichy, V.A., Lyubishkin, V.A., & Belabbasi, Yu. (1980). O regularizovannykh summakh kornei tseloi funktsii odnogo klassa [On regularized sums of roots of an entire function of one class]. *Doklady AN SSSR*. 254, 6, 1346–1348 [in Russian].
- 13 Sadovnichy, V.A., Lyubishkin, V.A., & Belabbasi, Yu. (1982). O nuliakh tselykh funktsii odnogo klassa [On zeros of entire functions of one class]. *Trudy seminara imeni I.G. Petrovskogo*, 8, 211–217 [in Russian].
- 14 Imanbaev, N.S. (2018). On zeros of a quasi-polynomial of exponential type connected with a regular third order differential operator. *Mathematical Journal*, 18, 2(68), 124–132.
- 15 Levin, B.Ya. (1956). *Raspredelenie kornei tselykh funktsii [Distribution of Roots of Entire Functions]*. Moscow [in Russian].
- 16 Sedletskiy, A.M.(1993). O nuliakh preobrazovaniia Fure finitnoi mery [On zeros of the Fourier transform of a finite measure]. *Matematicheskie seminary – Mathematical Notes*. 53, 1, 111–119 [in Russian].
- 17 Kanguzhin, B.E., & Sadybekov, M.A. (1996). *Differentsialnye operatory na otrezke. Raspredelenie sobstvennykh znachenii [Differential Operators on a Segment. Distribution of eigenvalues]*. Shymkent: Gylym [in Russian].
- 18 Imanbaev, N.S., & Kanguzhin, B.E. (1995). O nuliakh tselykh funktsii, imeiushchikh integralnoe predstavlenie [On zeros of entire functions having an integral representation]. *News of the National Academy of Sciences of the Republic of Kazakhstan. Series Phys.-Math*, 3, 47–52 [in Russian].
- 19 Imanbaev, N.S., Kanguzhin, B.E., & Kalimbetov, B.E.(2013). On zeros of the characteristic determinant of the spectral problem for a third-order differential operator on a segment with nonlocal boundary conditions *Advances in Difference Equations*. doi: 10.1186/1687-1847-20113-110.
- 20 Lunyov, A.A., & Malamud, M.M. (2015). On completeness and Riesz basis property of root subspaces of boundary value problems for first order systems and applications. *Journal of Spectral Theory*, 5, 17–70. doi 10.4171/jst/90
- 21 Sadybekov, M.A., & Imanbaev, N.S. (2017). A Regular Differential Operator with Perturbed Boundary Condition *Mathematical Notes*. 101, 5, 878–887.
- 22 Imanbaev, N.S., & Ospanov, M.N. (2019). On eigen values of third order composite type equations with regular boundary value conditions. *Bulletin of the Karaganda University. Mathematics series*, 4(96), 44–51. doi 10.31489/2019M4/44-51
- 23 Imanbaev, N.S., & Ospanov, M.N. (2019). Raspredelenie sobstvennykh znachenii uravneni tretiego poriadka sostavnogo tipa s reguliarnymi kraevymi usloviiami [Distribution of eigenvalues of the third order equations of composite type with regular boundary value conditions]. *Bulletin of the Institute of Mathematics. Tashkent*, 5, 28–33. doi http://mib.mathinst.UZ [in Russian].
- 24 Makin, A.S.(2006). On a nonlocal perturbation of a periodic eigenvalue problem. *Differential Equations*, 42, 4, 599–602.

- 25 Kritskov, L.V., Sadybekov, M.A., & Sarsenbi, A.M. (2018). Nonlocal spectral problem for a second-order differential equation with an involution. *Bulletin of the Karaganda University. Mathematics series. Special issue.*, 3(91), 53–61.
- 26 Baranetsky, Ya.O., & Kalenyuk, P.I.(2020). Nonlocal Multipoint Problem with Multiple Spectrum for an Ordinary (2n) Order Differential Equation. *Journal of Mathematical Sciences*, 246, 152–169.
- 27 Imanbaev, N.S. (2015). On stability of the basis property of root vectors system of the Sturm-Liouville operator with an integral perturbation of conditions in nonstrongly regular Samarskii-Ionkin type problems. *International Journal of Differential Equations*, Article ID 641481, 6 pages. doi <http://dx.doi.org/10.1155/2015/641481>
- 28 Sadybekov, M.A., & Imanbaev, N.S. (2017). Characteristic Determinant of a Boundary Value Problem, which does not have the basis Property. *Eurasian Mathematical Journal*, 8, 2, 40–46.
- 29 Sadybekov, M.A., & Imanbaev, N.S. (2019). On the integral perturbation of the boundary condition of one problem that does not have a basic property. *Kazakh Mathematical Journal*, 19(3), 55–65.
- 30 Imanbaev, N.S. (2020). On a Problem that Does not Have Basis Property of Root Vectors, Associated with a Perturbed Regular Operator of Multiple Differentiation. *Journal of Siberian Federal University. Mathematics & Physics.*, 13, 5, 568–573.