

ORGANIC CHEMISTRY

Article

Received: 1 September 2022 | Revised: 12 January 2023 |
Accepted: 12 February 2023 | Published online: 06 March 2023

UDC 547.8

<https://doi.org/10.31489/2959-0663/1-23-8>

Venkatesh B. Gopula^{1*} , Mohsin K. Pathan² 

¹Department of Chemistry, Anandibai Raorane Arts, Commerce and Science College,
Vaibhavwadi, Dist. Sindhudurg, Maharashtra, India;

²Department of Analytical Method Development Lab (R&D), Jamjoom Pharmaceutical, Jeddah, Saudi Arabia
(*Corresponding author's e-mail: gopulavenkatesh@gmail.com)

Simple and Efficient Method for One Pot Multicomponent Synthesis of 3,4-Dihydropyrimidin-2-(1H)-One Derivatives Catalyzed by Organocatalyst: Benzoic Acid

The Biginelli reaction is one the useful multicomponent reactions and a very appropriate reaction for the synthesis of 3,4-dihydropyrimidin-2(1H)-ones derivatives. These 3,4-dihydropyrimidin-2(1H)-ones have biological and pharmacological properties which make them a very important class of medicinal chemistry. Although they are an important class of medicinal chemistry, the syntheses of these compounds have been catalyzed by large number of strong Bronsted acids and Lewis acids under thermal conditions. Small organic molecules, organocatalysts, have been used as catalysts for the Biginelli reaction in a small number as compared to Bronsted acids and Lewis acid. Benzoic acid, which is a small organic molecule, although an acid, has never been tested for the synthesis of 3,4-dihydropyrimidin-2(1H)-ones. Benzoic acid is an inexpensive, non-toxic molecule, it has been successfully tested here as a catalyst for the one-pot three component synthesis of 3,4-dihydropyrimidin-2-(1H)-one derivatives via Biginelli reaction between β -keto ester, a variety of aromatic aldehydes and urea or thiourea under thermal conditions using 20 mol% benzoic acid in acetonitrile solvent refluxed for 12 h to give good to high yields. This synthetic method includes inexpensive, non-toxic, easily available benzoic acid as a catalyst and is carried out in a simple operational procedure.

Keywords: organocatalyst, benzoic acid, Biginelli, 3,4-dihydropyrimidin-2-(1H)-one, β -keto ester, aldehydes, urea, thiourea.


Introduction

The Biginelli reaction is one of the multicomponent reactions. It is a very suitable reaction for preparation of the 3,4-dihydropyrimidin-2(1H)-ones. 3,4-Dihydropyrimidinones (DHPMs) are a very important class since the recent discovery of their biological and pharmacological properties, namely antiviral, antitumour, antibacterial, anti-inflammatory and antihypertensive ones. Dihydropyrimidinone cores are also found in many natural products, highlighting important efforts aimed at the synthesis of these heterocycles [1–6]. These DHPMs have become a crucial basis for several calcium channel blockers, antihypertensives, α_{1a} -adrenergic antagonists and neuropeptide Y (NPY) antagonists [7].

Conventionally, Biginelli reactions were performed under strong acidic conditions with heating, and the yields were low to moderate (Scheme 1) [8].

Traditionally, there are several suitable reaction conditions with strong Bronsted acid along with the use of Lewis acid in condensation to more efficiently synthesize these DHPMs molecules; e.g. H_2SO_4 [7], $BF_3 \cdot Et_2O/CuCl$ [9], $BiCl_3$ [10], $CeCl_3 \cdot 7H_2O$ [11], $Cu(OTf)_2$ [12], $TiCl_4$ [13], $LiBr$ [14], Gallium (III) halides [15], Metal triflimide [16], *p*-toluenesulfonic acid [17], polystyrenesulfonic acid (PSSA) [18], $Cu(OTf)_2$ [19],

ZrCl₄ [20], FeCl₃·6H₂O [21], RuCl₃ [22], Bi(NO₃)₃·5H₂O–TBAF [23], SmI₂ [24]. In addition to these, the Biginelli reaction has also been reported by using nanomaterials [25–27], nano-composite [28], zeolites [29–31], polymers [32], ultrasonic irradiation [33, 34], microwaves [35–38] and ball milling [39].



Scheme 1

The use of small organic molecules known as Organocatalysts [40–44] is a very rapidly developing area of synthetic organic chemistry that is replacing the use of metal-based Lewis acids. Organocatalysis offers many advantages for synthetic organic chemistry. In contrast to many transition metal catalysts, most organocatalysts are resistant to air and water, easily handled experimentally, relatively nontoxic, and readily separated from the crude reaction mixture [45]. Given the growing interest in developing green processes and procedures in organic synthesis [46], organocatalysts are considered to be a more eco-friendly and user-friendly alternative to traditional counterparts. Because of these many advantages of organocatalysts, few organocatalysts have been explored in the synthesis of DHPM derivatives such as bakers' yeast [47], hydrazine type [48], oxalic acid [49, 50], citric acid [51], boric acid [52], phenylboronic acid [53], L-proline [54] and Lactic acid [55].

In continuation of our work on new methodologies using organocatalysts [44, 56] in the synthetic organic chemistry we wish to report the efficient use of benzoic acid and a very simple effective approach for the synthesis of 3,4-dihydropyrimidin-2-(1H)-ones derivatives via the Biginelli reaction under thermal conditions having good to high yield.

In addition to those listed above, numerous methods for the synthesis of 3,4-dihydropyrimidin-2-(1H)-one derivatives by the Biginelli reaction are available in the literature. However, few organocatalysts are available for the synthesis of DHPMs. Benzoic acid is cheap, non-toxic and readily available in every laboratory. To our knowledge, so far the potential of benzoic acid as a mild organocatalyst has not been much tested in organic synthesis. We decided to explore the potential of benzoic acid as an organocatalyst for the synthesis of 3,4-dihydropyrimidin-2-(1H)-ones and their thione analogs as well as their derivatives by the Biginelli reaction under thermal conditions.

Experimental

The following reagents were used in this work without additional purification; benzaldehyde, 4-chlorobenzaldehyde, 3-nitrobenzaldehyde, 4-methoxybenzaldehyde, 4-nitrobenzaldehyde, 4-hydroxybenzaldehyde, ethylacetoacetate, methylacetoacetate, benzoic acid, urea, thiourea, acetonitrile, ethanol (loba Chemie), 4-methylbenzaldehyde, 3-methoxybenzaldehyde (Sigma-Aldrich).

The general procedure for the Biginelli reaction is as follows: a solution of the appropriate aldehyde **1** (1.0 mmol), urea or thiourea **2** (1.5 mmol), β -keto ester **3** (1.0 mmol), benzoic acid (20 mol%, 0.2 mmol) in acetonitrile (10 mL) is heated to reflux for 12 h. Then it is cooled to room temperature and poured into ice-water about 50mL. The solid products are filtered, washed with ice water, dried and recrystallized from ethanol to give pure product **4 (a-m)**.

4a. 5-Ethoxycarbonyl-6-methyl-4-phenyl-3,4-dihydropyrimidin-2(1H)-one, Melting point: Found 206–208 °C (200–202 °C) [57]. ¹H NMR (DMSO-d₆): δ = 9.18 (s, 1H), 7.73 (s, 1H), 7.20–7.30 (m, 5H), 5.14 (s, 1H), 3.98 (q, J = 7.2 Hz, 2H), 2.24 (s, 3H), 1.06 (t, J = 7.2 Hz, 3H). ¹³C NMR (DMSO-d₆): δ = 166.3, 152.1, 147.2, 144.2, 128.1, 127.2, 126.5, 99.0, 58.1, 53.9, 17.9, 14.0. Mass m/z [M+1]⁺ = 261.11.


4b. 4-(4-Chlorophenyl)-5-ethoxycarbonyl-6-methyl-3,4-dihydropyrimidin-2(1H)-one, Melting point: Found 214–216 °C (210–212 °C) [58]. ¹H NMR (DMSO-d₆): δ = 9.20 (s, 1H), 7.76 (s, 1H), 7.40 (d, J = 9.0 Hz, 2H), 7.26 (d, J = 9.0 Hz, 2H), 5.16 (s, 1H), 3.95 (q, J = 7.1 Hz, 2H), 2.19 (s, 3H), 1.10 (t, J = 7.1 Hz, 3H). ¹³C NMR (DMSO-d₆): δ = 167.1, 151.4, 146.4, 141.9, 130.1, 126.2, 125.5, 98.0, 56.1, 53.6, 19.7, 13.4. Mass m/z [M+1]⁺ = 295.07.

Results and Discussion

We started our study with the reaction of benzaldehyde, ethyl acetoacetate and urea using benzoic acid as a catalyst under thermal conditions (Scheme 2) as a model reaction and the results are summarized in Table 1.

Initially, the reaction was carried out without a catalyst for an appropriate time and no progress was observed in acetonitrile solvent (Table 1, entry 01). Catalyst 1 mole % was added and the reaction was carried out up to 18 h, the obtained yield was very low (Table 1, entry 05).

The reaction gradually progressed with increasing the amount of catalyst while maintaining the reaction time (Table 1, entries 06 & 07). Due to the gradual development of the reaction, the amount of catalyst was increased to 7.5 mol % to see the reaction development, reducing the reaction time to 15 hours, a yield of 60 % was obtained (Table 1, entry 08). To obtain high and excellent yields, we continued to vary the amount of catalyst (Table 1, entries 09 & 16). The best reaction conditions were obtained using 20 mol% catalyst and the reaction was completed after 12 hours with a yield of 92 % (Table 1, entry 17). To our curiosity, we continued the reaction again up to 15 hours and also increased the amount of catalyst, but there was no any significant improvement in either case (Table 1, entry 18 & 19). In addition to acetonitrile, we also optimized the reaction conditions using different solvents as shown in Table 1. Without using a catalyst none of the reactions progressed to give any product (Table 1, entries 02, 03 & 04). Instead of introducing lower amount of catalyst for solvents water, THF and ethanol, we purposefully used 10 mol% and 20 mol% catalyst for our study. Yields were obtained relatively lower as compared with acetonitrile using 10 mol% catalyst for 15 hours (Table 1, entries 10, 11 & 12). Also, approximately the same results were observed using 20 mol% catalyst for 12 hours (Table 1, entries 13, 14 & 15) as compared with acetonitrile solvent (Table 1, entry 17).



Scheme 2

Table 1


Optimization of the reaction conditions in the synthesis of DHPM^a

Entry	Solvent	Catalyst (mole %)	Time (h)	Yield ^b (%)
01	CH ₃ CN	None	18	0
02	H ₂ O	None	18	0
03	THF	None	18	0
04	EtOH	None	18	0
05	CH ₃ CN	1	18	20
06	CH ₃ CN	2	18	30
07	CH ₃ CN	5	18	46
08	CH ₃ CN	7.5	15	60
09	CH ₃ CN	10	15	72
10	H ₂ O	10	15	20
11	THF	10	15	35
12	EtOH	10	15	30
13	H ₂ O	20	15	32
14	THF	20	12	65
15	EtOH	20	12	54
16	CH ₃ CN	15	15	85
17	CH ₃ CN	20	12	92
18	CH ₃ CN	20	15	94
19	CH ₃ CN	25	15	96

^aReaction conditions: Benzaldehyde (1.0 mmol), Urea (1.5 mmol), Ethyl acetoacetate (1.0 mmol), catalyst, solvent (10 ml) refluxed for the appropriate time.

^bIsolated yields.

After obtaining the best reaction conditions; 1.0 mmol of benzaldehyde, 1.0 mmol of ethyl acetoacetate, 1.5 mmol of urea, 20 mol% benzoic acid as a catalyst in 10 ml of acetonitrile refluxed for 12 hrs, in order to explore the scope and generality of the present reaction conditions, various aldehydes and an urea analog, i.e. thiourea were taken for the Biginelli reaction using our optimized reaction conditions (Scheme 3). The results obtained are summarized in Table 2.



In addition to the parent benzaldehyde, aldehydes with electron-donating groups such as methyl-, methoxy- and chloro-type groups, also electron withdrawing group such as nitro- were treated with ethyl acetoacetate and urea, resulting in the formation of the corresponding DHPM derivative in 84–92 % yield (Table 2, entries 01–05). The urea analog, i.e., thiourea, was also tested in the same reaction with benzaldehydes bearing chloro-, hydroxyl-, nitro- and methoxy-groups yielding the corresponding product in 82–90 % yield (Table 2 entries 06–10). Reaction with methylacetoacetate showed results similar to the reaction with ethylacetoacetate and gave the corresponding DHPM (Table 2, entries 11–12) in high yield. This reaction was also suitable for aldehydes with α -hydrogen, such as *n*-valeraldehyde (Table 2, entry 13), to give the corresponding DHPM product in good yield, and a significant reduction in yield was observed compared with aromatic aldehydes.

Table 2

Scope of Aldehydes with urea/thiourea and β -keto esters in the synthesis of DHPM derivatives^a

Entry	Aldehyde 1 (R)	Urea/thiourea 2 (X)	β -keto ester 3 (R ₁)	Product 4	Yield ^b %
01	C ₆ H ₅	O	Et	4a	92
02	4-ClC ₆ H ₄	O	Et	4b	90
03	4-CH ₃ C ₆ H ₄	O	Et	4c	84
04	4-OCH ₃ C ₆ H ₄	O	Et	4d	87
05	4-NO ₂ C ₆ H ₄	O	Et	4e	90
06	C ₆ H ₅	S	Et	4f	84
07	4-OHC ₆ H ₄	S	Et	4g	82
08	3-NO ₂ C ₆ H ₄	S	Et	4h	87
09	3-OCH ₃ C ₆ H ₄	S	Et	4i	90
10	4-ClC ₆ H ₄	S	Et	4j	86
11	C ₆ H ₅	O	Me	4k	86
12	4-ClC ₆ H ₄	O	Me	4l	82
13	<i>n</i> -Bu	O	Et	4m	72

^aReaction conditions: Aldehyde (1.0 mmol), Urea/thiourea (1.5 mmol), β -keto esters (1.0 mmol), benzoic acid 20 mol%, in acetonitrile (10 ml) refluxed for the 12 h.
^bIsolated yields.

Conclusions

In conclusion, we have developed a simple and efficient method for one-pot three-component synthesis of 3,4-dihydropyrimidinone derivatives from aldehydes with β -keto esters and urea or thiourea using benzoic acid as an organocatalyst by refluxing in acetonitrile in high yields of 72–92 %. This method using benzoic acid as an organocatalyst is very useful to explore diversified aldehydes for synthesizing derivatives of 3,4-dihydropyrimidinones as well as thione analogues, which can play crucial role in the field of medicinal chemistry.

Acknowledgements

We would like to thank Anandibai Raorane Arts, Commerce and Science College, Vaibhavwadi for providing all necessary facilities for conducting this research work.

References

- 1 Atwal, K. S., Rovnyak, G. C., O'Reilly, B. C., & Schwartz, J. J. (1989). Substituted 1,4-dihydropyrimidines. 3. Synthesis of selectively functionalized 2-hetero-1,4-dihydropyrimidines. *Organic Chemistry*, 54, 5898–5907. <https://doi.org/10.1021/jo00286a020>
- 2 Atwal, K. S., Swanson, B. N., Unger, S. E., Floyd, D. M., Moreland, S., Hedberg, A., & O'Reilly, B. C. (1991). Dihydropyrimidine calcium channel blockers. 3. 3-Carbamoyl-4-aryl-1,2,3,4-tetrahydro-6-methyl-5-pyrimidinecarboxylic acid esters as orally effective antihypertensive agents. *Journal of Medicinal Chemistry*, 34, 806–811. <https://doi.org/10.1021/jm00106a048>
- 3 Rovnyak, G. C., Atwal, K. S., Hedberg, A., Kimball, S. D., Moreland, S., Gougoutas, J. Z., O'Reilly, B. C., Schwartz, J., Malley, M. F., & Semones, M. A. (1992). Dihydropyrimidine calcium channel blockers. 4. Basic 3-substituted-4-aryl-1,4-dihydropyrimidine-5-carboxylic acid esters. Potent antihypertensive agents. *Journal of Medicinal Chemistry*, 35, 3254–3263. <https://doi.org/10.1021/jm00095a023>
- 4 Kappe, C. O. & Fabian, W. M. F. (1997). Conformational analysis of 4-aryl-dihydropyrimidine calcium channel modulators. A comparison of ab initio, semiempirical and X-ray crystallographic studies. *Tetrahedron*, 53, 2803–2816. [https://doi.org/10.1016/S0040-4020\(97\)00022-7](https://doi.org/10.1016/S0040-4020(97)00022-7)
- 5 Kappe, C. O., Shishkin, O. V., Uray, G., & Verdino, P. (2000). X-Ray Structure, Conformational Analysis, Enantioseparation, and Determination of Absolute Configuration of the Mitotic Kinesin Eg5 Inhibitor Monastrol. *Tetrahedron*, 56, 1859–1862. [https://doi.org/10.1016/S0040-4020\(00\)00116-2](https://doi.org/10.1016/S0040-4020(00)00116-2)
- 6 Rovnyak, G. C., Kimball, S. D., Beyer, B., Cucinotta, G., DiMarco, J. D., Gougoutas, J., Hedberg, A., Malley, M., McCarthy, J. P., Zhong, R., & Moreland, S. (1995). Calcium Entry Blockers and Activators: Conformational and Structural Determinants of Dihydropyrimidine. *Calcium Channel Modulators Journal of Medicinal Chemistry*, 38, 119–129. <https://doi.org/10.1021/jm00001a017>
- 7 Azizian, J., Mohammadi, M. K., Firuzi, O., Mirza, B., & Miri, R. (2010). Microwave-Assisted Solvent-Free Synthesis of Bis(dihydropyrimidinone)benzenes and Evaluation of their Cytotoxic Activity. *Chemical Biology & Drug Design*, 75, 375–380. <https://doi.org/10.1111/j.1747-0285.2009.00937.x>
- 8 Kappe, C. O. (2000). Recent Advances in the Biginelli Dihydropyrimidine Synthesis. *New Tricks from an Old Dog. Accounts of Chemical Research*, 33, 879–888. <https://doi.org/10.1021/ar000048h>
- 9 Hu, E. H., Sidler, D. R., & Dolling, U. H. (1998). Unprecedented catalytic three component one-pot condensation Reaction: An Efficient Synthesis of 5-Alkoxy-carbonyl-4-aryl-3,4-dihydropyrimidin-2(1H)-ones. *The Journal of Organic Chemistry*, 63, 3454–3457. <https://doi.org/10.1021/jo970846u>
- 10 Ramalinga, K., Vijayalakshmi, P., & Kaimala, T. N. B. (2001). Bismuth (III)-catalyzed Synthesis of Dihydropyrimidones: Improved protocol conditions for the Biginelli Reaction. *Synlett*, 6, 863–865. <https://doi.org/10.1055/s-2001-14587>
- 11 Bose, D. S., Fatima, L., & Mereyala, H. B. (2003). Green Chemistry Approaches to the Synthesis of 5-Alkoxy-carbonyl-4-aryl-3,4-dihydropyrimidin-2(1H)-ones by a Three-Component Coupling of One-Pot Condensation Reaction: Comparison of Ethanol, Water, and Solvent-free Conditions. *The Journal of Organic Chemistry*, 68, 587–590. <https://doi.org/10.1021/jo0205199>
- 12 Paraskar, A. S., Dewkar, G. K., & Sudalai, A. (2003). Cu(OTf)₂: a reusable catalyst for high-yield synthesis of 3,4-dihydropyrimidin-2(1H)-ones. *Tetrahedron Letters*, 44, 3305–3308. [https://doi.org/10.1016/S0040-4039\(03\)00619-1](https://doi.org/10.1016/S0040-4039(03)00619-1)
- 13 Maiti, G., Kundua, P., & Guin, C. (2003). One-pot synthesis of dihydropyrimidinones catalysed by lithium bromide: an improved procedure for the Biginelli reaction. *Tetrahedron Letters*, 44, 2757–2758. [https://doi.org/10.1016/S0040-4039\(02\)02859-9](https://doi.org/10.1016/S0040-4039(02)02859-9)
- 14 Baruah, P. P., Gadhwal, S., Prajapati, D., & Sandhu, J. S. (2002). The Biginelli Condensation: A Novel and Efficient Regioselective Synthesis of Dihydropyrimidin-2(1H)-ones Using Lithium Bromide. *Chemistry Letters*, 31, 1038–1039. <https://doi.org/10.1246/cl.2002.1038>
- 15 Saini, A., Kumar, S., & Sandhu, J. S. (2007). Gallium (III) halides catalyzed, microwave enhanced, synthesis of 3,4-dihydropyrimidin-2(1H)-ones under solvent free condition. *Indian Journal of Chemistry*, 46B, 1886–1889.
- 16 Suzuki, I., Suzumura, Y., & Takeda, K. (2006). Metal triflimide as a Lewis acid catalyst for Biginelli reaction in water. *Tetrahedron Letters*, 47, 7861–7864. <https://doi.org/10.1016/j.tetlet.2006.09.019>
- 17 Bose, A. K., Manhas, M. S., Pednekar, S., Ganguly, S. N., Dang, H., He, W., & Mandadi, A. (2005). Large scale Biginelli reaction via water-based biphasic media: a green chemistry strategy. *Tetrahedron Letters*, 46, 1901–1903. <https://doi.org/10.1016/j.tetlet.2005.01.087>
- 18 Polshettiwar, V., & Varma, R. S. (2007). Biginelli reaction in aqueous medium: a greener and sustainable approach to substituted 3,4-dihydropyrimidin-2(1H)-ones. *Tetrahedron Letters*, 48, 7343–7346. <https://doi.org/10.1016/j.tetlet.2007.08.031>
- 19 Paraskar, A. S., Dewkar, G. K., & Sudalai, A. (2003). Cu(OTf)₂: a reusable catalyst for high-yield synthesis of 3,4-dihydropyrimidin-2(1H)-ones. *Tetrahedron Letters*, 44, 3305–3308. [https://doi.org/10.1016/S0040-4039\(03\)00619-1](https://doi.org/10.1016/S0040-4039(03)00619-1)
- 20 Reddy, C. V., Mahesh, M., Raju, P. V. K., Babu, T. R., & Reddy, V. V. N. (2002). Zirconium (IV) chloride catalyzed one-pot synthesis of 3,4-dihydropyrimidin-2(1H)-ones. *Tetrahedron Letters*, 43, 2657–2659. [https://doi.org/10.1016/S0040-4039\(02\)00280-0](https://doi.org/10.1016/S0040-4039(02)00280-0)

- 21 Lu, J. & Ma, H. R. (2000). Iron (III)-Catalyzed Synthesis of Dihydropyrimidinones, Improved conditions for the Biginelli Reaction. *Synlett*, 1, 63–64. <https://doi.org/10.1055/s-2000-6469>
- 22 Surya, K. De, & Gibbs, R. A. (2005). Ruthenium(III) chloride-catalyzed one pot synthesis of 3,4-dihydropyrimidin-2-(1H)-ones under solvent free conditions. *Synthesis*, 11, 1748–1750. <https://doi.org/10.1055/s-2005-869899>
- 23 Khodaei, M. M., Khosropour, A. R., & Jowkar, M. (2005). Bi(NO₃)₃·5H₂O – TBAF as an Efficient Reagent for in situ Oxidation: Dihydropyrimidinone formation from Benzyl Halides. *Synthesis*, 8, 1301–1304. <https://doi.org/10.1055/s-2005-861876>
- 24 Han, X., Xu, F., Luo, Y., & Shen, Q. (2005). An Efficient One-pot synthesis of Dihydropyrimidinones by a Samarium Diiodide Catalyzed Biginelli reaction under Solvent free Conditions. *European Journal of Organic Chemistry*, 8, 1500–1503. <https://doi.org/10.1002/ejoc.200400753>
- 25 Zamani, F. & Izadi, E. (2013). Synthesis and characterization of sulfonated-phenylacetic acid coated Fe₃O₄ nanoparticles as a novel acid magnetic catalyst for the Biginelli reaction. *Catalysis Communications*, 42, 104–108. <https://doi.org/10.1016/j.catcom.2013.08.006>
- 26 Nasr-Esfahani, M., Hoseini, S. J. & Mohammadi, F. (2011). Fe₃O₄ Nanoparticles as an Efficient and Magnetically Recoverable Catalyst for the Synthesis of 3,4-dihydropyrimidin-2(1H)-ones under solvent free conditions. *Chinese Journal of Catalysis*, 32, 1484–1489. [https://doi.org/10.1016/S1872-2067\(10\)60263-X](https://doi.org/10.1016/S1872-2067(10)60263-X)
- 27 Girija, D., Naik, H. S. B., Kumar, B. V., Sudhamani, C. N., & Harish, K. N. (2019). Fe₃O₄ Nanoparticle supported Ni(II) Complexes: A Magnetically recoverable Catalyst for Biginelli reaction. *Arabian Journal of Chemistry*, 12, 420–428. <https://doi.org/10.1016/j.arabjc.2014.08.008>
- 28 Safari, J. & Zarnegar, Z. (2013). Biginelli reaction on Fe₃O₄-MWNT nanocomposite: excellent reactivity and facile recyclability of the catalyst combined with ultrasound irradiation. *RSC Advances*, 3, 17962–17967. <https://doi.org/10.1039/C3RA43014F>
- 29 Mistry, S. R., Joshi, R. S., Sahoo, S. K., & Maheria, K. C. (2011). Synthesis of Dihydropyrimidinones Using Large Pore Zeolites. *Catalysis Letters*, 141, 1541–1547. <https://doi.org/10.1007/s10562-011-0639-6>
- 30 Rani, V. R., Srinivas, N., Kishan, M. R., Kulkarni, S. J., & Raghavan, K. V. (2001). Zeolite-catalyzed cyclocondensation reaction for the selective synthesis of 3,4-dihydropyrimidin-2(1H)-ones. *Green Chemistry*, 3, 305–306. <https://doi.org/10.1039/B107612B>
- 31 Kang, L., Jin, D., & Cai, Y. (2013). Silica-supported ionic liquid Si[SbSipim][PF₆]: An efficient catalyst for the synthesis of 3,4-dihydropyrimidin-2-(1H)-ones. *Synthetic Communications*, 43, 1896–1901. <https://doi.org/10.1080/00397911.2012.678462>
- 32 Pourjavadi, A., Salimi, H., Barzegar, S., & Eftekharis, B. (2007). A Novel Polymeric catalyst for One-Pot synthesis of 3,4-dihydropyrimidin-2-(1H)-ones via Biginelli Reaction. *Acta Chimica Slovenica*, 54, 140–143.
- 33 Li, J. T., Han, J. F., Yang, J. H., & Li, T. S. (2003). An efficient synthesis of 3,4-dihydropyrimidin-2-ones catalyzed by NH₂SO₃H under ultrasound irradiation. *Ultrasonics Sonochemistry*, 10, 119–122. [https://doi.org/10.1016/S1350-4177\(03\)00092-0](https://doi.org/10.1016/S1350-4177(03)00092-0)
- 34 Yadav, J. S., Reddy, B. V. S., Reddy, K. B., Raj, K. S., & Prasad, A. R. (2001). Ultrasound- accelerated synthesis of 3,4-dihydropyrimidin-2(1H)-ones with ceric ammonium nitrate. *Journal of the Chemical Society, Perkin Transactions 1*, 1939–1941. <https://doi.org/10.1039/B102565C>
- 35 Cheng, J. & Qi, D. Y. (2007). ChemInform Abstract: An Efficient and Solvent-Free One-Pot Synthesis of Dihydropyrimidinones under Microwave Irradiation. *Chinese Chemical Letters*, 18, 647–650. <https://doi.org/10.1016/j.ccllet.2007.04.002>
- 36 Felluga, F., Benedetti, F., Berti, F., Drioli, S., & Regini, G. (2018). Efficient Biginelli Synthesis of 2-Aminodihydropyrimidines under Microwave Irradiation. *Synlett*, 29, 986–992. <http://doi.org/10.1055/s-0036-1591900>
- 37 Misra, A. K., Agnihotri, G., & Madhusudan, S. K. (2001). Microwave induced eco-friendly solvent-free Biginelli reaction catalyzed by calcium chloride. *Indian Journal of Chemistry*, 43B, 2018–2020.
- 38 Hazarkhani, H. & Karimi, B. (2004). N-Bromosuccinimide as an Almost Neutral Catalyst for Efficient Synthesis of Dihydropyrimidinones Under Microwave Irradiation. *Synthesis*, 8, 1239–1242. <https://doi.org/10.1055/s-2004-822348>
- 39 Mohamed, M. O., Alshammari, A. G. & Lemine, O. M. (2016). Green High-Yielding one-pot approach to Biginelli Reaction under catalyst-free and solvent free Ball Milling conditions. *Applied Sciences*, 6, 431–437. <https://doi.org/10.3390/app6120431>
- 40 Carlone, A., Cabrera, S., Marigo, M. & Jorgensen, K. A. (2007). A new approach for an organocatalytic multicomponent domino asymmetric reaction. *Angewandte Chemie International Edition*, 46, 1101–1104. <https://doi.org/10.1002/anie.200604479>
- 41 Xie, H. Zu, Li, H., Wang, J. & Jiang, W. J. (2007). Organocatalytic Enantioselective cascade Michael-Alkylation reactions: Synthesis of chiral Cyclopropanes and Investigation of Unexpected organocatalyzed Stereoselective Ring opening of cyclopropanes. *Journal of the American Chemical Society*, 129, 10886–10894. <https://doi.org/10.1021/ja073262a>
- 42 Martin, N. J. A., Ozores, L. & List, B. (2007). Organocatalytic Asymmetric transfer Hydrogenation of nitroolefins. *Journal of the American Chemical Society*, 129, 8976–8977. <https://doi.org/10.1021/ja074045c>
- 43 Li, G., Liang, Y. & Antilla, J.C. (2007). A vaulted Biaryl Phosphoric acid-catalyzed reduction of α -Imino Esters: The Highly Enantioselective preparation of α -amino esters. *Journal of the American Chemical Society*, 129, 5830–5831. <https://doi.org/10.1021/ja070519w>
- 44 Gopula, V. B. (2021) Mandelic acid: organocatalyst for efficient synthesis of secondary amines from aldehydes and primary amines. *Journal of Advanced Scientific Research*, 12(03), 292-296. <https://doi.org/10.55218/JASR.202112341>
- 45 Louis S. Hegeudus (2009). Organocatalysis in Organic Synthesis. *Journal of the American Chemical Society*, 131, 17995–17997. <https://doi.org/10.1021/ja908581u>
- 46 List, B. (2007). Introduction: Organocatalysis. *Chemical Reviews*, 107, 5413–5415. <https://doi.org/10.1021/cr078412e>

- 47 Kumar, A. & Maurya, R. M. (2007). An efficient baker's yeast catalyzed synthesis of 3,4-dihydropyrimidin-2-(1H)-ones. *Tetrahedron Letters*, 48, 4569–4571. <https://doi.org/10.1016/j.tetlet.2007.04.130>
- 48 Suzuki, I., Iwata, Y. & Takeda, K. (2008). Biginelli Reactions catalyzed by hydrazine type organocatalyst. *Tetrahedron Letters*, 49, 3238–3241. <https://doi.org/10.1016/j.tetlet.2008.03.080>
- 49 Sangshetti, J. N., Shinde, D. B. & Kokare, N. D. (2008). Oxalic acid as a versatile catalyst for one pot facile synthesis of 3,4-dihydropyrimidin-2-(1H)-ones and their thione analogues. *Journal of Heterocyclic Chemistry*, 45, 1191–1193. <https://doi.org/10.1002/jhet.5570450440>
- 50 Mohamadpour, F., Maghsoodlou, M. T., Heydari, R. & Lashkari, M. (2016). Oxalic acid dehydrate catalyzed synthesis of 3,4-dihydropyrimidin-2-(1H)-one derivatives under thermal and solvent free conditions. *Iranian Journal of Catalysis*, 6(3), 127–131.
- 51 Ramu, E., Kotra, V., Bansal, N., Varala, R. & Adapa, S. R. (2008). Green approach for the efficient synthesis of Biginelli compounds promoted by citric acid under solvent free conditions. *Rasayan Journal of Chemistry*, 1, 188–190.
- 52 Tu, S., Fang, F., Miao, C., Jiang, H., Feng, Y., Shi, D. & Wang, X. (2003). One-pot synthesis of 3,4-dihydropyrimidin-2-(1H)-ones using boric acid as catalyst. *Tetrahedron Letters*, 44, 6153–6155. [https://doi.org/10.1016/S0040-4039\(03\)01466-7](https://doi.org/10.1016/S0040-4039(03)01466-7)
- 53 Debache, A., Boumoud, B., Amimour, M., Belfaitah, A., Rhoutia, S. & Carboni, B. (2006). Phenylboronic acid as a mild and efficient catalyst for Biginelli Reaction. *Tetrahedron Letters*, 47, 5697–5699. <https://doi.org/10.1016/j.tetlet.2006.06.015>
- 54 Mabry, J. & Ganem, B. (2006). Studies on the Biginelli reaction: a mild and selective route to 3,4-dihydropyrimidin-2-(1H)-ones via enamine intermediates. *Tetrahedron Letters*, 47, 55–56. <https://doi.org/10.1016/j.tetlet.2005.10.124>
- 55 Saini, S. A., Kumar, D. & Sandhu, J. S. (2009). Multicomponent eco-friendly synthesis of 3,4-dihydropyrimidine-2-(1H)-ones using an organocatalyst Lactic acid. *Green Chemistry Letters and Reviews*, 1, 29–33. <https://doi.org/10.1080/17518250902973833>
- 56 Tale, R. H., Shinde, S. V., Gopula, V. B. & Wankhede, D. S. (2016). Methylboronic acid: A Mild, Green and recyclable Organocatalyst for Transformation of β -keto esters. *Journal of Chemical and Pharmaceutical Research*, 8(10), 169–175.
- 57 Ma, Y., Qian, C., Wang, L & Yang, M. (2000). Lanthanide triflate catalyzed Biginelli reaction. one-pot synthesis of dihydropyrimidinones under solvent-free conditions. *Journal of Organic Chemistry*, 65(12), 3864–3868. <https://doi.org/10.1021/jo9919052>
- 58 Shaabani, A., Bazgir, A. & Teimouri, F. (2003). Ammonium chloride-catalyzed one-pot synthesis of 3,4-dihydropyrimidin-2-(1H)-ones under solvent-free conditions. *Tetrahedron Letters*, 44(4), 857–859. [https://doi.org/10.1016/S0040-4039\(02\)02612-6](https://doi.org/10.1016/S0040-4039(02)02612-6)

Information about authors*

Gopula, Venkatesh Balkishan (*corresponding author*) — Assistant Professor, Department of Chemistry, Anandibai Raorane Arts, Commerce and Science College, Vaibhavwadi, Dist. Sindhudurg, Maharashtra, 416810, India; e-mail: gopulavenkatesh@gmail.com; <https://orcid.org/0000-0002-6193-5921>;

Pathan, Mohsin Khan Moin Khan — Sr. Executive, Dept. of Analytical Method Development Lab (R&D), Jamjoom Pharmaceutical, Jeddah, 21442, Saudi Arabia; e-mail: mohsin0506@gmail.com; <https://orcid.org/0000-0001-9307-3732>

*The author's name is presented in the order: Last Name, First and Middle Names