

A.H. Kalidolday^{1,*}, E.D. Nursultanov^{1,2}¹*L.N. Gumilyov Eurasian National University, Nur-Sultan, Kazakhstan;*²*M.V. Lomonosov Moscow State University (Kazakhstan branch), Nur-Sultan, Kazakhstan
(E-mail: aitolkynnur@gmail.com, er-nurs@yandex.kz)*

Interpolation of nonlinear integral Urysohn operators in net spaces

In this paper, we study the interpolation properties of the net spaces $N_{p,q}(M)$, in the case when M is a sufficiently general arbitrary system of measurable subsets from \mathbb{R}^n . The integral Urysohn operator is considered. This operator generalizes all linear, integral operators, and non-linear integral operators. The Urysohn operator is not a quasilinear or subadditive operator. Therefore, the classical interpolation theorems for these operators do not hold. A certain analogue of the Marcinkiewicz-type interpolation theorem for this class of operators is obtained. This theorem allows to obtain, in a sense, a strong estimate for Urysohn operators in net spaces from weak estimates for these operators in net spaces with local nets. For example, in order for the Urysohn integral operator in a net space, where the net is the set of all balls in \mathbb{R}^n , it is sufficient for it to be of weak type for net spaces, where the net is concentric balls.

Keywords: interpolation spaces, net spaces, Urysohn integral operators.

Introduction

Let (V, ν) , (U, μ) are measurable spaces and $Z(U)$, $M(V)$ are normed spaces of ν -measurable and μ -measurable functions, respectively. Let $K : \mathbb{R} \times U \times V \rightarrow \mathbb{R}$, and the operator $T : Z(U) \rightarrow M(V)$ is defined by the following equality: For any $f \in Z(U)$

$$T(f, y) = \int_U K(f(x), x, y) d\mu, \quad y \in V \quad (1)$$

and assume that this integral exists and is finite for almost all $y \in V$. This operator is called the Urysohn integral operator.

In the paper [1], new interpolation theorems were proved for these operators in Morrey-type spaces. Analogs of the interpolation theorems of Marcinkiewicz-Calderon, Stein-Weiss, Petre were obtained.

In this paper, we study the interpolation properties of the net spaces $N_{p,q}(M)$. Also, we prove a certain analogue of the Marcinkiewicz-type interpolation theorem for the Urysohn operator (1). We use the ideas developed in [1-3], where an interpolation theorem of Marcinkiewicz type for Morrey spaces was obtained.

Let in \mathbb{R}^n is given n -dimensional Lebesgue measure μ , M is an arbitrary system of measurable subsets from \mathbb{R}^n . For a function $f(x)$, defined and integrable on each e from M , we define the function

$$\bar{f}(t, M) = \sup_{\substack{e \in M \\ |e| \geq t}} \frac{1}{|e|} \left| \int_e f(x) dx \right|, \quad t > 0,$$

where the supremum is taken over all $e \in M$, whose measure is $|e| \stackrel{\text{def}}{=} \mu e \geq t$. In the case, when $\sup\{|e| : e \in M\} = \alpha < \infty$ and $t > \alpha$ assuming that $\bar{f}(t, M) = 0$.

Let p, q parameters satisfy the conditions $0 < p \leq \infty$, $0 < q \leq \infty$. We define the net spaces $N_{p,q}(M)$, as the set of all functions f , such that for $q < \infty$

$$\|f\|_{N_{p,q}(M)} = \left(\int_0^\infty \left(t^{\frac{1}{p}} \bar{f}(t, M) \right)^q \frac{dt}{t} \right)^{\frac{1}{q}} < \infty,$$

*Corresponding author.

E-mail: aitolkynnur@gmail.com

and for $q = \infty$

$$\|f\|_{N_{p,\infty}(M)} = \sup_{t>0} t^{\frac{1}{p}} \bar{f}(t, M) < \infty.$$

These spaces were introduced in the work [4]. Net spaces have found important applications in various problems of harmonic analysis, operator theory and the theory of stochastic processes [5–13].

Marcinkiewicz-type interpolation theorem for Urysohn operators

A family of measurable sets $G = \{G_t\}_{t>0}$ is called a local net if it satisfies the following conditions $G_t \subset G_s$ for $t \leq s$ and $|G_t| = t$. An example of a local net is the set $\{B_t(x)\}_{t>0}$ of all balls centered at the point x .

Let $G = \{G_t\}_{t>0}$ be a local net. We define the net $F_{G,\Omega} = \bigcup_{x \in \Omega} G + x$, where $G + x = \{G_t + x\}_{t>0}$. The net $F_{G,\Omega}$ will be called the generated by local net G and the set Ω .

Example. Let $\Omega = \mathbb{R}^n$, $G = \{Q_t\}_{t>0}$ be a set of cubes centered at 0, then $F_{G,\Omega} = \{Q_t + x\}_{t>0, x \in \mathbb{R}^n}$ is the set of all cubes in \mathbb{R}^n .

Lemma 1. Let T be the Urysohn operator of the form (1), then for an arbitrary function $f \in Z(U)$ from the domain and for any μ measurable set $w \subset U$ the following condition holds:

$$T(f, y) = T(f\chi_w, y) + T(f\chi_{U \setminus w}, y) - T(0, y).$$

Proof. Due to the additivity of the integral with respect to measure

$$\begin{aligned} T(f, y) - T(f\chi_w, y) &= \int_U K(f(x), xy) d\mu - \int_U K(f(x)\chi_w(x), x, y) d\mu \\ &= \int_{U \setminus w} K(f(x)\chi_{U \setminus w}(x), x, y) d\mu + \int_w K(f(x)\chi_w(x), x, y) d\mu \\ &\quad - \int_{U \setminus w} K(0, x, y) d\mu - \int_w K(f(x)\chi_w(x), x, y) d\mu \\ &= \int_U K(f(x)\chi_{U \setminus w}(x), x, y) d\mu - \int_w K(0, x, y) d\mu - \int_{U \setminus w} K(0, x, y) d\mu \\ &= \int_U K(f(x)\chi_{U \setminus w}(x), x, y) d\mu - \int_U K(0, x, y) d\mu = T(f\chi_{U \setminus w}, y) - T(0, y). \end{aligned}$$

Lemma 2. (Hardy’s inequalities) Let $\mu > 0, -\infty < \nu < \infty$ and $0 < \sigma \leq \tau \leq \infty$, then the following inequalities hold

$$\left(\int_0^\infty \left(y^{-\mu} \left(\int_0^y \left(r^{-\nu} |g(r)| \right)^\sigma \frac{dr}{r} \right)^{\frac{1}{\sigma}} \right)^\tau \frac{dy}{y} \right)^{\frac{1}{\tau}} \leq (\mu\sigma)^{-\frac{1}{\sigma}} \left(\int_0^\infty \left(y^{-\mu-\nu} |g(y)| \right)^\tau \frac{dy}{y} \right)^{\frac{1}{\tau}}$$

and

$$\left(\int_0^\infty \left(y^\mu \left(\int_0^y \left(r^{-\nu} |g(r)| \right)^\sigma \frac{dr}{r} \right)^{\frac{1}{\sigma}} \right)^\tau \frac{dy}{y} \right)^{\frac{1}{\tau}} \leq (\mu\sigma)^{-\frac{1}{\sigma}} \left(\int_0^\infty \left(y^{\mu-\nu} |g(y)| \right)^\tau \frac{dy}{y} \right)^{\frac{1}{\tau}}.$$

Theorem 1. Let $\Omega \subset \mathbb{R}^n$, $G = \{G_t\}_{t>0}$ is the local net, $F = \bigcup_{x \in \Omega} G + x$. Let $0 < p_0 < p_1 < \infty$ and $0 < q_0, q_1 \leq \infty, q_0 \neq q_1, 0 < \theta < 1, 1 \leq \tau \leq \infty$,

$$\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}, \quad \frac{1}{q} = \frac{1-\theta}{q_0} + \frac{\theta}{q_1}.$$

If for the Urysohn operator T and some $M_0, M_1 > 0$ the following inequalities hold

$$\|T(f) - T(0)\|_{N_{q_i,\infty}(G+x)} \leq M_i \|f\|_{N_{p_i,1}(G+x)}, \quad i = 0, 1, \quad x \in \Omega, \tag{2}$$

then

$$\|T(f) - T(0)\|_{N_{q,\tau}(F)} \leq c M_0^{1-\theta} M_1^\theta \|f\|_{N_{p,\tau}(F)}, \tag{3}$$

for all functions $f \in N_{p,\tau}(F)$, where $c > 0$ depends only on the parameters $p_0, p_1, q_0, q_1, p, q, \tau, \theta$.

Proof.

Let $1 \leq \tau < \infty$, $f \in N_{p,\tau}(F)$, for arbitrary $x \in \Omega$, $s > 0$, we define the functions

$$f_{0,s} = f\chi_{G_s+x}, \quad f_{1,s} = f - f_{0,s},$$

where χ_{G_s+x} denotes the characteristic function of the set $G_s + x$. It is easily seen that $f_{0,s} \in N_{p_0,1}$ and $f_{1,s} \in N_{p_1,1}$. Then $f = f_{0,s} + f_{1,s}$ and

$$\begin{aligned} & \sup_{\xi \geq t} \frac{1}{|G_\xi|} \left| \int_{G_\xi+x} (T(f, y) - T(0, y)) dy \right| \\ &= \sup_{\xi \geq t} \frac{1}{|G_\xi|} \left| \int_{G_\xi+x} (T(f\chi_{G_s+x}, y) + T(f\chi_{\mathbb{R}^n \setminus G_s+x}, y) - 2T(0, y)) dy \right| \\ &\leq \sup_{\xi \geq t} \frac{1}{|G_\xi|} \left| \int_{G_\xi+x} (T(f_{0,s}, y) - T(0, y)) dy \right| \\ &+ \sup_{\xi \geq t} \frac{1}{|G_\xi|} \left| \int_{G_\xi+x} (T(f_{1,s}, y) - T(0, y)) dy \right| = I_1 + I_2. \end{aligned}$$

First, we estimate I_1 , according to the inequality (2) we have

$$\begin{aligned} I_1 &= \sup_{\xi \geq t} \frac{1}{|G_s|} \left| \int_{G_\xi+x} (T(f_{0,s}, y) - T(0, y)) dy \right| \\ &\leq t^{-\frac{1}{q_0}} \sup_{r>0} r^{\frac{1}{q_0}} \sup_{\xi \geq r} \frac{1}{|G_\xi|} \left| \int_{G_\xi+x} (T(f_{0,s}, y) - T(0, y)) dy \right| \\ &= t^{-\frac{1}{q_0}} \|(T(f_{0,s}, y) - T(0, y))\|_{N_{q_0,\infty}(G+x)} \leq M_0 t^{-\frac{1}{q_0}} \|f_{0,s}\|_{N_{p_0,1}(G+x)} \\ &= M_0 t^{-\frac{1}{q_0}} \left(\int_0^s r^{\frac{1}{p_0}} \sup_{\xi \geq r} \frac{1}{|G_\xi|} \left| \int_{G_\xi+x} f_{0,s}(y) dy \right| \frac{dr}{r} + \int_s^\infty r^{\frac{1}{p_0}} \sup_{\xi \geq r} \frac{1}{|G_\xi|} \left| \int_{G_\xi+x} f_{0,s}(y) dy \right| \frac{dr}{r} \right). \end{aligned}$$

Let $0 < r \leq s$, if $\xi \leq s$, $y \in G_\xi + x$, we have $f_{0,s}(y) = f(y)\chi_{G_s+x} = f(y)$, if $\xi > s$, then

$$\left| \int_{G_\xi+x} f_{0,s}(y) dy \right| = \left| \int_{G_s+x} f(y) dy \right|.$$

By the first integral, we have the following,

$$\begin{aligned} & \int_0^s r^{\frac{1}{p_0}} \sup_{\xi \geq r} \frac{1}{|G_\xi|} \left| \int_{G_\xi+x} f_{0,s}(y) dy \right| \frac{dr}{r} = \int_0^s r^{\frac{1}{p_0}} \sup_{s \geq \xi \geq r} \frac{1}{|G_\xi|} \left| \int_{G_\xi+x} f(y) dy \right| \frac{dr}{r} \\ &\leq \int_0^s r^{\frac{1}{p_0}} \sup_{\xi \geq r} \frac{1}{|G_\xi|} \left| \int_{G_\xi+x} f(y) dy \right| \frac{dr}{r} \leq \int_0^s r^{\frac{1}{p_0}} \bar{f}(r, F) \frac{dr}{r}. \end{aligned}$$

By the second integral, we have

$$\begin{aligned} & \int_s^\infty r^{\frac{1}{p_0}} \sup_{\xi \geq r} \frac{1}{|G_\xi|} \left| \int_{G_\xi+x} f_{0,s}(y) dy \right| \frac{dr}{r} = \int_s^\infty r^{\frac{1}{p_0}} \sup_{\xi \geq r} \frac{1}{|G_\xi|} \left| \int_{G_s+x} f(y) dy \right| \frac{dr}{r} \\ &= \left| \int_{G_s+x} f(y) dy \right| \int_s^\infty r^{\frac{1}{p_0}} \sup_{\xi \geq r} \frac{1}{|G_\xi|} \frac{dr}{r} = \left| \int_{G_s+x} f(y) dy \right| \int_s^\infty r^{\frac{1}{p_0}-1} \frac{dr}{r} \\ &= p'_0 s^{\frac{1}{p_0}} \frac{1}{|G_s|} \left| \int_{G_s+x} f(y) dy \right| \leq p'_0 s^{\frac{1}{p_0}} \bar{f}(s, F). \end{aligned}$$

Thus, we get

$$I_1 \lesssim M_0 t^{-\frac{1}{q_0}} \left(\int_0^s r^{\frac{1}{p_0}} \bar{f}(r, F) \frac{dr}{r} + s^{\frac{1}{p_0}} \bar{f}(s, F) \right).$$

We estimate I_2 in a similar way applying the inequality (2), we obtain

$$\begin{aligned}
 I_2 &= \sup_{s \geq t} \frac{1}{|G_s|} \left| \int_{G_{s+x}} (T(f_{1,s}, y) - T(0, y)) dy \right| \\
 &\leq t^{-\frac{1}{q_1}} \sup_{r > 0} r^{\frac{1}{q_1}} \sup_{s \geq r} \frac{1}{|G_s|} \left| \int_{G_{s+x}} (T(f_{1,s}, y) - T(0, y)) dy \right| \\
 &= t^{-\frac{1}{q_1}} \|(T(f_{1,s}, y) - T(0, y))\|_{N_{q_1, \infty}(G+x)} \leq M_1 t^{-\frac{1}{q_1}} \|f_{1,s}\|_{N_{p_1, 1}(G+x)} \\
 &= M_1 t^{-\frac{1}{q_1}} \left(\int_0^\infty r^{\frac{1}{p_1}} \sup_{s \geq r} \frac{1}{|G_s|} \left| \int_{G_{s+x}} f_{1,s}(y) dy \right| \frac{dr}{r} \right) \\
 &= M_1 t^{-\frac{1}{q_1}} \left(\int_0^s r^{\frac{1}{p_1}} \sup_{\xi \geq r} \frac{1}{|G_\xi|} \left| \int_{G_{\xi+x}} f_{1,s}(y) dy \right| \frac{dr}{r} + \int_s^\infty r^{\frac{1}{p_1}} \sup_{\xi \geq r} \frac{1}{|G_\xi|} \left| \int_{G_{\xi+x}} f_{1,s}(y) dy \right| \frac{dr}{r} \right) \\
 &= M_1 t^{-\frac{1}{q_1}} (J_1 + J_2).
 \end{aligned}$$

To estimate J_1, J_2 note that

$$\begin{aligned}
 \left| \int_{G_{\xi+x}} f_{1,s}(y) dy \right| &= \begin{cases} 0, & \xi \leq s, \\ \left| \int_{G_{\xi+x} \setminus G_{s+x}} f(y) dy \right|, & \xi > s \end{cases} \\
 &= \begin{cases} 0, & \xi \leq s, \\ \left| \int_{G_{\xi+x}} f(y) dy - \int_{G_{s+x}} f(y) dy \right| \leq \left| \int_{G_{\xi+x}} f(y) dy \right| + \left| \int_{G_{s+x}} f(y) dy \right|, & \xi > s. \end{cases}
 \end{aligned}$$

Further,

$$\begin{aligned}
 J_1 &\leq \int_0^s r^{\frac{1}{p_1}} \sup_{\xi \geq r} \frac{1}{|G_\xi|} \left(\left| \int_{G_{\xi+x}} f(y) dy \right| + \left| \int_{G_{s+x}} f(y) dy \right| \right) \frac{dr}{r} \\
 &\leq \int_0^s r^{\frac{1}{p_1}} \left(\bar{f}(s, F) + \left| \int_{G_{s+x}} f(y) dy \right| \sup_{\xi \geq r} \frac{1}{|G_\xi|} \right) \frac{dr}{r} \\
 &\leq 2\bar{f}(s, F) \int_0^s r^{\frac{1}{p_1}} \frac{dr}{r} = 2p_1 s^{\frac{1}{p_1}} \bar{f}(s, F),
 \end{aligned}$$

and

$$\begin{aligned}
 J_2 &\leq \int_s^\infty r^{\frac{1}{p_1}} \sup_{\xi \geq r} \frac{1}{|G_\xi|} \left(\left| \int_{G_{\xi+x}} f(y) dy \right| + \left| \int_{G_{s+x}} f(y) dy \right| \right) \frac{dr}{r} \\
 &\leq \int_s^\infty r^{\frac{1}{p_1}} \left(\bar{f}(s, F) + \left| \int_{G_{s+x}} f(y) dy \right| \sup_{\xi \geq r} \frac{1}{|G_\xi|} \right) \frac{dr}{r} \leq \int_s^\infty r^{\frac{1}{p_1}} \bar{f}(r, F) \frac{dr}{r} \\
 &\quad + \left| \int_{G_{s+x}} f(y) dy \right| \int_s^\infty r^{\frac{1}{p_1}} \sup_{\xi \geq r} \frac{1}{|G_\xi|} \frac{dr}{r} = \int_s^\infty r^{\frac{1}{p_1}} \bar{f}(r, F) \frac{dr}{r} \\
 &\quad + \left| \int_{G_{s+x}} f(y) dy \right| \frac{s^{\frac{1}{p_1}-1}}{p_1-1} p_1 \lesssim \int_s^\infty r^{\frac{1}{p_1}} \bar{f}(r, F) \frac{dr}{r} + s^{\frac{1}{p_1}} \bar{f}(s, F).
 \end{aligned}$$

Combining the estimates, we obtain the following estimate

$$I_2 \lesssim M_1 t^{\frac{1}{q_1}} \left(\int_s^\infty r^{\frac{1}{p_1}} \bar{f}(r, F) \frac{dr}{r} + s^{\frac{1}{p_1}} \bar{f}(s, F) \right).$$

So, we got the following estimate

$$\sup_{s \geq t} \frac{1}{|G_s|} \left| \int_{G_{s+x}} (T(f, y) - T(0, y)) dy \right| \lesssim M_0 t^{-\frac{1}{q_0}} \left(\int_0^s r^{\frac{1}{p_0}} \bar{f}(r, F) \frac{dr}{r} + s^{\frac{1}{p_0}} \bar{f}(s, F) \right)$$

$$+M_1 t^{-\frac{1}{q_1}} \left(\int_s^\infty r^{\frac{1}{p_1}} \bar{f}(r, F) \frac{dr}{r} + s^{\frac{1}{p_1}} \bar{f}(s, F) \right).$$

Assuming that $s = ct^\gamma$, where $\gamma = \left(\frac{1}{q_1} - \frac{1}{q_0} \right) / \left(\frac{1}{p_1} - \frac{1}{p_0} \right)$, then, taking into account the above estimates, we obtain

$$\begin{aligned} \|T(f) - T(0)\|_{N_{q,\tau}(F)} &= \left(\int_0^\infty \left(t^{\frac{1}{q}} \sup_{\substack{s \geq t \\ x \in \mathbb{R}^n}} \frac{1}{|G_s|} \left| \int_{G_{s+x}} f(x) dx \right| \right)^\tau \frac{dt}{t} \right)^{\frac{1}{\tau}} \\ &\lesssim M_0 A_1 + M_0 A_2 + M_1 A_3 + M_1 A_4, \end{aligned}$$

where

$$\begin{aligned} A_1 &= \left(\int_0^\infty \left(t^{\frac{1}{q} - \frac{1}{q_0}} \int_0^{ct^\gamma} r^{\frac{1}{p_0}} \bar{f}(r, F) \frac{dr}{r} \right)^\tau \frac{dt}{t} \right)^{\frac{1}{\tau}}, \\ A_2 &= \left(\int_0^\infty \left(t^{\frac{1}{q} - \frac{1}{q_0}} (ct^\gamma)^{\frac{1}{p_0}} \bar{f}(ct^\gamma, F) \right)^\tau \frac{dt}{t} \right)^{\frac{1}{\tau}}, \\ A_3 &= \left(\int_0^\infty \left(t^{\frac{1}{q} - \frac{1}{q_1}} \int_{ct^\gamma}^\infty r^{\frac{1}{p_1}} \bar{f}(r, F) \frac{dr}{r} \right)^\tau \frac{dt}{t} \right)^{\frac{1}{\tau}}, \\ A_4 &= \left(\int_0^\infty \left(t^{\frac{1}{q} - \frac{1}{q_1}} (ct^\gamma)^{\frac{1}{p_1}} \bar{f}(ct^\gamma, F) \right)^\tau \frac{dt}{t} \right)^{\frac{1}{\tau}}. \end{aligned}$$

Using the change of variable $ct^\gamma = y$, we get

$$\begin{aligned} A_1 &= \gamma^{-\frac{1}{\tau}} c^{-\theta \left(\frac{1}{p_1} - \frac{1}{p_0} \right)} B_1, \quad A_2 = \gamma^{-\frac{1}{\tau}} c^{-\theta \left(\frac{1}{p_1} - \frac{1}{p_0} \right)} B_2, \\ A_3 &= \gamma^{-\frac{1}{\tau}} c^{(1-\theta) \left(\frac{1}{p_1} - \frac{1}{p_0} \right)} B_3, \quad A_4 = \gamma^{-\frac{1}{\tau}} c^{(1-\theta) \left(\frac{1}{p_1} - \frac{1}{p_0} \right)} B_3, \end{aligned}$$

where

$$\begin{aligned} B_1 &= \left(\int_0^\infty \left(y^{\theta \left(\frac{1}{p_1} - \frac{1}{p_0} \right)} \int_0^y r^{\frac{1}{p_0}} \bar{f}(r, F) \frac{dr}{r} \right)^\tau \frac{dy}{y} \right)^{\frac{1}{\tau}}, \\ B_2 &= \left(\int_0^\infty \left(y^{\frac{1}{p}} \bar{f}(y, F) \right)^\tau \frac{dy}{y} \right)^{\frac{1}{\tau}} = \|f\|_{N_{p,\tau}(F)}, \\ B_3 &= \left(\int_0^\infty \left(y^{-(1-\theta) \left(\frac{1}{p_1} - \frac{1}{p_0} \right)} \int_y^\infty r^{\frac{1}{p_1}} \bar{f}(r, F) \frac{dr}{r} \right)^\tau \frac{dy}{y} \right)^{\frac{1}{\tau}}, \\ B_4 &= \left(\int_0^\infty \left(y^{\frac{1}{p}} \bar{f}(y, F) \right)^\tau \frac{dy}{y} \right)^{\frac{1}{\tau}} = \|f\|_{N_{p,\tau}(F)}. \end{aligned}$$

To estimate B_1, B_3 we apply Hardy's inequalities from the lemma 2 and we obtain

$$\begin{aligned} B_1 &\lesssim \left(\int_0^\infty \left(y^{\theta \left(\frac{1}{p_1} - \frac{1}{p_0} \right) + \frac{1}{p_0}} \bar{f}(r, F) \right)^\tau \frac{dy}{y} \right)^{\frac{1}{\tau}} \lesssim \|f\|_{N_{p,\tau}(F)}, \\ B_3 &\lesssim \left(\int_0^\infty \left(y^{(\theta-1) \left(\frac{1}{p_1} - \frac{1}{p_0} \right) + \frac{1}{p_1}} \bar{f}(r, F) \right)^\tau \frac{dy}{y} \right)^{\frac{1}{\tau}} \lesssim \|f\|_{N_{p,\tau}(F)}. \end{aligned}$$

Thus, from the above estimates, the following estimate was obtained

$$\|T(f) - T(0)\|_{N_{q,\tau}(F)} \lesssim \left(M_0 c^{-\theta \left(\frac{1}{p_1} - \frac{1}{p_0} \right)} + M_1 c^{(1-\theta) \left(\frac{1}{p_1} - \frac{1}{p_0} \right)} \right) \|f\|_{N_{p,\tau}(F)},$$

where the corresponding constants depend only on $p_0, p_1, q_0, q_1, p, q, \tau$ and θ .

Let $c = \left(\frac{M_1}{M_0}\right)^{\frac{p_0 p_1}{p_1 - p_0}}$, then

$$\|T(f) - T(0)\|_{N_{q,\tau}(F)} \lesssim M_0^{1-\theta} M_1^\theta \|f\|_{N_{p,\tau}(F)}.$$

Consequently, we obtain the required estimate (3). The theorem is proved.

Corollary. Let F be the set of all balls in \mathbb{R}^n , F_x – the set of all balls centered at the point $x \in \mathbb{R}^n$. Let $0 < p_0 < p_1 < \infty$ and $0 < q_0, q_1 \leq \infty$, $q_0 \neq q_1$, $0 < \theta < 1$, $1 \leq \tau \leq \infty$,

$$\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}, \quad \frac{1}{q} = \frac{1-\theta}{q_0} + \frac{\theta}{q_1}.$$

If the following inequalities hold for the Urysohn operator T and some $M_0, M_1 > 0$

$$\|T(f) - T(0)\|_{N_{q_i,\infty}(G+x)} \leq M_i \|f\|_{N_{p_i,1}(G+x)}, \quad i = 0, 1, \quad x \in \mathbb{R}^n,$$

then for all functions $f \in N_{p,\tau}(F)$, holds

$$\|T(f) - T(0)\|_{N_{q,\tau}(F)} \leq c M_0^{1-\theta} M_1^\theta \|f\|_{N_{p,\tau}(F)},$$

where $c > 0$ depends only on parameters $p_0, p_1, q_0, q_1, p, q, \tau, \theta$.

Acknowledgments

This research was supported by the Ministry of Education and Science of the Republic of Kazakhstan (Project No. AP08856479).

References

- 1 Буренков В.И. Интерполяционные теоремы для нелинейных операторов в общих пространствах типа Морри и их приложения / В.И. Буренков, Е.Д. Нурсултанов // Тр. МИАН. — 2021. — 312. — С. 124–149.
- 2 Burenkov V.I. Marcinkiewicz-type interpolation theorem and estimates for convolutions for Morrey-type spaces / V.I. Burenkov, D.K. Chigambayeva, E.D. Nursultanov // Eurasian Math. J. — 2018. — 9. — No. 2. — P. 82–88.
- 3 Burenkov V.I. Interpolation theorems for nonlinear urysohn integral operators in general Morrey-type spaces / V.I. Burenkov, E.D. Nursultanov // Eurasian Math. J. — 2020. — 11. No. 4. — P. 87–94.
- 4 Нурсултанов Е.Д. Сетевые пространства и неравенства типа Харди–Литтлвуда / Е.Д. Нурсултанов // Мат. сб. — 1998. — 189. — № 3. — С. 399–419.
- 5 Akylzhanov R. $L_p - L_q$ multipliers on locally compact groups / R. Akylzhanov, M. Ruzhansky // J. Fun. Anal. — 2020. — 278. — No. 3. — P. 108324.
- 6 Akylzhanov R. Net spaces on lattices, Hardy-Littlewood type inequalities, and their converses / R. Akylzhanov, M. Ruzhansky // Eurasian Math. J. — 2017. — 8. — No. 3. — P. 10–27.
- 7 Akylzhanov R. Hardy-Littlewood, Hausdorff-Young-Paley inequalities, and $L_p - L_q$ Fourier multipliers on compact homogeneous manifolds / R. Akylzhanov, M. Ruzhansky, E.D. Nursultanov // J. Math. Anal. Appl. — 2019. — 479. — No. 2. — P. 1519–1548.
- 8 Костюченко А.Г. Теория управления «катастрофами» / А.Г. Костюченко, Е.Д. Нурсултанов // УМН. — 1998. — 53. — № 3. — С. 628, 629.
- 9 Нурсултанов Е.Д. Нижние и верхние оценки нормы мультипликаторов кратных тригонометрических рядов Фурье в пространствах Лебега / Е.Д. Нурсултанов, Н.Т. Тлеуханова // Функци. анализ и его прил. — 2000. — 34. — № 2. — С. 151–153.
- 10 Нурсултанов Е.Д. О коэффициентах кратных рядов Фурье из L_p -пространств / Е.Д. Нурсултанов // Изв. РАН. Сер. мат. — 2000. — 64. — № 1. — С. 95–122.
- 11 Нурсултанов Е.Д. О применении интерполяционных методов к исследованию свойств функций многих переменных / Е.Д. Нурсултанов // Мат. заметки. — 2004. — 75. — № 3. — С. 372–383.

- 12 Nursultanov E.D. Net spaces and boundedness of integral operators / E.D. Nursultanov, S. Tikhonov // Geom. Anal. — 2011. — 21. — P. 950–981.
- 13 Nursultanov E.D. Interpolation methods for stochastic processes spaces / E.D. Nursultanov, T.U. Aubakirov // Abstr. Appl. Anal. — 2013. — 2013. — P. 1–12.

А.Х. Калидолдай¹, Е.Д. Нұрсұлтанов²

¹Л.Н.Гумилев атындағы Еуразия ұлттық университеті, Нұр-Сұлтан, Қазақстан;

²М.В. Ломоносов атындағы ММУ Қазақстан филиалы, Нұр-Сұлтан, Қазақстан

Сызықтық емес интегралды Урысон операторларының торлы кеңістіктеріндегі интерполяциясы

Мақалада $N_{p,q}(M)$ торлы кеңістіктерінің интерполяциялық қасиеттері зерттелген, мұндағы $M - \mathbb{R}^n$ жиынының өлшенетін ішкі жиындардың жеткілікті жалпы ерікті жүйесі. Интегралды Урысон операторы қарастырылған. Бұл оператор барлық сызықтық, интегралды операторларды, сондай-ақ сызықты емес интегралды операторларды жалпылайды. Урысон операторы, әдетте, квазисызықты немесе субаддитивті оператор емес, сондықтан бұл операторлар үшін классикалық интерполяциялық теоремалар орындалмайды. Осы операторлар класы үшін Марцинкевич типіндегі интерполяциялық теоремасының белгілі бір аналогы алынды. Бұл теорема белгілі бір мағынада локальды торы бар торлы кеңістіктердегі Урысон операторлары үшін әлсіз бағалаулардан торлы кеңістіктердегі осы операторлар үшін күшті бағалау алуға мүмкіндік береді. Мысалы, тор \mathbb{R}^n -дегі барлық шарлар жиынтығы болатын торлы кеңістігінде Урысон интегралды операторы болу үшін, оның тор концентрлі шарлар болатын торлы кеңістіктер үшін әлсіз типті болуы жеткілікті.

Кілт сөздер: интерполяция кеңістіктері, торлы кеңістіктер, Урысон операторлары.

А.Х. Калидолдай¹, Е.Д. Нұрсұлтанов²

¹Евразийский национальный университет имени Л.Н. Гумилева, Нур-Султан, Казахстан;

²Казахстанский филиал Московского государственного университета им. М.В. Ломоносова, Нур-Султан, Казахстан

Интерполяция нелинейных интегральных операторов Урысона в сетевых пространствах

В статье изучены интерполяционные свойства сетевых пространств $N_{p,q}(M)$, в случае когда M есть достаточно общая произвольная система измеримых подмножеств из \mathbb{R}^n . Рассмотрен интегральный оператор Урысона. Данный оператор обобщает все линейные, интегральные, а также нелинейные интегральные операторы. Оператор Урысона, вообще говоря, не является квазилинейным либо субаддитивным, поэтому классические интерполяционные теоремы для этих операторов не имеют места. Получен некий аналог интерполяционной теоремы типа Марцинкевича для этого класса операторов. Настоящая теорема позволяет получать в некотором смысле сильную оценку для операторов Урысона в сетевых пространствах из слабых оценок для них. Так, например, для того, чтобы был интегральный оператор Урысона в сетевом пространстве, где сеть есть множество всех шаров в \mathbb{R}^n , достаточно, чтобы он был слабого типа для сетевых пространств, где сеть есть концентрические шары.

Ключевые слова: интерполяционные пространства, сетевое пространство, операторы Урысона, интегральный оператор.

References

- 1 Burenkov, V.I., & Nursultanov, E.D. (2021). Interpoliatsionnye teoremy dlia nelineinykh operatorov v obshchikh prostranstvakh tipa Morri i ikh prilozheniia [Interpolation Theorems for Nonlinear Operators in General Morrey-Type Spaces and Their Applications]. *Trudy MIAN — Proc. Steklov Inst. Math.*, 312, 124–149 [in Russian].
- 2 Burenkov, V.I., Chigambayeva, D.K., & Nursultanov, E.D. (2018). Marcinkiewicz-type interpolation theorem and estimates for convolutions for Morrey-type spaces. *Eurasian Math. J.*, 9 (2), 82–88.
- 3 Burenkov, V. I., & Nursultanov, E.D. (2020). Interpolation theorems for nonlinear urysohn integral operators in general Morrey-type spaces. *Eurasian Math. J.*, 11 (4), 87–94 .
- 4 Nursultanov, E.D. (1998). Setevye prostranstva i neravenstva tipa Khardi–Littlvuda [Net spaces and inequalities of Hardy–Littlewood type]. *Matematicheskii sbornik*, 189 (3), 399–419 [in Russian].
- 5 Akylzhanov, R., & Ruzhansky, M. (2020). $L_p - L_q$ multipliers on locally compact groups. *J. Fun. Anal.*, 278 (3), 108324.
- 6 Akylzhanov, R., & Ruzhansky, M. (2017). Net spaces on lattices, Hardy–Littlewood type inequalities, and their converses. *Eurasian Math. J.*, 8 (3), 10–27.
- 7 Akylzhanov, R., Ruzhansky, M., & Nursultanov, E.D. (2019). Hardy–Littlewood, Hausdorff–Young–Paley inequalities, and $L_p - L_q$ Fourier multipliers on compact homogeneous manifolds. *J. Math. Anal. Appl.*, 479 (2), 1519–1548.
- 8 Nursultanov, E.D., & Kostyuchenko, A.G. (1998). Teoriia upravleniia «katastrofami» [Theory of control of “catastrophes”]. *Russ. Math. Surv.*, 53 (3), 628–629 [in Russian].
- 9 Nursultanov, E.D., & Tleukhanova, N.T. (2000). Nizhnie i verkhnie otsenki normy multiplikatorov kratnykh trigonometricheskikh riadov Fure v prostranstvakh Lebega [Lower and upper bounds for the norm of multipliers of multiple trigonometric Fourier series in Lebesgue spaces]. *Func. Anal. Appl.*, 34 (2), 151–153 [in Russian].
- 10 Nursultanov, E.D. (2000). O koeffitsientakh kratnykh riadov Fure iz L_p -prostranstv [On the coefficients of multiple Fourier series in L_p - spaces]. *Izvestiia RAN. Serii Matematika — Izvestiia Mathematics*, 64 (1), 93–120 [in Russian].
- 11 Nursultanov, E.D. (2004). O primeneniі interpoliatsionnykh metodov k issledovaniu svoistv funktsii mnogikh peremennykh [Application of Interpolational Methods to the Study of Properties of Functions of Several Variables]. *Matematicheskii zametki — Mathematical Notes*, 75, 341–351 [in Russian].
- 12 Nursultanov, E.D., & Tikhonov, S. (2011). Net spaces and boundedness of integral operators. *J. Geom. Anal.*, 21, 950–981.
- 13 Nursultanov, E.D., & Aubakirov, T.U. (2013). Interpolation methods for stochastic processes spaces. *Abstr. Appl. Anal.*, 2013, 1–12.