

## Punctual numberings for families of sets

A. Askarbekkyzy<sup>1</sup>, R. Bagaviev<sup>2</sup>, V. Isakov<sup>3</sup>, B. Kalmurzayev<sup>1</sup>, D. Nurlanbek<sup>1,\*</sup>,  
F. Rakymzhankyzy<sup>1</sup>, A. Slobozhanin<sup>3</sup>

<sup>1</sup>Kazakh-British Technical University, Almaty, Kazakhstan;

<sup>2</sup>Kazan Federal University, Kazan, Russia;

<sup>3</sup>Novosibirsk State University, Novosibirsk, Russia

(E-mail: [ms.askarbekkyzy@gmail.com](mailto:ms.askarbekkyzy@gmail.com), [nurlanbek.dias21@gmail.com](mailto:nurlanbek.dias21@gmail.com), [fariza.rakymzhankyzy@gmail.com](mailto:fariza.rakymzhankyzy@gmail.com),  
[birzhan.kalmurzayev@gmail.com](mailto:birzhan.kalmurzayev@gmail.com), [v.isakov@q.nsu.ru](mailto:v.isakov@q.nsu.ru), [a.slobozhanin@q.nsu.ru](mailto:a.slobozhanin@q.nsu.ru), [ramilbagaviev@mail.ru](mailto:ramilbagaviev@mail.ru))

This work investigates the structure of punctual numberings for families of punctually enumerable sets with respect to primitive recursively reducibility. We say that a numbering of a certain family is primitive recursively reducible to another numeration of the same family if there exists a primitive recursively procedure (an algorithm not employing unbounded search) mapping the numbers of objects in the first numbering to the numbers of the same objects in the second numbering. This study was motivated by the work of Bazhenov, Mustafa, and Ospichev on punctual Rogers semilattices for families of primitive recursively enumerable functions. The concept of punctually enumerable sets was introduced in the paper, and it was proven that not all recursively enumerable sets are punctually enumerable, but in all  $m$ -degrees, recursively enumerable sets include punctually enumerable sets. For two-element families of punctual sets, it was demonstrated that punctual Rogers semilattices can be of at least three types: (1) one-element family, (2) isomorphic to the upper semilattice of recursively enumerable sets with respect to primitive recursively  $m$ -reducibility, (3) without the greatest element. It was also proven that the set of all punctually enumerable sets does not have a punctual numbering, and punctual families with a Friedberg numbering do not have the least numbering.

*Keywords:* primitive recursive functions, punctually enumerable sets, Rogers semilattice, quick functions, punctual numberings.

*2020 Mathematics Subject Classification:* 03D25, 03D30.

### Introduction

Theory of computable numberings is one of the actively developing areas in the computability theory. *Numbering* of a countable set  $S$  is any surjective mapping  $\nu : \omega \rightarrow S$  (Here and further as  $\omega$  we denote the set of natural numbers). A numbering  $\nu$  is called computable if the set

$$\{(n, x) : n \in \omega, x \in \nu(n)\}$$

is computably enumerable (c.e.) set.

The set of all computable numberings for family  $\mathcal{S}$  denotes as  $\text{Com}(\mathcal{S})$ . Let  $\nu$  and  $\mu$  are numberings for family  $S$ . Numbering  $\nu$  is *reducible* to  $\mu$  if there is computable function  $f$  such that  $\nu = \mu \circ f$  (denotes  $\nu \leq \mu$ ). This reducibility induces a partially preordered set structure, which factor structure is called *Rogers semilattice for family S* and denoted as  $\mathcal{R}(\mathcal{S})$ .

There are several interesting results known about Rogers semilattice. For example, if  $\mathcal{S}$  is a family of c.e. sets, then either  $|\mathcal{R}(\mathcal{S})| = 1$  or  $|\mathcal{R}(\mathcal{S})| = \infty$  [1]. In the case when  $|\mathcal{R}(\mathcal{S})| = \infty$  the semilattice is not

\*Corresponding author. E-mail: [nurlanbek.dias21@gmail.com](mailto:nurlanbek.dias21@gmail.com)

The work of A. Askarbekkyzy, B. Kalmurayev, D. Nurlanbek, F. Rakymzhankyzy was supported by the Ministry of Science and Higher Education of the Republic of Kazakhstan, grant AP19576325 “Algorithmic complexities of presentations for algebraic structures”.

*Received:* 16 January 2024; *Accepted:* 10 July 2024.

© 2024 The Authors. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

a lattice [2]. There is a universal numbering of all partial computable functions [3] and a computable numbering of all c.e. sets without repetitions [4] (these numberings are called friedberg numbering). For more information about the properties of the classical Rogers semilattice, refer to the following articles: [5–8].

In recent years, under the influence of the work [9], interest in primitive recursive (or punctual) properties of algebraic structures has increased. The next articles will help you to find more information about punctual structures [10–27]. In connection with this, studying the punctual properties of numberings is also relevant. Bazhenov, Mustafa, and Ospichev considered punctual numberings of families of functions in their article [28]. The authors established punctual reducibility between numberings, induced by primitive recursive functions, leading to the creation of upper semilattices of degrees known as Rogers *pr*-semilattices. They demonstrated that any infinite, uniformly primitive recursive family  $S$  induces an infinite Rogers *pr*-semilattice  $\mathcal{R}$ . It was proven that the semilattice  $\mathcal{R}$  is downwards dense, with every nontrivial interval within  $\mathcal{R}$  containing an infinite antichain. Additionally, every non-greatest element in  $\mathcal{R}$  is a part of an infinite antichain. The authors showed that the  $\Sigma_1$ -fragment of the theory  $Th(\mathcal{R})$  is decidable. Several examples were provided to emphasize the contrasts between the punctual framework and the classical theory of computable numberings. Notably, it was demonstrated that some infinite Rogers *pr*-semilattices  $\mathcal{R}$  are lattices, while others are not. The authors obtained a series of results concerning special classes of punctual numberings, including Friedberg numberings and decidable numberings with primitive recursive numeration equivalence. This paper is a logical continuation of the article [28] and aims to investigate punctual numberings for families of sets.

In Chapter 2, we introduce punctual analogs of concepts standard in the theory of computable numberings and define the punctual Rogers semilattice for sets. Chapter 3 is devoted to the structural properties of c.e. degrees induced by the restriction of  $m$ -reducibility by primitive recursive functions (called *prm*-reducibility). Chapters 4 and 5 present some properties of the punctual Rogers semilattice for finite and infinite families, including its connection with the structure of c.e. *prm*-degrees.

We adhere to the notations and terminology adopted in [29, 30]. We denote by  $\{p_e\}_{e \in \omega}$  the computable numbering of all primitive recursive functions. In this article we will consider restricted Church-Turing thesis for primitive recursive functions. We can define this thesis as follows: a function is primitive recursive if and only if it can be described by an algorithm that uses only bounded loops. More about restricted Church-Turing thesis you can find in the work [31].

### 1 Punctually enumerable sets and numberings

In the paper [28], the numbering  $\nu$  of a family of primitive recursive functions is called “punctual” if the function  $g_\nu(n, x) := (\nu(n))(x)$  is primitive recursive. It seems natural to attempt to extend this definition to a family of c.e. sets, but here we face some difficulties.

The thing is, such a definition of punctual numbering yields the same class of computable numberings for families of c.e. sets because any c.e. set can be represented as the range of a primitive recursive function, which means that a family can be enumerated in a punctual way. On the other hand, even with the presence of a punctual enumeration of a c.e. set, it is not always possible to use primitive recursive constructions, for example, due to the unbounded repetition of elements in the enumeration. In this regard, it makes sense to consider families of sets with stricter enumeration constraints than c.e. sets.

*Definition 1.* A set  $A$  is called *punctually enumerable*, if there is a primitive recursive function  $p$ , such that

- 1)  $A = \text{range}(p)$ , and
- 2) If  $p(x) = p(y)$  for some  $x < y$ , then  $\text{range}(p) = \{p(0), p(1), \dots, p(x)\}$ .

We will call the function  $p$  as a *quick* function for  $A$ .

Thus, the quick function  $p$  from the definition is injective for an infinite set  $A$ , and for a finite set it eventually enumerates all of its elements and then starts repeating them.

*Definition 2.* A numbering  $\nu$  of a family  $\mathcal{S}$  of punctually enumerable sets is called *punctual*, if there exists a primitive recursive function  $g_\nu(n, x)$  such that  $\lambda x.g_\nu(n, x)$  is a quick function for  $\nu(n)$  for any  $n$ . The set of all punctual numberings for family  $\mathcal{S}$  we will denote as  $\text{Com}_{pr}(\mathcal{S})$ .

Reductions on numberings are defined analogously to [28].

*Definition 3.* We say that numbering  $\nu$  is *punctually reducible* to numbering  $\mu$  (denoted as  $\nu \leq_{pr} \mu$ ), if there is primitive recursive function  $f$  such that  $\nu = \mu \circ f$ .

Numberings  $\nu$  and  $\mu$  are punctually equivalent and denote as  $\nu \equiv_{pr} \mu$ , if  $\nu \leq_{pr} \mu$  and  $\mu \leq_{pr} \nu$ .

As in the computable case, the least upper bound of the numberings  $\nu$  and  $\mu$  is the numbering  $\nu \oplus \mu$ , which is defined as

$$(\nu \oplus \mu)(2x) = \nu(x), (\nu \oplus \mu)(2x + 1) = \mu(x).$$

As punctual Rogers semilattice of the family of punctually enumerable sets  $\mathcal{S}$ , we will call partially ordered set  $\mathcal{R}_{pr}(\mathcal{S}) = (\text{Com}_{pr}(\mathcal{S}) / \equiv_{pr}, \leq_{pr}, \oplus)$ .

The following theorem demonstrates the independence of the concepts of primitive recursive set (having primitive recursive characteristic function) and punctually enumerable.

*Theorem 1.* There exist sets  $A$  and  $B$  such that  $A$  is punctually enumerable but not primitive recursive, and  $B$  is primitive recursive but not punctually enumerable.

*Proof.* As a set  $A$  we can choose the set  $K \oplus \omega$ . For this set its quick function we can construct as follows: we fix a primitive recursive approximation of the creative set  $K$ , denoted  $K_i$ , which at each step enumerates at most one element. Then, we set  $f(0) = 1$ , and  $f(x) = 2s$ , where  $s \in K_x \setminus K_{x-1}$ , if such  $s$  exists. If there is no such  $s$ , then  $f(x)$  is defined as the smallest odd number that has not been used before. It is clear that  $f$  is injective primitive recursive function and  $A$  is the range of  $f$ .

For set  $B$  we will construct its primitive recursive characteristic function  $\phi$  such that for  $B$  there is no quick function. We fix a computable numbering of all injective primitive recursive functions with the following condition

$$i_e(x)[t] \downarrow = a \Rightarrow a < t.$$

We will define  $B$  as a infinite set. So, it is sufficient that there is no injective quick function for  $B$ .

At step  $s$  we will define  $\phi(s)$  as follow: Assume that  $k$  is the cardinality of the set  $\{t : \phi(t) = 1 \ \& \ t < s\}$ . If there is more that  $k$  elements  $x \leq s$  such that  $i_k(x)[s] \downarrow$ , then define  $\phi(s) = 1$ . Otherwise, define  $\phi(s) = 0$ .

It is not hard to see that  $B$  is infinite set and any  $i_k$  can not enumerate  $B$ .

## 2 The structure of pr $m$ -degrees

In recent work [32] considered a many-one reductions for computable sets under primitive recursive functions, and have been proven that first-order theory of upper semilattice of degrees of computable sets with respect to primitive recursive many-one reducibility is hereditarily undecidable.

*Definition 4.* [32] The set  $A$  is *pr $m$ -reducible* to the set  $B$  (written as  $A \leq_m^{pr} B$ ), if there exists a primitive recursive function  $f$  such that  $A \leq_m B$  via  $f$ .

*Remark 1.* The computable  $m$ -degree contains infinitely many *pr $m$ -degrees*.

*Theorem 2.* For any c.e., but not computable set  $A$ , there is c.e. set  $B$  such that  $A \equiv_m B$  and  $A \not\leq_m^{pr} B$ .

*Proof.* Let's define computable majorant for all primitive recursive functions:

$$f(0) = p_0(0) + 1,$$

$$f(x + 1) = \max_{i,j \leq x+1} \{p_i(j), f(x)\} + 1.$$

Note that  $f$  is not primitive recursive but the set  $\text{range}(f)$  is primitive recursive.

Let  $B = f(A) = \{f(x) : x \in A\}$ .

It is clear, that  $A \leq_m^f B$ , since  $f$  is strongly increasing, then  $x \in B \Leftrightarrow \exists n \leq x (n \in A \ \& \ f(n) = x)$ , which means that reverse reducibility is correct.

Let's show that if  $A \leq_m^{pr} B$ , then  $A$  is computable. We fix primitive recursive function  $p_e$ , which reduces  $A$  to  $B$ , and also step  $s$  such that  $A \upharpoonright e = A_s \upharpoonright e$  (here by  $A \upharpoonright e$  denotes the set  $\{x : x \in A \ \& \ x \leq e\}$ ).

Let  $x$  be an arbitrary number. If  $x \leq e$ , then  $x \in A \Leftrightarrow x \in A_s$ . Otherwise, we check the following condition:  $p_e(x) \in \text{range}(f)$ ? If it is not, then  $x \notin A$ . If it is, we effectively find  $z_0$  such that  $f(z_0) = p_e(x)$ .

Repeat for  $z_0$  same procedure as we did for  $x$ , and, if  $z_0 > e$ , then we find number  $z_1$  such that  $f(z_1) = p_e(z_0)$  and so on. As a result, we receive sequence  $(z_k)_k$ . Since  $f(z_0) = p_e(x) < f(x)$ , by definition of  $f$ , then  $z_0 < x$ , consequently, the sequence  $(z_k)_k$  decreases and we find  $k$ , such that  $p_e(z_k) \notin \text{range}(f)$  or  $z_k \leq e$ . Then  $x \in A \Leftrightarrow p_e(x) = f(z_0) \Leftrightarrow z_0 \in A \Leftrightarrow p_e(z_0) = f(z_1) \Leftrightarrow \dots \Leftrightarrow p_e(z_{k-1}) = f(z_k) \Leftrightarrow z_k \in A$ . If  $p_e(z_k) \notin \text{range}(f)$ , then  $z_k \notin A$ , otherwise  $z_k \leq e$  and  $z_k \in A \Leftrightarrow z_k \in A_s$ .

Thus, we can effectively define that  $x$  belongs to  $A$  or not.

*Corollary 1.* Every non-computable c.e.  $m$ -degree contains infinitely many  $prm$ -degrees.

*Proof.* Let  $A_0$  be non-computable c.e. set. By using the previous theorem, we will build c.e. set  $A_1$  such that  $A_0 \equiv_m A_1$  and  $A_0 \not\leq_m^{pr} A_1$ , for  $A_1$  similarly build  $A_2$ , and for  $A_2$  build  $A_3$  and so on. All sets  $A_n, n \in \omega$  are  $m$ -equivalent, and for  $i < j$  set  $A_i$   $m$ -reduces to  $A_j$  by  $f^{j-i}(x)$  ( $(j-i)$ -th composition of function  $f$  from the previous theorem), consequently,  $A_i \not\leq_m^{pr} A_j$ . Here, note that  $B = f(A) \leq_m^{pr} A$ . (Proof is similar).

### 3 Punctual semilattice of two-element families

In the work [28] it was shown that the punctual Rogers semilattice of a finite family of functions always has exactly one element. However, it turns out that this is not the case for families of sets.

In this chapter we assume, that  $\mathcal{S} = \{A, B\}$ , where  $A, B$  are different punctually enumerable sets.

Note that in this case  $\mathcal{R}_{pr}(\mathcal{S}) \neq \emptyset$ , since the function

$$\alpha(n)(x) = \begin{cases} f(x), & \text{for } n = 2k, \\ g(x), & \text{for } n = 2k + 1, \end{cases}$$

where  $f$  and  $g$  are quick functions for  $A$  and  $B$  respectively, gives the punctual numbering of the family  $\mathcal{S}$ .

*Proposition 1.* Let  $\mathcal{S}$  is punctual two-element family such that  $A$  or  $B$  is finite then  $|\mathcal{R}_{pr}(\mathcal{S})| = 1$ .

*Proof.* Let  $|A| = N \leq |B|$ ;  $f$  and  $g$  are quick functions of the sets  $A$  and  $B$ , respectively.

Let  $\nu, \mu \in \text{Com}_{pr}(\mathcal{S})$  are arbitrary and  $k_\nu, k_\mu$  their quick functions. Let's show that  $\nu \equiv_{pr} \mu$ .

Fix numbers  $a$  and  $b$  such that  $\mu(a) = A$  and  $\mu(b) = B$ . Then  $\nu \leq_{pr} \mu$  by primitive recursive function  $h$ , which defines as:

$$h(n) = \begin{cases} a, & \text{if } |k_\nu(n, \cdot) \upharpoonright N| \leq N, \\ b, & \text{otherwise.} \end{cases}$$

Really,  $|k_\nu(n, \cdot) \upharpoonright N| \leq N$  means that quick function  $k_\nu$  on  $N$ -th argument starts to repeat the values, consequently,  $|\nu(n)| \leq N$  and that's why  $\nu(n) = A = \mu(a)$ .

Reverse reducibility is proved similarly.

*Proposition 2.* Let  $\mathcal{S}$  be a two-element family such that  $A$  and  $B$  are infinite,  $A \cap B$  is finite and one of the sets is primitive recursive, then  $|\mathcal{R}_{pr}(\mathcal{S})| = 1$ .

*Proof.* Let  $|A \cap B| = N$  and  $A$  is primitive recursive.

Let's take two numberings  $\nu$  and  $\mu$  of the family  $\mathcal{S}$  and fix numbers  $a$  and  $b$  such that  $\mu(a) = A$  and  $\mu(b) = B$ .

We define

$$h(n) = \begin{cases} a, & \text{if } \exists x \in (k_\nu(n, \cdot) \upharpoonright N) \cap (A \setminus B), \\ b, & \text{otherwise.} \end{cases}$$

Note, that  $|k_\nu(n, \cdot) \upharpoonright N| > |A \cap B| \Rightarrow \exists x \in (k_\nu(n, \cdot) \upharpoonright N) \setminus (A \cap B)$ . We can check that  $x$  belongs to  $A$  by primitive recursive procedure, and  $\nu(n) = A \Leftrightarrow x \in A$ . Consequently,  $\nu \leq_{pr} \mu$  by function  $h$ . It is clear that reverse reducibility is true, then  $\nu \equiv_{pr} \mu$ .

*Theorem 3.* There exists family  $\mathcal{S} = \{A, B\}$ , where  $|A \cap B| < \infty$ , such that there is no universal numbering for  $\mathcal{S}$ .

*Proof.* We will build the sets  $A, B$  and numbering  $\alpha_e$  for family  $\mathcal{S} = \{A, B\}$ , satisfying the following requirements:

$$\mathcal{P}_{e,i} : \pi_e \in \text{Com}_{pr}(\mathcal{S}) \rightarrow \alpha_e \not\leq_{pr} \pi_e \text{ by function } p_i,$$

where  $\pi_e$  is computable numbering of all primitive recursive numberings,  $p_i$  is computable numbering of all primitive recursive functions. Let  $k_e$  be primitive recursive quick function for numbering  $\pi_e$ .

*Strategy for  $\mathcal{P}_{e,i}$ :*

- 1) Pick  $w_{e,i}$  – the least number, that we do not use before.
- 2) Wait until  $p_i(w_{e,i}) \downarrow$  and  $k_e(p_i(w_{e,i}), 0) \downarrow$  on the step  $t$ . While we are waiting, list to  $\alpha_e(w_{e,i})$  new numbers.
- 3) We perform one of the following cases:  
 Case 1: If  $k_e(p_i(w_{e,i}), 0) \in B$ , then all elements that we listed to  $\alpha_e(w_{e,i})$  until this step, we add to  $A$ . Also, we add to  $\alpha(w_{e,i})$  all elements from  $A$ . After this, we add to  $\alpha_e(w_{e,i})$  all elements that we add to  $A$ .  
 Case 2: If  $k_e(p_i(w_{e,i}), 0) \in A$ , then all elements that we listed to  $\alpha_e(w_{e,i})$  until this step, we add to  $B$ . Also, we add to  $\alpha(w_{e,i})$  all elements from  $B$ . After this, we add to  $\alpha_e(w_{e,i})$  all elements that we add to  $B$ .  
 Case 3: If  $k_e(p_i(w_{e,i}), 0) \notin A \cup B$ , then  $k_e(p_i(w_{e,i}), 0) \in B$  and return to the Case 1.

*Construction.* Fix effective linear order of requirements:

$$\mathcal{P}_{0,0} < \mathcal{P}_{1,0} < \mathcal{P}_{0,1} < \mathcal{P}_{2,0} < \mathcal{P}_{1,1} < \mathcal{P}_{0,2} < \dots$$

On step  $s$  of the construction we visit the first  $s$  strategies from the list. At every step, fresh numbers are selected and thrown into the sets  $A$  or  $B$ .

Let  $\pi_e \in \text{Com}_{pr}(\mathcal{S})$ , by this we build  $\alpha_e$  and let there is primitive recursive function  $p_i$ , such that  $\alpha_e \leq_{pr} \pi_e$  by function  $p_i$ . So the following should be performed:

$$\forall x[\alpha_e(x) = \pi_e(p_i(x))].$$

Let's check the witness  $w_{e,i}$  in  $\mathcal{P}_{e,i}$ . By construction, while  $p_i(w_{e,i})$  defines on step  $t$ , in  $\alpha_e(w_{e,i})$  we add new numbers. Suppose that on stage 3 the number  $k_e(p_i(w_{e,i}), 0)$  is in the set  $B$ , since  $A \cap B = \emptyset$ , the following performed  $\pi_e(p_i(w_{e,i})) = B$ . But, by construction  $\alpha_e(w_{e,i}) = A$ . This contradicts to reducibility  $\alpha_e$  to  $\pi_e$ .

For case, when the number  $k_e(p_i(w_{e,i}), 0)$  is in the set  $A$ , we do the same action. If number  $k_e(p_i(w_{e,i}), 0) \notin A \cup B$ , then the action is performed as in the first case.

*Proposition 3.* Let  $\mathcal{S}$  be two-elemented family, such that  $A \subset B$ , then  $|\mathcal{R}_{pr}(\mathcal{S})| = \infty$  with universal numbering.

*Proof.* Since the sets  $A$  and  $B$  are punctually enumerable, then there are functions  $p, q$  – quick functions for  $A$  and  $B$ , respectively.

We will build the numberings for family  $\mathcal{S}$  as follows:

Let  $W$  be arbitrary c.e. set, which is not empty and not  $\omega$ , then numbering  $\nu$  defines as:

$$\nu_W(x) = \begin{cases} A, & \text{if } x \notin W, \\ B, & \text{if } x \in W. \end{cases}$$

$$W = \cup_s W_s.$$

Quick function  $\lambda x.h(x, y)$  of the numbering  $\nu$  will add elements as:

On step 0.  $h_\nu(x, 0) = p(0)$ .

On step  $s$ .

1)  $h_\nu(x, s) = p(s)$ , if  $x \notin W_s$ ,

2)  $h_\nu(x, s) = q(\mu_{z \leq s+1}[q(z) \notin h_\nu(x, \cdot) \upharpoonright s])$ , if  $x \in W_s$ .

It is easy to check, that for an arbitrary  $W_i$ , which is not empty and not  $\omega$ , we can decide that  $\nu_{W_i} \in \text{Com}_{pr}(\mathcal{S})$ .

$W_i$  reduces to  $W_j$  by primitive recursive function if and only if  $\nu_{W_i}$  reduces by primitive recursive function to  $\nu_{W_j}$ .

Now, let  $\alpha$  be an arbitrary punctual numbering of the family  $\mathcal{S}$ . Then we can find c.e. set  $W_i$  such that  $\alpha = \nu_{W_i}$ . Since  $\alpha$  is numbering of the family  $\mathcal{S}$ , and  $W_i$  is not empty, then there is an element  $b \in B \setminus A$ , then we can define the function  $\varphi(x) = \mu_z[h_\alpha(x, z) = b]$ , which is range of c.e. set  $W_i$ . Which means that  $\mathcal{R}_{pr}(\mathcal{S})$  is isomorphic to  $\mathcal{L}_{prm}^0$ .

It is known, that the set  $K_0 = \{\langle x, y \rangle | x \in W_y\}$  is universal in  $\mathcal{L}_{prm}^0$ . Consequently,  $\nu_{K_0}$  is universal punctual numbering of the family  $\mathcal{S}$ .

Note that  $\mathcal{R}_{pr}(\mathcal{S})$  is isomorphic to the upper semilattice of all c.e. sets under  $pr$ -many-one reducibility. By [32] we can say that first-order theory of  $\mathcal{R}_{pr}(\mathcal{S})$  is undecidable.

*Proposition 4.* There exists the family  $\mathcal{S} = \{A, B\}$  such that  $|A \cap B| = \infty$  and  $|\mathcal{R}_{pr}(\mathcal{S})| = \infty$  without universal numbering.

*Proof.* Let  $A = \omega \setminus \{0\}$  and  $B = \omega \setminus \{1\}$ . And let  $\mu$  be an arbitrary numbering of the family  $\mathcal{S}$ . We will build the numbering  $\nu \in \text{Com}_{pr}(\mathcal{S})$  so that  $\nu \not\leq_{pr} \mu$ .

We will construct quick function  $q_\nu$  for  $\nu$  as follows:

$$q_\nu(w, y) = \begin{cases} y + 2, & \{0, 1\} \cap h_\mu(p_e(w) \downarrow, \cdot) \upharpoonright y = \emptyset; \\ 0, & h_\mu(p_e(w), y) = 1; \\ 1, & h_\mu(p_e(w), y) = 0; \\ y + 1, & \{0, 1\} \cap h_\mu(p_e(w) \downarrow, \cdot) \upharpoonright y \neq \emptyset, \end{cases}$$

where  $h$  is quick function of the numbering  $\mu$  and  $p_e$  is function that does not reduce  $\nu$  to  $\mu$ .

Let numbering  $\nu$  reduces to numbering  $\mu$  by primitive recursive function  $p$ , then there is  $w$  such that  $\nu(w) = 0$  and  $\mu(p(w)) = 1$  or  $\nu(w) = 1$  and  $\mu(p(w)) = 0$ . Contradiction.

Since  $\mu$  is an arbitrary numbering from the  $\text{Com}_{pr}(\mathcal{S})$ , then for any numbering from  $\text{Com}_{pr}(\mathcal{S})$  we can construct  $\nu$ , which does not reduce to  $\mu$ , which means that there is no universal numbering in this family.

#### 4 Punctual semilattice of the infinite families

*Theorem 4.* There is no punctual numbering for the family of all punctually enumerable sets.

*Proof.* Suppose that there is the numbering  $\nu$  for the family of all punctually enumerable sets. Let  $g(x, y)$  be a quick function for  $\nu$ . We construct punctually enumerable set  $A$  such that  $\nu(e) \neq A$  for all  $e \in \omega$ ; thus we will come to the contradiction.

Let  $q(0) = p(0, 0) + 1$  and  $q(n + 1) = \max(\{p(n + 1, z) : z \leq n + 1\} \cup \{q(n)\}) + 1$ .

It is clear that  $q$  is primitive recursive increasing function. Assume  $A = \text{range}(q)$ .

By contradiction, assume  $\nu(n) = A$  for some  $n \in \omega$ . Since,  $A$  is infinite, the function  $\lambda y.g(n, y)$  is injective. It is clear that the set  $\{p(n, z) : z \leq n\}$  has  $n+1$  different elements, hence in  $\{p(n, z) : z \leq n\}$  there is a number greater than  $q(n)$ . Contradiction.

*Corollary 2.* There is no punctual numbering for the family of all primitive recursive punctually enumerable sets.

The proof of corollary is the same as the proof of the theorem.

*Definition 5.* The numbering  $\nu \in \text{Com}_{pr}(\mathcal{S})$  is called *friedberg*, if it is injective.

*Proposition 5.* If the infinite family  $\mathcal{S}$  has friedberg numbering, then

- 1)  $\mathcal{R}_{pr}(\mathcal{S})$  does not have the least element,
- 2)  $|\mathcal{R}_{pr}(\mathcal{S})| = \infty$ .

*Proof.* 1) Let  $\nu$  be friedberg numbering for the infinite family  $\mathcal{S}$ . Suppose that  $\alpha$  is the least numbering of the family  $\mathcal{S}$ . Then  $\alpha \leq_{pr} \nu$  by primitive recursive function  $g$ , which means that *alpha* is punctually decidable, since  $\forall n, m \alpha(m) = \alpha(n) \Leftrightarrow \nu(g(m)) = \nu(g(n)) \Leftrightarrow g(m) = g(n)$ . Consequently ([28], Proposition 3.1(ii)), there is *spd*-numbering  $\mu \equiv_{pr} \alpha$ . By using the construction from the Theorem 4.1 of the same paper, we can construct the numbering  $\mu_0 <_{pr} \mu$ , which contradicts to choice of  $\alpha$ .

2) Let  $\nu$  be friedberg numbering of the infinite family  $\mathcal{S}$ . Consider  $\mu = \nu \circ f$ , where  $f$  is a primitive recursive bijective function such that  $f^{-1}$  is not primitive recursive (existence of such function is shown in [33]). It is clear, that  $\mu \leq_{pr} \nu$  and  $\mu$  friedberg:  $\mu(m) = \mu(n) \Leftrightarrow \nu(f(m)) = \nu(f(n)) \Leftrightarrow f(m) = f(n) \Leftrightarrow m = n$ . Wherein,  $\nu = \mu \circ f^{-1}$  and  $f^{-1}$  is not primitive recursive, which means that  $\nu \not\leq_{pr} \mu$ . Thus  $\mu <_{pr} \nu$ . Continuing the process, you can build an endless-waning chain of friedberg numberings, from where we get required.

#### Acknowledgments

The work was done during The Workshop at the Mathematical Center in Akademgorodok in 2022 (website of the workshop: <https://bmm2022.mca.nsu.ru/>). The authors would like to thank the organizers of the workshop, and are immensely thankful to the project mentors Nikolay Bazhenov and Iskander Kalimullin for their valuable guidance.

The work of A. Askarbekkyzy, B. Kalmurayev, D. Nurlanbek, F. Rakymzhankyzy was supported by the Ministry of Science and Higher Education of the Republic of Kazakhstan, grant AP19576325 “Algorithmic complexities of presentations for algebraic structures”.

*Author Contributions*

A. Askarbekkyzy proved Theorem 3. R. Bagaviev worked on Proposition 5. V. Isakov proved Proposition 1. B. Kalmurzayev set the direction of the research and participated in discussions of all the evidence and propositions in this article. D. Nurlanbek proved Proposition 3. F. Rakymzhankyzy proved Theorem 4. A. Slobozhanin proved Theorem 1. All authors participated in discussions of the definitions that were introduced in this article. All authors participated in the revision of the manuscript and approved the final submission.

All authors contributed equally to this work.

*Conflict of Interest*

The authors declare no conflict of interest.

References

- 1 Khutoretskii, A.B. (1971). On the cardinality of the upper semilattice of computable enumerations. *Algebra and Logic*, 10(5), 348–352. <https://doi.org/10.1007/bf02219842>
- 2 Selivanov, V.L. (1976). Two theorems on computable enumerations. *Algebra Logika*, 15(4), 297–306.
- 3 Rogers, H. (1958). Godel numberings of partial recursive functions. *The Journal of Symbolic Logic*, 23(3), 331–341. <https://doi.org/10.2307/2964292>
- 4 Friedberg, R.M. (1958). Three theorems on recursive enumeration. I. Decomposition. II. Maximal set. III. Enumeration without duplication. *The Journal of Symbolic Logic*, 23(3), 309–316. <https://doi.org/10.2307/2964290>
- 5 Badaev, S.A. (1975). On incomparable enumerations. *Siberian Mathematical Journal*, 15(4), 519–524. <https://doi.org/10.1007/bf00967428>
- 6 Badaev S.A. (1992). On minimal enumerations. *Siberian Advances in Mathematics*, 2(1), 1–30.
- 7 Goncharov, S.S., & Badaev, S.A. (1998). Families with one-element Rogers semilattice. *Algebra and Logic*, 37(1), 21–34. <https://doi.org/10.1007/bf02684082>
- 8 Badaev, S., & Goncharov, S. (1996). On computable minimal enumerations. In Y. Ershov, E. Khukhro, V. Levchuk & N. Podufalov (Ed.), *Algebra: Proceedings of the Third International Conference on Algebra held in Krasnoyarsk, August 23–28, 1993 (pp. 21–34)*. Berlin, Boston: De Gruyter. <https://doi.org/10.1515/9783110813418-005>
- 9 Kalimullin, I., Melnikov, A., & Ng, K.M. (2017). Algebraic structures computable without delay. *Theoretical Computer Science*, 674, 73–98. <https://doi.org/10.1016/j.tcs.2017.01.029>
- 10 Kalimullin, I.S., Melnikov, A.G., & Ng, K.M. (2017). The diversity of categoricity without delay. *Algebra and Logic*, 56(2), 171–177. <https://doi.org/10.1007/s10469-017-9437-6>
- 11 Melnikov, A.G., & Ng, K.M. (2019). The back-and-forth method and computability without delay. *Israel Journal of Mathematics*, 234(2), 959–1000. <https://doi.org/10.1007/s11856-019-1948-5>
- 12 Zubkov, M.V., Kalimullin, I.S., Melnikov, A.G., & Frolov, A.N. (2019). Punctual copies of algebraic structures. *Siberian Mathematical Journal*, 60(6), 993–1002. <https://doi.org/10.1134/s0037446619060077>
- 13 Downey, R., Harrison-Trainor, M., Kalimullin, I., Melnikov, A., & Turetsky, D. (2020). Graphs are not universal for online computability. *Journal of Computer and System Sciences*, 112, 1–12. <https://doi.org/10.1016/j.jcss.2020.02.004>

- 14 Melnikov, A., & Ng, K.M. (2020). A structure of punctual dimension two. *Proceedings of the American Mathematical Society*, 148(7), 3113–3128. <https://doi.org/10.1090/proc/15020>
- 15 Bazhenov, N., Kalimullin, I., Melnikov, A., & Ng, K.M. (2020). Online presentations of finitely generated structures. *Theoretical Computer Science*, 844, 195–216. <https://doi.org/10.1016/j.tcs.2020.08.021>
- 16 Kalimullin, I.S., & Melnikov, A.G. (2021). Punctual categoricity relative to a computable oracle. *Lobachevskii Journal of Mathematics*, 42(4), 735–742. <https://doi.org/10.1134/s1995080221040107>
- 17 Greenberg, N., Harrison-Trainor, M., Melnikov, A., & Turetsky, D. (2021). Non-density in punctual computability. *Annals of Pure and Applied Logic*, 172(9), 102985. <https://doi.org/10.1016/j.apal.2021.102985>
- 18 Kalimullin, I., Melnikov, A., & Montalban, A. (2021). Punctual definability on structures. *Annals of Pure and Applied Logic*, 172(8), 102987. <https://doi.org/10.1016/j.apal.2021.102987>
- 19 Kalimullin, I., Melnikov, A., & Zubkov, M. (2023). Punctual degrees and lattice embeddings. In *Lecture notes series, Institute For Mathematical Sciences* (pp. 315–334). [https://doi.org/10.1142/9789811278631\\_0012](https://doi.org/10.1142/9789811278631_0012)
- 20 Downey, R., Melnikov, A., & Ng, K.M. (2021). Foundations of Online Structure Theory II: The Operator Approach. *Logical Methods in Computer Science*, 17(3). [https://doi.org/10.46298/lmcs-17\(3:6\)2021](https://doi.org/10.46298/lmcs-17(3:6)2021)
- 21 Bazhenov, N.A., & Kalimullin, I.S. (2021). Punctual categoricity spectra of computably categorical structures. *Algebra and Logic*, 60(3), 223–228. <https://doi.org/10.1007/s10469-021-09645-9>
- 22 Kalimullin, I.S. (2020). Constructing punctually categorical semigroups. *Algebra and Logic*, 59(5), 408–411. <https://doi.org/10.1007/s10469-020-09614-8>
- 23 Bazhenov, N., Ng, K.M., Mauro, L.S., & Sorbi, A. (2022). Primitive recursive equivalence relations and their primitive recursive complexity. *Computability*, 11(3–4), 187–221. <https://doi.org/10.3233/com-210375>
- 24 Bazhenov, N., Mustafa, M., & Ospichev, S. (2022). Rogers semilattices of punctual numberings. *MSCS. Mathematical Structures in Computer Science*, 32(2), 164–188. <https://doi.org/10.1017/s0960129522000093>
- 25 Bazhenov, N. (2020). Definable subsets of Polynomial-Time algebraic structures. In *Lecture notes in computer science* (pp. 142–154). [https://doi.org/10.1007/978-3-030-50026-9\\_10](https://doi.org/10.1007/978-3-030-50026-9_10)
- 26 Bazhenov, N., Harrison-Trainor, M., Kalimullin, I., Melnikov, A., & Ng, K.M. (2019). Automatic and polynomial-time algebraic structures. *The Journal of Symbolic Logic*, 84(4), 1630–1669. <https://doi.org/10.1017/jsl.2019.26>
- 27 Bazhenov, N., Fiori-Carones, M., Liu, L., & Melnikov, A. (2024). Primitive recursive reverse mathematics. *Annals of Pure and Applied Logic*, 175(1), 103354. <https://doi.org/10.1016/j.apal.2023.103354>
- 28 Bazhenov, N., Mustafa, M., & Ospichev, S. (2022). Rogers semilattices of punctual numberings. *Mathematical Structures in Computer Science*, 1–25.
- 29 Ershov, Iu.L. (1977). *Teoriia numeratsii [Theory of Numbers]*. Moscow: Nauka [in Russian].
- 30 Soare, R.I. (1999). *Recursively enumerable sets and degrees: A Study of Computable Functions and Computably Generated Sets*. Springer Science & Business Media.
- 31 Bazhenov, N., Downey, R., Kalimullin, I., & Melnikov, A. (2019). Foundations of online structure theory. *Bulletin of Symbolic Logic*, 25(2), 141–181. <https://doi.org/10.1017/bsl.2019.20>
- 32 Kalmurzayev, B.S., Bazhenov, N.A., & Isakov, A.M. (2023). A note on the degree structure of primitive recursive  $m$ -reducibility. *Preprint, arXiv:2311.15253*.

- 33 Downey, R., Greenberg, N., Melnikov, A., Ng, K.M., & Turetsky, D. (2020). Punctual categoricity and universality. *The Journal of Symbolic Logic*, 85(4), 1427–1466. <https://doi.org/10.1017/jsl.2020.51>

*Author Information\**

**Aknur Askarbekkyzy** — PhD student, Lecturer, Kazakh-British Technical University, Almaty, Kazakhstan; e-mail: [ms.askarbekkyzy@gmail.com](mailto:ms.askarbekkyzy@gmail.com); <https://orcid.org/0000-0003-0075-4438>

**Ramil Bagaviev** — Master's student, Kazan Federal University, Kazan, Russia; e-mail: [ramilbagaviev@mail.ru](mailto:ramilbagaviev@mail.ru)

**Valeriy Isakov** — Bachelor's student, Novosibirsk State University, Novosibirsk, Russia; e-mail: [v.isakov@g.nsu.ru](mailto:v.isakov@g.nsu.ru)

**Birzhan Kalmurzayev** — Associate Professor, PhD, Kazakh-British Technical University, Almaty, Kazakhstan; e-mail: [birzhan.kalmurzayev@gmail.com](mailto:birzhan.kalmurzayev@gmail.com); <https://orcid.org/0000-0002-4386-5915>

**Dias Nurlanbek** (*corresponding author*) — PhD student, Tutor, Kazakh-British Technical University, Almaty, Kazakhstan; e-mail: [nurlanbek.dias21@gmail.com](mailto:nurlanbek.dias21@gmail.com); <https://orcid.org/0000-0002-1275-1413>

**Fariza Rakymzhankyzy** — PhD candidate, Senior-Lecturer, Kazakh-British Technical University, Almaty, Kazakhstan; e-mail: [fariza.rakymzhankyzy@gmail.com](mailto:fariza.rakymzhankyzy@gmail.com); <https://orcid.org/0000-0002-6517-5560>

**Artyom Slobozhanin** — Master's student, Novosibirsk State University, Novosibirsk, Russia; e-mail: [a.slobozhanin@g.nsu.ru](mailto:a.slobozhanin@g.nsu.ru)

---

\*The author's name is presented in the order: First, Middle and Last Names.