

MATHEMATICS

<https://doi.org/10.31489/2024M2/4-21>

Research article

On estimates of M -term approximations of the Sobolev class in the Lorentz space

G. Akishev¹, A.Kh. Myrzagalieva^{2,*}

¹Kazakhstan Branch of Lomonosov Moscow State University, Astana, Kazakhstan;

²Astana IT University, Astana, Kazakhstan

(E-mail: akishev_g@mail.ru, aigul.myrzagalieva@astanait.edu.kz)

In the paper spaces of periodic functions of several variables were considered, namely the Lorentz space $L_{2,\tau}(\mathbb{T}^m)$, the class of functions with bounded mixed fractional derivative $W_{2,\tau}^{\bar{\tau}}$, $1 \leq \tau < \infty$, and the order of the best M -term approximation of a function $f \in L_{p,\tau}(\mathbb{T}^m)$ by trigonometric polynomials was studied. The article consists of an introduction, a main part, and a conclusion. In the introduction, basic concepts, definitions and necessary statements for the proof of the main results were considered. One can be found information about previous results on the mentioned topic. In the main part, exact-order estimates are established for the best M -term approximations of functions of the Sobolev class $W_{2,\tau_1}^{\bar{\tau}}$ in the norm of the space $L_{p,\tau_2}(\mathbb{T}^m)$ for various relations between the parameters p, τ_1, τ_2 .

Keywords: Lorentz space, Sobolev class, mixed derivative, trigonometric polynomial, M -term approximation.

2020 Mathematics Subject Classification: 41A10, 41A25, 42A05.

Introduction

Let \mathbb{N} , \mathbb{Z} , \mathbb{R} be the sets of natural, integer, and real numbers, respectively, and $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$, \mathbb{R}^m is m -dimensional Euclidean space of points $\bar{x} = (x_1, \dots, x_m)$ with real coordinates; $\mathbb{T}^m = [0, 2\pi]^m$ and $\mathbb{I}^m = [0, 1]^m$ are m -dimensional cubes.

We denote by $L_{p,\tau}(\mathbb{T}^m)$ the Lorentz space of all real-valued Lebesgue measurable functions f that have 2π -period in each variable and for which the quantity

$$\|f\|_{p,\tau} = \left\{ \frac{\tau}{p} \int_0^1 (f^*(t))^{\tau} t^{\frac{\tau}{p}-1} dt \right\}^{\frac{1}{\tau}}, \quad 1 < p < \infty, \quad 1 \leq \tau < \infty$$

is finite, where $f^*(t)$ is a non-increasing rearrangement of the function $|f(2\pi\bar{x})|$, $\bar{x} \in \mathbb{I}^m$ (see [1]).

*Corresponding author. E-mail: aigul.myrzagalieva@astanait.edu.kz

This research was funded by the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan (Grant No. AP22683029).

Received: 24 August 2023; Accepted: 05 February 2024.

In case when $\tau = p$, the Lorentz space $L_{p,\tau}(\mathbb{T}^m)$ coincides with the Lebesgue space $L_p(\mathbb{T}^m)$ with the norm (see, for example, [2])

$$\|f\|_p = \left[\int_0^{2\pi} \dots \int_0^{2\pi} |f(x_1, \dots, x_m)|^p dx_1 \dots dx_m \right]^{\frac{1}{p}}, \quad 1 \leq p < \infty.$$

Let us begin by introducing some notation: $a_{\bar{n}}(f)$ are Fourier coefficients of the function $f \in L_1(\mathbb{T}^m)$ by the system $\{e^{i\langle \bar{n}, \bar{x} \rangle}\}_{\bar{n} \in \mathbb{Z}^m}$ and $\langle \bar{y}, \bar{x} \rangle = \sum_{j=1}^m y_j x_j$;

$$\delta_{\bar{s}}(f, \bar{x}) = \sum_{\bar{n} \in \rho(\bar{s})} a_{\bar{n}}(f) e^{i\langle \bar{n}, \bar{x} \rangle},$$

where

$$\rho(\bar{s}) = \{\bar{k} = (k_1, \dots, k_m) \in \mathbb{Z}^m : [2^{s_j-1}] \leq |k_j| < 2^{s_j}, j = 1, \dots, m\},$$

and $[a]$ is an integer part of a , $\bar{s} = (s_1, \dots, s_m)$, $s_j = 0, 1, 2, \dots$

For a given vector $\bar{r} = (r_1, \dots, r_m) > \bar{0} = (0, \dots, 0)$ we set $\bar{\gamma} = \frac{\bar{r}}{r_1}$ and

$$Q_n^{(\bar{\gamma})} = \cup_{\langle \bar{s}, \bar{\gamma} \rangle < n} \rho(\bar{s}),$$

$S_n^{(\bar{\gamma})}(f, \bar{x}) = \sum_{\bar{k} \in Q_n^{(\bar{\gamma})}} a_{\bar{k}}(f) e^{i\langle \bar{k}, \bar{x} \rangle}$ is a partial sum of the Fourier series of the function f (see [2]).

Let us consider an one-dimensional Bernoulli kernel (see, for example, [2])

$$F_r(x) = 1 + 2 \sum_{k=1}^{\infty} k^{-r} \cos(kx - r\pi/2), \quad r > 0.$$

Next, for the vector $\bar{r} = (r_1, \dots, r_m)$, $r_j > 0$, $j = 1, \dots, m$, we set

$$F_{\bar{r}}(\bar{x}) = \prod_{j=1}^m F_{r_j}(x_j).$$

Let us consider a Sobolev functional class

$$W_{p,\tau}^{\bar{r}} = \{f : f = \varphi \star F_{\bar{r}}, \|\varphi\|_{p,\tau} \leq 1\},$$

where $1 < p < \infty$, $1 \leq \tau < \infty$,

$$(\varphi \star F_{\bar{r}})(\bar{x}) = \frac{1}{(2\pi)^m} \int_{\mathbb{T}^m} \varphi(\bar{x} - \bar{u}) F_{\bar{r}}(\bar{u}) d\bar{u}.$$

In case when $\tau = p$, the class $W_{p,\tau}^{\bar{r}}$ has been considered in [3] and [4], so in this case, instead of $W_{p,p}^{\bar{r}}$ we write $W_p^{\bar{r}}$.

The value

$$e_M(f)_{p,\tau} = \inf_{\bar{k}^{(j)}, b_j} \left\| f - \sum_{j=1}^M b_j e^{i\langle \bar{k}^{(j)}, \bar{x} \rangle} \right\|_{p,\tau}$$

is called the best M -term trigonometric approximation of the function $f \in L_{p,\tau}(\mathbb{T}^m)$, $n \in \mathbb{N}$.

If $F \subset L_{p,\tau}(\mathbb{T}^m)$ is some functional class, then we set $e_M(F)_{p,\tau} = \sup_{f \in F} e_M(f)_{p,\tau}$. In case when $\tau = p$, instead of $e_M(F)_{p,\tau}$ we write $e_M(F)_p$.

The best M -term approximation of a function $f \in L_2[0, 1]$ by polynomials in an orthonormal system has been first determined by S.B. Stechkin [5] and he has established a criterion for the absolute convergence of the Fourier series in this system. The advantage of the M -term approximation with respect to the one-dimensional trigonometric system over the linear approximation by M -order trigonometric polynomials has been shown by R.S. Ismagilov [6].

Exact order estimates of the best M -term approximation of the Bernoulli kernel have been established by V.E. Maiorov [7] and Yu. Makovoz [8], E.S. Belinsky [9, 10]. In the one-dimensional case, the value $e_M(W_q^r)_p$ has been estimated by S. Belinsky [9]. At present, many important results on estimates of M -term approximations of functions from various Sobolev, Nikol'skii-Besov and Lizorkin-Triebel classes are known [11, 12]. In the multidimensional case, for $1 < q \leq p < 2$ and $r_1 > \frac{1}{2}(\frac{1}{q} - \frac{1}{p})$, order-exact estimates of the best M -term approximation of functions of W_q^r in the norm of $L_p(\mathbb{T}^m)$ have been obtained by V.N. Temlyakov [3, 4], and for $1 < q \leq p < 2$ and $r_1 \leq \frac{1}{2}(\frac{1}{q} - \frac{1}{p})$, E.S. Belinsky [10] has proved the following theorem:

Theorem. Let $1 < q \leq 2 < p < \infty$ and $r_1 = \dots = r_\nu < r_{\nu+1} \leq \dots r_m$. Then

$$e_M(W_q^r)_p \asymp M^{-\frac{p}{2}(r_1 + \frac{1}{p} - \frac{1}{q})} (\log M)^{(\nu-1)(p-1)(r_1 - p'(\frac{1}{q} - \frac{1}{p}))_+}$$

in case $\frac{1}{q} - \frac{1}{p} < r_1 < \frac{1}{q}$, where $q' = \frac{q}{q-1}$.

Note that a generalization of this theorem on the Lorentz space $L_{p,\tau}(\mathbb{T}^m)$ has been proved in [13–15].

Throughout the paper, $A_n \asymp B_n$ means that there are positive numbers C_1, C_2 independent of $n \in \mathbb{N}$ such that $C_1 A_n \leq B_n \leq C_2 A_n$ for $n \in \mathbb{N}$ and $\log M$, where $\log M$ is the logarithm with base 2 of the number $M > 1$.

By the constructive method, V.N. Temlyakov [16, 17] has established estimates for M -term approximations of functions of the class W_q^r in the space $L_p(\mathbb{T}^m)$ for $1 < q \leq 2 < p < \infty$ and $(\frac{1}{q} - \frac{1}{p})p' < r_1 < \frac{1}{q}$, $p' = \frac{p}{p-1}$ and has raised the question of finding constructive evaluation method for $\frac{1}{q} - \frac{1}{p} < r_1 \leq (\frac{1}{q} - \frac{1}{p})p'$. Further application of the constructive method is given in [18, 19].

In the first section, some auxiliary assertions are formulated that are necessary for proving main results. The main results of the article are formulated as a theorem and proved in the second section. In conclusion, we compare the proved Theorem 1 with previously known results.

1 Auxiliary statements

Theorem A. [20] Let $1 < q < \lambda < \infty$, $1 < \tau$, $\theta < \infty$. If a function $f \in L_{q,\tau}(\mathbb{T}^m)$, then

$$\|f\|_{q,\tau} \geq C \left(\sum_{\bar{s} \in \mathbb{Z}_+^m} \prod_{l=1}^m 2^{s_l(1/\lambda - 1/q)\tau} \|\delta_{\bar{s}}(f)\|_{\lambda,\theta}^\tau \right)^{1/\tau}.$$

Theorem B. [20] Let $1 < p < q < \infty$, $1 < \tau_1, \tau_2 < \infty$. If a function $f \in L_{p,\tau_1}(\mathbb{T}^m)$ satisfies the condition

$$\sum_{\bar{s} \in \mathbb{Z}_+^m} \prod_{j=1}^m 2^{s_j \tau_2(1/p - 1/q)} \|\delta_{\bar{s}}(f)\|_{p,\tau_1}^{\tau_2} < \infty,$$

then $f \in L_{q,\tau_2}(\mathbb{T}^m)$ and the inequality

$$\|f\|_{q,\tau_2} \leq C \left(\sum_{\bar{s} \in \mathbb{Z}_+^m} \prod_{j=1}^m 2^{s_j \tau_2(1/p - 1/q)} \|\delta_{\bar{s}}(f)\|_{p,\tau_1}^{\tau_2} \right)^{1/\tau_2}$$

holds.

For a function $f \in L_1(\mathbb{T}^m)$ we set

$$f_{l,\bar{r}}(\bar{x}) = \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} \delta_{\bar{s}}(f, \bar{x}), l \in \mathbb{Z}_+,$$

where $\bar{\gamma} = (\gamma_1, \dots, \gamma_m)$, $\gamma_1 = \dots = \gamma_\nu < \gamma_{\nu+1} \leq \dots \leq \gamma_m$, $\gamma_j = \frac{r_j}{r_1}$, $r_j > 0$, $j = 1, \dots, m$.

Let us consider the following class defined in [5, 6]

$$W_A^{a,b,\bar{r}} = \left\{ f \in L_1(\mathbb{T}^m) : \|f_{l,\bar{r}}\|_A \leq 2^{-la} l^{(\nu-1)b} \right\},$$

where

$$\|f_{l,\bar{r}}\|_A = \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} \sum_{\bar{n} \in \rho(\bar{s})} |a_{\bar{n}}(f)|.$$

The following lemma is a consequence of Lemma 6.1 in [16] (see also Lemma 2.1 in [17]), which we often use in proofs of main results.

Lemma 1. [15] Let $2 \leq p < \infty$ and $1 < \tau < \infty$, $a > 0$. Then for $f \in W_A^{a,b,\bar{r}}$ there are constructive approximation methods of the greedy algorithm type of $G_M(f)$ with the property:

$$\|f - G_M(f)\|_{p,\tau} \leq C(m) M^{-a-\frac{1}{2}} (\log M)^{(\nu-1)(a+b)}.$$

2 Main results

Theorem 1. Let $0 < r_1 = \dots = r_\nu < r_{\nu+1} \leq \dots \leq r_m$, $2 < p < \infty$, $1 < \max\{\tau_1, 2\} \leq \tau_2 < \infty$, $\tau_2' = \frac{\tau_2}{\tau_2-1}$.

a) If $\frac{1}{2} - \frac{1}{p} < r_1 < \left(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}\right)\tau_2'$, then

$$e_M(W_{2,\tau_1}^{\bar{r}})_{p,\tau_2} \leq CM^{-\frac{p}{2}(r_1+\frac{1}{p}-\frac{1}{2})} (\log_2 M)^{\frac{1}{2}-\frac{1}{\tau_1}}, M > 1.$$

b) If $\tau_2' \left(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}\right) < r_1 < \frac{1}{2}$, then

$$e_M(W_{2,\tau_1}^{\bar{r}})_{p,\tau_2} \leq CM^{-\frac{p}{2}(r_1+\frac{1}{p}-\frac{1}{2})} (\log_2 M)^{\frac{1}{2}-\frac{1}{\tau_1}} (\log_2 M)^{(\nu-1)\frac{p}{\tau_2'} \left(r_1 - \tau_2' \left(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}\right)\right)}.$$

Proof. Let us introduce some notation

$$Q_{n,\bar{\gamma}} = \cup_{\langle \bar{s}, \bar{\gamma} \rangle \leq n} \rho(\bar{s}), S_{Q_{n,\bar{\gamma}}}(f, \bar{x}) = \sum_{\langle \bar{s}, \bar{\gamma} \rangle \leq n} \delta_{\bar{s}}(f, \bar{x}).$$

For a natural number M , there exists a number $n \in \mathbb{N}$ such that $M \asymp 2^n n^{\nu-1}$.

Let $\nu \geq 2$. We set

$$n_1 = \frac{p}{2}n - p\left(\frac{1}{2} - \frac{1}{\tau_2}\right)(\nu-1)\log n,$$

$$n_2 = \frac{p}{2}n + \frac{p}{2}(\nu-1)\log n.$$

Also, let us introduce

$$S_l = \left(2^{lr_1\tau_1} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} 2^{\langle \bar{s}, \bar{1} \rangle \left(\frac{1}{2} - \frac{1}{q}\right)\tau_1} \|\delta_{\bar{s}}(\varphi)\|_2^{\tau_1} \right)^{1/\tau_1}$$

and

$$m_l = \left[2^{-l \frac{\tau_2'}{p}} S_l^{\tau_1} 2^{n \frac{\tau_2'}{2}} n^{(\nu-1) \frac{\tau_2'}{2}} \right] + 1,$$

where $\langle \bar{s}, \bar{1} \rangle = \sum_{j=1}^m s_j$, $p' = \frac{p}{p-1}$ and $[y]$ is an integer part of a number y .

By $G(l)$ is denoted the set of indices \bar{s} , $l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1$, with the largest $\|\delta_{\bar{s}}(\varphi)\|_2$, and $m_l = |G(l)|$ is the number of elements of $G(l)$.

Let us consider the functions

$$\begin{aligned} F_1(\bar{x}) &= \sum_{n \leq l < n_1} f_l(\bar{x}), \\ F_2(\bar{x}) &= \sum_{n_1 \leq l < n_2} \sum_{\bar{s} \notin G(l)} \delta_{\bar{s}}(f, \bar{x}), \\ F_3(\bar{x}) &= \sum_{n_1 \leq l < n_2} \sum_{\bar{s} \in G(l)} \delta_{\bar{s}}(f, \bar{x}). \end{aligned}$$

Let us estimate $\|F_1\|_A$. Applying Hölder's inequality for the sum and Parseval's equality, we have

$$\begin{aligned} \|F_1\|_A &= \sum_{l=n}^{n_1-1} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} \sum_{\bar{k} \in \rho(\bar{s})} |a_{\bar{k}}(f)| \leq 2^{-\frac{m}{2}} \sum_{l=n}^{n_1-1} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} 2^{\langle \bar{s}, \bar{1} \rangle \frac{1}{2}} \|\delta_{\bar{s}}(f)\|_2 = \\ &= 2^{-\frac{m}{2}} \sum_{l=n}^{n_1-1} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} 2^{\langle \bar{s}, \bar{1} \rangle (\frac{1}{2} - \frac{1}{q})} \|\delta_{\bar{s}}(f)\|_2 2^{\langle \bar{s}, \bar{1} \rangle \frac{1}{q}}. \end{aligned} \tag{1}$$

It is known that the Fourier coefficients of the convolution $f = \varphi \star F_{\bar{r}}$ are equal to $a_{\bar{k}}(\varphi) a_{\bar{k}}(F_{\bar{r}})$, $\bar{k} \in \mathbb{Z}^m$. Therefore, using Parseval's equality, it is easy to verify that

$$\|\delta_{\bar{s}}(f)\|_2 \ll 2^{-\langle \bar{s}, \bar{r} \rangle} \|\delta_{\bar{s}}(\varphi)\|_2, \quad \bar{s} \in \mathbb{Z}_+^m. \tag{2}$$

Hence, from (1) and (2) we get

$$\|F_1\|_A \leq 2^{-\frac{m}{2}} \sum_{l=n}^{n_1-1} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} 2^{\langle \bar{s}, \bar{1} \rangle \frac{1}{2}} \|\delta_{\bar{s}}(f)\|_2 \leq C \sum_{l=n}^{n_1-1} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} 2^{\langle \bar{s}, \bar{1} \rangle \frac{1}{2}} 2^{-\langle \bar{s}, \bar{r} \rangle} \|\delta_{\bar{s}}(\varphi)\|_2. \tag{3}$$

If $2 < \tau_1 < \infty$, then according to the inequality of different metrics for trigonometric polynomials in the Lorentz space [20] we have

$$\|\delta_{\bar{s}}(\varphi)\|_2 \leq C \left(\sum_{j=1}^m (s_j + 1) \right)^{\frac{1}{2} - \frac{1}{\tau_1}} \|\delta_{\bar{s}}(\varphi)\|_{2, \tau_1}.$$

From Lemma 1.6 [21] for $p = 2$ and $2 < \tau_1 < \infty$ we get

$$\left(\sum_{\bar{s} \in \mathbb{Z}_+^m} \left(\sum_{j=1}^m (s_j + 1) \right)^{\left(\frac{1}{\tau_1} - \frac{1}{2}\right)\tau_1} \|\delta_{\bar{s}}(\varphi)\|_2^{\tau_1} \right)^{\frac{1}{\tau_1}} \leq C \left(\sum_{\bar{s} \in \mathbb{Z}_+^m} \|\delta_{\bar{s}}(\varphi)\|_{2, \tau_1}^{\tau_1} \right)^{\frac{1}{\tau_1}} \leq C \|\varphi\|_{2, \tau_1}. \tag{4}$$

By virtue of inequality (4) and Hölder's inequality, we obtain

$$\begin{aligned} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} 2^{\langle \bar{s}, \bar{1} \rangle \frac{1}{2}} \|\delta_{\bar{s}}(f)\|_2 &\leq \left(\sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} \left(\sum_{j=1}^m (s_j + 1) \right)^{\left(\frac{1}{\tau_1} - \frac{1}{2}\right)\tau_1} \|\delta_{\bar{s}}(\varphi)\|_2^{\tau_1} \right)^{\frac{1}{\tau_1}} \times \\ &\times \left(\sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} 2^{-\langle \bar{s}, \bar{\gamma} \rangle (r_1 - \frac{1}{2})\tau_1'} \left(\sum_{j=1}^m (s_j + 1) \right)^{\left(\frac{1}{2} - \frac{1}{\tau_1}\right)\tau_1'} \right)^{\frac{1}{\tau_1}} \leq \\ &\leq C \|\varphi\|_{2, \tau_1} \left(\sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} 2^{-\langle \bar{s}, \bar{\gamma} \rangle (r_1 - \frac{1}{2})\tau_1'} \left(\sum_{j=1}^m (s_j + 1) \right)^{\left(\frac{1}{2} - \frac{1}{\tau_1}\right)\tau_1'} \right)^{\frac{1}{\tau_1}} \leq \\ &\leq C 2^{-l(r_1 - \frac{1}{2})} l^{(\nu-1)\frac{1}{\tau_1}} l^{\frac{1}{2} - \frac{1}{\tau_1}} \|\varphi\|_{2, \tau_1}, \end{aligned} \tag{5}$$

where $\tau_1' = \frac{\tau_1}{\tau_1 - 1}$, $1 < \tau_1 < \infty$.

(3) and (5) imply that

$$\|F_1\|_A \leq C \sum_{l=n}^{n_1-1} 2^{\frac{l}{2}} l^{(\nu-1)\frac{1}{\tau_1}} l^{\frac{1}{2} - \frac{1}{\tau_1}} 2^{-lr_1} \leq C 2^{-n_1(r_1 - \frac{1}{2})} n_1^{(\nu-1)\frac{1}{\tau_1}} n_1^{\frac{1}{2} - \frac{1}{\tau_1}} \tag{6}$$

for a function $f \in W_{2, \tau_1}^{\bar{r}}$ when $r_1 < \frac{1}{2}$ and $2 < \tau_1 < \infty$.

By Lemma 1 for the function F_1 using a constructive method, one can find an M -term trigonometric polynomial $G_M(F_1)$ such that

$$\|F_1 - G_M(F_1)\|_{p, \tau_2} \leq CM^{-\frac{1}{2}} 2^{-n_1(r_1 - \frac{1}{2})} n_1^{(\nu-1)\frac{1}{\tau_1}} n_1^{\frac{1}{2} - \frac{1}{\tau_1}}. \tag{7}$$

Therefore, according to inequality (6) and (7) and taking into account the definition of the number n_1 and the relation $M \asymp 2^n n^{\nu-1}$, we obtain

$$\|F_1 - G_M(F_1)\|_{p, \tau_2} \leq CM^{-\frac{p}{2}(r_1 + \frac{1}{p} - \frac{1}{2})} (\log M)^{\frac{1}{2} - \frac{1}{\tau_1}} \tag{8}$$

in case when $q = 2 < p < \infty$, $2 < \tau_1 < \infty$, $1 < \tau_2 < \infty$, $r_1 < \frac{1}{2}$.

Let us estimate $\|F_3\|_A$. Applying Hölder's inequality for the sum and Parseval's equality, we obtain

$$\begin{aligned} \|F_3\|_A &= \sum_{l=n_1}^{n_2-1} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1, \bar{s} \in G(l)} \sum_{\bar{k} \in \rho(\bar{s})} |a_{\bar{k}}(f)| \leq 2^{-\frac{m}{2}} \sum_{l=n_1}^{n_2-1} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1, \bar{s} \in G(l)} 2^{\langle \bar{s}, \bar{1} \rangle \frac{1}{2}} \|\delta_{\bar{s}}(f)\|_2 \leq \\ &\leq C \sum_{l=n_1}^{n_2-1} 2^{\frac{l}{2}} (l+1)^{\frac{1}{2} - \frac{1}{\tau_1}} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1, \bar{s} \in G(l)} \left(\sum_{j=1}^m (s_j + 1) \right)^{\left(\frac{1}{\tau_1} - \frac{1}{2}\right)\tau_1} \|\delta_{\bar{s}}(f)\|_2. \end{aligned} \tag{9}$$

Now, to the inner sum on the right side of inequality (9), applying Hölder's inequality for $\frac{1}{\tau_1} + \frac{1}{\tau_1} = 1$, $1 < \tau_1 < \infty$, we have

$$\|F_3\|_A \leq C \sum_{l=n_1}^{n_2-1} 2^{\frac{l}{2}} (l+1)^{\frac{1}{2} - \frac{1}{\tau_1}} \left(\sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1, \bar{s} \in G(l)} \left(\sum_{j=1}^m (s_j + 1) \right)^{\left(\frac{1}{\tau_1} - \frac{1}{2}\right)\tau_1} \|\delta_{\bar{s}}(f)\|_2^{\tau_1} \right)^{\frac{1}{\tau_1}} |G(l)|^{\frac{1}{\tau_1}}.$$

Then, using (2) we get

$$\begin{aligned} \|F_3\|_A &\leq C \sum_{l=n_1}^{n_2-1} 2^{\frac{l}{2}}(l+1)^{\frac{1}{2}-\frac{1}{\tau_1}} \times \\ &\times \left(\sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1, \bar{s} \in G(l)} \left(\sum_{j=1}^m (s_j+1) \right)^{\left(\frac{1}{\tau_1}-\frac{1}{2}\right)\tau_1} 2^{-\langle \bar{s}, \bar{r} \rangle} \|\delta_{\bar{s}}(\varphi)\|_2^{\tau_1} \right)^{\frac{1}{\tau_1}} |G(l)|^{\frac{1}{\tau_1}} \leq \\ &\leq C \sum_{l=n_1}^{n_2-1} 2^{l\left(\frac{1}{2}-r_1\right)}(l+1)^{\frac{1}{2}-\frac{1}{\tau_1}} \times \\ &\times \left(\sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1, \bar{s} \in G(l)} \left(\sum_{j=1}^m (s_j+1) \right)^{\left(\frac{1}{\tau_1}-\frac{1}{2}\right)\tau_1} \|\delta_{\bar{s}}(\varphi)\|_2^{\tau_1} \right)^{\frac{1}{\tau_1}} |G(l)|^{\frac{1}{\tau_1}}. \end{aligned} \tag{10}$$

We set

$$\tilde{S}_l = \left(\sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} \left(\sum_{j=1}^m (s_j+1) \right)^{\left(\frac{1}{\tau_1}-\frac{1}{2}\right)\tau_1} \|\delta_{\bar{s}}(f)\|_2^{\tau_1} \right)^{1/\tau_1}$$

and

$$m_l := |G(l)| := \left[2^{-l\frac{\tau_2'}{p}} \tilde{S}_l^{\tau_1} 2^{n\frac{\tau_2'}{2}} n^{(\nu-1)\frac{\tau_2'}{2}} \right] + 1.$$

Then (10) implies that

$$\begin{aligned} \|F_3\|_A &\leq C \sum_{l=n_1}^{n_2-1} 2^{-l\left(r_1-\frac{1}{2}\right)}(l+1)^{\frac{1}{2}-\frac{1}{\tau_1}} \tilde{S}_l m_l^{\frac{1}{\tau_1}} \leq \\ &\leq C \sum_{l=n_1}^{n_2-1} 2^{-l\left(r_1-\frac{1}{2}\right)}(l+1)^{\frac{1}{2}-\frac{1}{\tau_1}} \tilde{S}_l \left\{ 2^{-l\frac{\tau_2'}{p}} \tilde{S}_l^{\tau_1} 2^{n\frac{\tau_2'}{2}} n^{(\nu-1)\frac{\tau_2'}{2}} + 1 \right\}^{\frac{1}{\tau_1}} \leq \\ &\leq C \left\{ \left(2^n n^{\nu-1} \right)^{\frac{\tau_2'}{2\tau_1}} \sum_{l=n_1}^{n_2-1} 2^{-l\left(r_1-\frac{1}{2}+\frac{\tau_2'}{p\tau_1}\right)}(l+1)^{\frac{1}{2}-\frac{1}{\tau_1}} \tilde{S}_l^{1+\frac{\tau_1}{\tau_1}} + \sum_{l=n_1}^{n_2-1} 2^{-l\left(r_1-\frac{1}{2}\right)}(l+1)^{\frac{1}{2}-\frac{1}{\tau_1}} \tilde{S}_l \right\}. \end{aligned} \tag{11}$$

Since $\tilde{S}_l^{1+\frac{\tau_1}{\tau_1}} = \tilde{S}_l^{\tau_1}$ and $-\frac{1}{2} + \frac{\tau_2'}{p\tau_1} = \tau_2' \left(-\frac{1}{2} + \frac{1}{p} - \frac{1}{p\tau_1} + \frac{1}{2\tau_2} \right)$, then by (4) we have

$$\begin{aligned} \sum_{l=n_1}^{n_2-1} 2^{-l\left(r_1-\frac{1}{2}+\frac{\tau_2'}{p\tau_1}\right)}(l+1)^{\frac{1}{2}-\frac{1}{\tau_1}} \tilde{S}_l^{1+\frac{\tau_1}{\tau_1}} &= \sum_{l=n_1}^{n_2-1} 2^{-l\left(r_1-\tau_2'\left(\frac{1}{2}-\frac{1}{p}+\frac{1}{p\tau_1}-\frac{1}{2\tau_2}\right)\right)}(l+1)^{\frac{1}{2}-\frac{1}{\tau_1}} \tilde{S}_l^{\tau_1} \leq \\ &\leq C \sum_{l=n_1}^{n_2-1} 2^{-l\left(r_1-\tau_2'\left(\frac{1}{2}-\frac{1}{p}+\frac{1}{p\tau_1}-\frac{1}{2\tau_2}\right)\right)}(l+1)^{\frac{1}{2}-\frac{1}{\tau_1}} \|\varphi\|_{2,\tau_1}^{\tau_1} \leq \\ &\leq C \sum_{l=n_1}^{n_2-1} 2^{-l\left(r_1-\tau_2'\left(\frac{1}{2}-\frac{1}{p}+\frac{1}{p\tau_1}-\frac{1}{2\tau_2}\right)\right)} l^{\frac{1}{2}-\frac{1}{\tau_1}} \end{aligned}$$

for a function $f \in W_{2,\tau_1}^{\bar{r}}$ and $2 < \tau_1 < \infty$. Since $r_1 - \tau_2' \left(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2} \right) < 0$, then, taking into

account the definition of the number n_2 , from here we obtain

$$\begin{aligned} \sum_{l=n_1}^{n_2-1} 2^{-l(r_1-\frac{1}{2}+\frac{\tau_2'}{p\tau_1})} (l+1)^{\frac{1}{2}-\frac{1}{\tau_1}} \tilde{S}_l^{1+\frac{\tau_1}{\tau_1}} &\leq C 2^{-n_2(r_1-\tau_2'(\frac{1}{2}-\frac{1}{p}+\frac{1}{p\tau_1}-\frac{1}{2\tau_2}))} n_2^{\frac{1}{2}-\frac{1}{\tau_1}} \leq \\ &\leq C 2^{-n\frac{p}{2}(r_1-\tau_2'(\frac{1}{2}-\frac{1}{p}+\frac{1}{p\tau_1}-\frac{1}{2\tau_2}))} n^{-(\nu-1)\frac{p}{2}(r_1-\tau_2'(\frac{1}{2}-\frac{1}{p}+\frac{1}{p\tau_1}-\frac{1}{2\tau_2}))} n^{\frac{1}{2}-\frac{1}{\tau_1}} \end{aligned} \quad (12)$$

for a function $f \in W_{2,\tau_1}^{\bar{r}}$, $2 < \tau_1 < \infty$.

Next, due to inequality (4), taking into account that a function $f \in W_{2,\tau_1}^{\bar{r}}$ and $r_1 - \frac{1}{2} < 0$, we have

$$\begin{aligned} \sum_{l=n_1}^{n_2-1} 2^{-l(r_1-\frac{1}{2})} (l+1)^{\frac{1}{2}-\frac{1}{\tau_1}} \tilde{S}_l &\leq C \sum_{l=n_1}^{n_2-1} 2^{-l(r_1-\frac{1}{2})} (l+1)^{\frac{1}{2}-\frac{1}{\tau_1}} \|\varphi\|_{2,\tau_1} \leq \\ &\leq C \sum_{l=n_1}^{n_2-1} 2^{-l(r_1-\frac{1}{2})} (l+1)^{\frac{1}{2}-\frac{1}{\tau_1}} \leq C 2^{-n_2(r_1-\frac{1}{2})} (n_2+1)^{\frac{1}{2}-\frac{1}{\tau_1}} \leq \\ &\leq C 2^{-n\frac{p}{2}(r_1-\frac{1}{2})} n^{-(\nu-1)\frac{p}{2}(r_1-\frac{1}{2})} n^{\frac{1}{2}-\frac{1}{\tau_1}}. \end{aligned} \quad (13)$$

Now it follows from inequalities (11)–(13) that

$$\begin{aligned} \|F_3\|_A &\leq C \left\{ \left(2^n n^{\nu-1} \right)^{\frac{\tau_2'}{2\tau_1}} 2^{-n\frac{p}{2}(r_1-\tau_2'(\frac{1}{2}-\frac{1}{p}+\frac{1}{p\tau_1}-\frac{1}{2\tau_2}))} n^{-(\nu-1)\frac{p}{2}(r_1-\tau_2'(\frac{1}{2}-\frac{1}{p}+\frac{1}{p\tau_1}-\frac{1}{2\tau_2}))} n^{\frac{1}{2}-\frac{1}{\tau_1}} + \right. \\ &\quad \left. + (2^n n^{\nu-1})^{-\frac{p}{2}(r_1-\frac{1}{2})} n^{\frac{1}{2}-\frac{1}{\tau_1}} \right\} \end{aligned}$$

for a function $f \in W_{2,\tau_1}^{\bar{r}}$, $2 < \tau_1 < \infty$, $1 < \tau_2 < \infty$, $r_1 - \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}) < 0$.

Since $\frac{p}{2}(r_1 - \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2})) - \frac{\tau_2'}{2\tau_1} = \frac{p}{2}(r_1 - \frac{1}{2})$, then it follows that

$$\|F_3\|_A \leq C (2^n n^{\nu-1})^{-\frac{p}{2}(r_1-\frac{1}{2})} n^{\frac{1}{2}-\frac{1}{\tau_1}}. \quad (14)$$

Since $2 < p < \infty$, then by Lemma 1 for the function F_3 , by a constructive method, there is an M -term trigonometric polynomial $G_M(F_3)$ such that

$$\|F_3 - G_M(F_3)\|_{p,\tau_2} \leq CM^{-\frac{1}{2}} (2^n n^{\nu-1})^{-\frac{p}{2}(r_1-\frac{1}{2})} n^{\frac{1}{2}-\frac{1}{\tau_1}}.$$

Hence, in accordance with (14), we have

$$\|F_3 - G_M(F_3)\|_{p,\tau_2} \leq CM^{-\frac{p}{2}(r_1+\frac{1}{p}-\frac{1}{2})} (\log M)^{\frac{1}{2}-\frac{1}{\tau_1}} \quad (15)$$

for a function $f \in W_{2,\tau_1}^{\bar{r}}$ for $2 < p < \infty$, $2 < \tau_1 < \infty$, $1 < \tau_2 < \infty$ and $r_1 < \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2})$.

Let us estimate $\|F_2\|_{p,\tau_2}$. So,

$$\|F_2\|_{p,\tau_2} \leq C \left(\sum_{l=n_1}^{n_2-1} \sum_{l < \langle \bar{s}, \bar{\gamma} \rangle < l+1, \bar{s} \notin G(l)} 2^{\langle \bar{s}, \bar{1} \rangle (\frac{1}{2}-\frac{1}{p})\tau_2} \|\delta_{\bar{s}}(f)\|_2^{T_2-\tau_1} \|\delta_{\bar{s}}(f)\|_2^{\tau_1} \right)^{1/\tau_2}.$$

Taking into account that

$$\|\delta_{\bar{s}}(f)\|_2 \leq m_l^{-\frac{1}{\tau_1}} 2^{-lr_1} l^{\frac{1}{2}-\frac{1}{\tau_1}} \tilde{S}_l$$

for $\bar{s} \notin G(l)$ and substituting the values of the numbers m_l for $\tau_2 - \tau_1 \geq 0$, we have

$$\begin{aligned} \|F_2\|_{p,\tau_2} &\leq C \left(\sum_{l=n_1}^{n_2-1} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1, \bar{s} \notin G(l)} 2^{\langle \bar{s}, \bar{1} \rangle (\frac{1}{2} - \frac{1}{p}) \tau_2} \|\delta_{\bar{s}}(f)\|_2^{\tau_1} \left(m_l^{-\frac{1}{\tau_1}} 2^{-lr_1} l^{\frac{1}{2} - \frac{1}{\tau_1}} \tilde{S}_l \right)^{\tau_2 - \tau_1} \right)^{1/\tau_2} \leq \\ &\leq C \left(\sum_{l=n_1}^{n_2-1} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1, \bar{s} \notin G(l)} 2^{l(\frac{1}{2} - \frac{1}{p}) \tau_2} 2^{-lr_1 \tau_1} \|\delta_{\bar{s}}(\varphi)\|_2^{\tau_1} \left(m_l^{-\frac{1}{\tau_1}} 2^{-lr_1} l^{\frac{1}{2} - \frac{1}{\tau_1}} \tilde{S}_l \right)^{\tau_2 - \tau_1} \right)^{1/\tau_2} \leq \\ &\leq C \left(\sum_{l=n_1}^{n_2-1} \left(\left(2^{-l \frac{\tau_2'}{p}} \tilde{S}_l^{\tau_1} 2^{n \frac{\tau_2'}{2}} n^{(\nu-1) \frac{\tau_2'}{2}} \right)^{-\frac{1}{\tau_1}} 2^{-lr_1} \tilde{S}_l^{\frac{1}{2} - \frac{1}{\tau_1}} \right)^{\tau_2 - \tau_1} \times \right. \\ &\quad \times 2^{l(\frac{1}{2} - \frac{1}{p}) \tau_2} 2^{-lr_1 \tau_1} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1, \bar{s} \notin G(l)} \|\delta_{\bar{s}}(\varphi)\|_2^{\tau_1} \left. \right)^{1/\tau_2} = \\ &= C(2^n n^{\nu-1})^{-\frac{\tau_2'}{2} \frac{\tau_2 - \tau_1}{\tau_1 \tau_2}} \left(\sum_{l=n_1}^{n_2-1} 2^{-l(r_1 - \frac{\tau_2'}{p \tau_1})(\tau_2 - \tau_1)} l^{(\frac{1}{2} - \frac{1}{\tau_1})(\tau_2 - \tau_1)} l^{-(\frac{1}{\tau_1} - \frac{1}{2}) \tau_1} \tilde{S}_l^{\tau_1} \right)^{1/\tau_2}. \end{aligned} \tag{16}$$

Using inequality (4), it is easy to verify that

$$\begin{aligned} \tilde{S}_l &= \left(\sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} \left(\sum_{j=1}^m (s_j + 1) \right)^{\left(\frac{1}{\tau_1} - \frac{1}{2}\right) \tau_1} \|\delta_{\bar{s}}(\varphi)\|_2^{\tau_1} \right)^{1/\tau_1} \leq \\ &\leq C \left\| \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} \delta_{\bar{s}}(\varphi) \right\|_{2,\tau_1} \leq C \|\varphi\|_{2,\tau_1} \end{aligned} \tag{17}$$

for a function $f \in W_{2,\tau_1}^{\bar{r}}$, $2 < \tau_1 \leq \tau_2 < \infty$.

Now it follows from inequalities (16) and (17) that

$$\begin{aligned} \|F_2\|_{p,\tau_2} &\leq C(2^n n^{\nu-1})^{-\frac{\tau_2'}{2} \frac{\tau_2 - \tau_1}{\tau_1 \tau_2}} \times \\ &\times \left(\sum_{l=n_1}^{n_2-1} 2^{-l(r_1 - \frac{\tau_2'}{p \tau_1})(\tau_2 - \tau_1)} l^{(\frac{1}{2} - \frac{1}{\tau_1})(\tau_2 - \tau_1)} 2^{l(\frac{1}{2} - \frac{1}{p}) \tau_2} l^{(\frac{1}{2} - \frac{1}{\tau_1}) \tau_1} 2^{-lr_1 \tau_1} \right)^{1/\tau_2} = \\ &= C(2^n n^{\nu-1})^{-\frac{\tau_2'}{2} \frac{\tau_2 - \tau_1}{\tau_1 \tau_2}} \left(\sum_{l=n_1}^{n_2-1} 2^{-l\tau_2(r_1 - \frac{\tau_2'}{p \tau_1 \tau_2}(\tau_2 - \tau_1) - (\frac{1}{2} - \frac{1}{p}))} l^{(\frac{1}{2} - \frac{1}{\tau_1}) \tau_2} \right)^{1/\tau_2}. \end{aligned}$$

Since

$$r_1 - \frac{\tau_2'}{p \tau_1 \tau_2} (\tau_2 - \tau_1) - \left(\frac{1}{2} - \frac{1}{p}\right) = r_1 - \tau_2' \left(\frac{1}{2} - \frac{1}{p} + \frac{1}{p \tau_1} - \frac{1}{2\tau_2}\right),$$

then taking into account the definition of the number n_2 , from here we get

$$\begin{aligned} \|F_2\|_{p,\tau_2} &\leq C(2^n n^{\nu-1})^{-\frac{\tau_2'}{2} \frac{\tau_2 - \tau_1}{\tau_1 \tau_2}} 2^{-n_2(r_1 - \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p \tau_1} - \frac{1}{2\tau_2}))} n_2^{\frac{1}{2} - \frac{1}{\tau_1}} \leq \\ &\leq C 2^{-n \frac{p}{2}(r_1 - \frac{1}{p} - \frac{1}{2})} n^{\frac{1}{2} - \frac{1}{\tau_1}} \end{aligned} \tag{18}$$

for function $f \in W_{2,\tau_1}^{\bar{r}}$ when $2 < p < \infty$, $2 < \tau_1 \leq \tau_2 < \infty$, $r_1 < \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p \tau_1} - \frac{1}{2\tau_2})$.

Now it follows from inequalities (8), (15), and (18) that

$$\begin{aligned} & \|f - (S_{Q_{n,\bar{\gamma}}}(f) + G_M(F_1) + G_M(F_3))\|_{p,\tau_2} \leq \\ & \leq \|F_1 - G_M(F_1)\|_{p,\tau_2} + \|F_3 - G_M(F_3)\|_{p,\tau_2} + \|F_2\|_{p,\tau_2} + \\ & + \left\| \sum_{\langle \bar{s}, \bar{\gamma} \rangle \geq n_2} \delta_{\bar{s}}(f) \right\|_{p,\tau_2} \leq CM^{-\frac{p}{2}(r_1 + \frac{1}{p} - \frac{1}{2})} (\log M)^{\frac{1}{2} - \frac{1}{\tau_1}} + \left\| \sum_{\langle \bar{s}, \bar{\gamma} \rangle \geq n_2} \delta_{\bar{s}}(f) \right\|_{p,\tau_2} \end{aligned}$$

for a function $f \in W_{2,\tau_1}^{\bar{r}}$ when $2 < p < \infty$, $2 < \tau_1 \leq \tau_2 < \infty$, $\frac{1}{2} - \frac{1}{p} < r_1 < \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2})$.

Further, taking into account that $2 < \tau_1 < \tau_2 < \infty$ and $r_1 + \frac{1}{p} - \frac{1}{2} > 0$, and successively applying Theorem B, Jensen's inequality, Theorem A, then Lemma 1.3 [21] and Theorem 1.1 [21], we obtain

$$\begin{aligned} & \left\| \sum_{\langle \bar{s}, \bar{\gamma} \rangle \geq n_2} \delta_{\bar{s}}(f) \right\|_{p,\tau_2} = \left\| \sum_{l=n_2}^{\infty} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} \delta_{\bar{s}}(f) \right\|_{p,\tau_2} \leq \\ & \leq C \left(\sum_{l=n_2}^{\infty} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} \prod_{j=1}^m 2^{s_j(\frac{1}{2} - \frac{1}{p})\tau_2} \|\delta_{\bar{s}}(f)\|_{2,\tau_1}^{\tau_2} \right)^{\frac{1}{\tau_2}} \leq \\ & \leq C \left(\sum_{l=n_2}^{\infty} 2^{l(\frac{1}{2} - \frac{1}{p})\tau_2} \left[\sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} \|\delta_{\bar{s}}(f)\|_{2,\tau_1}^{\tau_1} \right]^{\frac{\tau_2}{\tau_1}} \right)^{\frac{1}{\tau_2}} \leq \\ & \leq C \left(\sum_{l=n_2}^{\infty} 2^{l(\frac{1}{2} - \frac{1}{p})\tau_2} \left\| \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} \delta_{\bar{s}}(f) \right\|_{2,\tau_1}^{\tau_2} \right)^{\frac{1}{\tau_2}} \leq C \left(\sum_{l=n_2}^{\infty} 2^{-l(r_1 + \frac{1}{p} - \frac{1}{2})p} \right)^{\frac{1}{p}} \leq \\ & \leq C 2^{-n_2(r_1 + \frac{1}{p} - \frac{1}{2})} \leq CM^{-\frac{p}{2}(r_1 + \frac{1}{p} - \frac{1}{2})}, \end{aligned}$$

that leads to

$$e_M(f)_{p,\tau_2} \leq \|f - (S_{Q_{n,\bar{\gamma}}}(f) + G_M(F_1) + G_M(F_3))\|_{p,\tau_2} \leq CM^{-\frac{p}{2}(r_1 + \frac{1}{p} - \frac{1}{2})} (\log M)^{\frac{1}{2} - \frac{1}{\tau_1}}$$

For a function $f \in W_{2,\tau_1}^{\bar{r}}$ when $2 < p < \infty$, $2 < \tau_1 \leq \tau_2 < \infty$, $\frac{1}{2} - \frac{1}{p} < r_1 < \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2})$.

Assume that $\tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}) < r_1 < \frac{1}{2}$. Then, taking into account the definition of the number n_1 , we get

$$\begin{aligned} & \sum_{l=n_1}^{n_2-1} 2^{-l(r_1 - \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}))} (l+1)^{\frac{1}{2} - \frac{1}{\tau_1}} \leq C 2^{-n_1(r_1 - \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}))} n_1^{\frac{1}{2} - \frac{1}{\tau_1}} \leq \\ & \leq C 2^{-\frac{p}{2}n(r_1 - \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}))} n^{p(\frac{1}{2} - \frac{1}{\tau_2})(\nu-1)} (r_1 - \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2})) n^{\frac{1}{2} - \frac{1}{\tau_1}} \end{aligned} \tag{19}$$

for a function $f \in W_{2,\tau_1}^{\bar{r}}$ when $\tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}) < r_1 < \frac{1}{2}$.

(11), (13) and (19) imply that

$$\|F_3\|_A \leq (2^n n^{\nu-1})^{-\frac{p}{2}(r_1 - \frac{1}{2})} n^{\frac{p}{\tau_2}(\nu-1)} (r_1 - \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2})) n^{\frac{1}{2} - \frac{1}{\tau_1}}$$

for a function $f \in W_{2,\tau_1}^{\bar{r}}$ when $\tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}) < r_1 < \frac{1}{2}$.

Hence, by Lemma 1 we obtain

$$\|F_3 - G_M(F_3)\|_{p,\tau_2} \leq CM^{-\frac{p}{2}(r_1 + \frac{1}{p} - \frac{1}{2})} (\log M)^{\frac{p}{\tau_2}(\nu-1)} (r_1 - \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2})) (\log M)^{\frac{1}{2} - \frac{1}{\tau_1}} \tag{20}$$

for a function $f \in W_{2,\tau_1}^{\bar{r}}$ when $\tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}) < r_1 < \frac{1}{2}$.

Let us estimate $\|F_2\|_{p,\tau_2}$ in case when $\tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}) < r_1 < \frac{1}{2}$. (16) and (17) imply that

$$\begin{aligned} \|F_2\|_{p,\tau_2} &\leq C(2^n n^{\nu-1})^{-\frac{\tau_2'}{2} \frac{\tau_2 - \tau_1}{\tau_1 \tau_2}} \left(\sum_{l=n_1}^{n_2-1} 2^{-l\tau_2(r_1 - \frac{\tau_2'}{p\tau_1\tau_2}(\tau_2 - \tau_1) - (\frac{1}{2} - \frac{1}{p}))} l^{(\frac{1}{2} - \frac{1}{\tau_1})\tau_2} \right)^{1/\tau_2} \leq \\ &\leq CM^{-\frac{p}{2}(r_1 + \frac{1}{p} - \frac{1}{2})} (\log M)^{\frac{p}{\tau_2}(\nu-1)(r_1 - \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}))} (\log M)^{\frac{1}{2} - \frac{1}{\tau_1}} \end{aligned} \quad (21)$$

in case when $\tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}) < r_1 < \frac{1}{2}$.

Since $\tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}) < r_1 < \frac{1}{2}$, (8) implies that

$$\begin{aligned} \|F_1 - G_M(F_1)\|_{p,\tau_2} &\leq CM^{-\frac{p}{2}(r_1 + \frac{1}{p} - \frac{1}{2})} (\log M)^{\frac{1}{2} - \frac{1}{\tau_1}} \leq \\ &\leq CM^{-\frac{p}{2}(r_1 + \frac{1}{p} - \frac{1}{2})} (\log M)^{\frac{p}{\tau_2}(\nu-1)(r_1 - \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}))} (\log M)^{\frac{1}{2} - \frac{1}{\tau_1}} \end{aligned} \quad (22)$$

(20)–(22) (see (18)) imply that

$$\begin{aligned} &\|f - (S_{Q_{n,\bar{\gamma}}}(f) + G_M(F_1) + G_M(F_3))\|_{p,\tau_2} \leq \\ &\leq \|F_1 - G_M(F_1)\|_{p,\tau_2} + \|F_3 - G_M(F_3)\|_{p,\tau_2} + \|F_2\|_{p,\tau_2} + \left\| \sum_{\langle \bar{s}, \bar{\gamma} \rangle \geq n_2} \delta_{\bar{s}}(f) \right\|_{p,\tau_2} \leq \\ &\leq CM^{-\frac{p}{2}(r_1 + \frac{1}{p} - \frac{1}{2})} (\log M)^{\frac{p}{\tau_2}(\nu-1)(r_1 - \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}))} (\log M)^{\frac{1}{2} - \frac{1}{\tau_1}} + \left\| \sum_{\langle \bar{s}, \bar{\gamma} \rangle \geq n_2} \delta_{\bar{s}}(f) \right\|_{p,\tau_2}. \end{aligned}$$

Then, taking into account that $\tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}) < r_1 < \frac{1}{2}$ and following the same steps as in [20], we have

$$\left\| \sum_{\langle \bar{s}, \bar{\gamma} \rangle \geq n_2} \delta_{\bar{s}}(f) \right\|_{p,\tau_2} \leq CM^{-\frac{p}{2}(r_1 + \frac{1}{p} - \frac{1}{2})} (\log M)^{\frac{p}{\tau_2}(\nu-1)(r_1 - \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}))} (\log M)^{\frac{1}{2} - \frac{1}{\tau_1}}.$$

Hence,

$$\begin{aligned} e_M(f)_{p,\tau_2} &\leq \|f - (S_{Q_{n,\bar{\gamma}}}(f) + G_M(F_1) + G_M(F_3))\|_{p,\tau_2} \leq \\ &\leq CM^{-\frac{p}{2}(r_1 + \frac{1}{p} - \frac{1}{2})} (\log M)^{\frac{p}{\tau_2}(\nu-1)(r_1 - \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}))} (\log M)^{\frac{1}{2} - \frac{1}{\tau_1}} \end{aligned}$$

for a function $f \in W_{2,\tau_1}^{\bar{r}}$ when $2 < p < \infty$, $2 < \tau_1 \leq \tau_2 < \infty$, $\tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2}) < r_1 < \frac{1}{2}$.

Let $1 < \tau_1 \leq 2$. Then by Lemma 1.5 [21] the inequality

$$\left(\sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} \|\delta_{\bar{s}}(f)\|_{2,\tau_1}^2 \right)^{1/2} \leq C \left\| \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} \delta_{\bar{s}}(f) \right\|_{2,\tau_1}. \quad (23)$$

Since $1 < \tau_1 \leq 2$, then (see [1; 217])

$$\|\delta_{\bar{s}}(f)\|_2 \leq C \|\delta_{\bar{s}}(f)\|_{2,\tau_1}. \quad (24)$$

It follows from inequalities (1), (23), and (24) that

$$\|F_1\|_A \leq C \sum_{l=n}^{n_1-1} 2^{l/2} \left\| \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} \delta_{\bar{s}}(f) \right\|_{2,\tau_1}.$$

Now, given that the function $f \in W_{2,\tau_1}^{\bar{r}}$ and the choice of the number n_1 , we get

$$\|F_1\|_A \leq CM^{-\frac{p}{2}(r_1-\frac{1}{2})}(\log M)^{(\nu-1)\frac{p}{\tau_2}(r_1-\frac{1}{2})}$$

for $r_1 < 1/2$. Further, arguing as in the proof of inequality (8), we obtain

$$\|F_1 - G_M(F_1)\|_{p,\tau_2} \leq CM^{-\frac{p}{2}(r_1+\frac{1}{p}-\frac{1}{2})}(\log M)^{\frac{1}{2}-\frac{1}{\tau_1}} \leq CM^{-\frac{p}{2}(r_1+\frac{1}{p}-\frac{1}{2})} \tag{25}$$

in case when $q = 2 < p < \infty$, $1 < \tau_1 \leq 2$, $1 < \tau_2 < \infty$, $r_1 < \frac{1}{2}$.

Let us estimate $\|F_3\|_A$. For this we set

$$\tilde{S}_l = \left(2^{lr_1\tau_1} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} \|\delta_{\bar{s}}(f)\|_2^2\right)^{1/2}$$

and

$$\tilde{m}_l := |G(l)| := \left[2^{-l\frac{\tau_2}{p}} \tilde{S}_l^2 2^{n\frac{\tau_2}{2}} n^{(\nu-1)\frac{\tau_2}{2}}\right] + 1.$$

In inequality (9) it is proved that

$$\begin{aligned} \|F_3\|_A &\leq 2^{-\frac{m}{2}} \sum_{l=n_1}^{n_2-1} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1, \bar{s} \in G(l)} 2^{\langle \bar{s}, \bar{1} \rangle \frac{1}{2}} \|\delta_{\bar{s}}(f)\|_2 \leq \\ &\leq 2^{-\frac{m}{2}} \sum_{l=n_1}^{n_2-1} 2^{(l+1)/2} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1, \bar{s} \in G(l)} \|\delta_{\bar{s}}(f)\|_2. \end{aligned} \tag{26}$$

Applying Hölder's inequality to the inner sum and substituting the value of the number $\tilde{m}_l := |G(l)|$ from (26), we obtain

$$\begin{aligned} \|F_3\|_A &\leq 2^{-\frac{m}{2}} \sum_{l=n_1}^{n_2-1} 2^{(l+1)/2} \left(\sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1, \bar{s} \in G(l)} \|\delta_{\bar{s}}(f)\|_2^2 \right)^{1/2} |G(l)|^{1/2} \times \\ &\times 2^{-\frac{m-1}{2}} \left\{ \sum_{l=n_1}^{n_2-1} 2^{l(\frac{1}{2}-r_1)} 2^{-l\frac{\tau_2}{2p}} \tilde{S}_l^2 (2^n n^{(\nu-1)})^{\frac{\tau_2}{4}} + \sum_{l=n_1}^{n_2-1} 2^{l(\frac{1}{2}-r_1)} \tilde{S}_l \right\}. \end{aligned} \tag{27}$$

Using inequalities (23) and (24) and taking into account the value of the numbers \tilde{S}_l , we obtain

$$\sum_{l=n_1}^{n_2-1} 2^{-l(r_1-\frac{1}{2}+\frac{\tau_2}{2p})} \tilde{S}_l^2 \leq \sum_{l=n_1}^{n_2-1} 2^{-l(r_1-\frac{1}{2}+\frac{\tau_2}{2p})} \left(2^{lr_1} \left\| \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1} \delta_{\bar{s}}(f) \right\|_{2,\tau_1}\right). \tag{28}$$

Since a function $f \in W_{2,\tau_1}^{\bar{r}}$ and

$$r_1 - \frac{1}{2} + \frac{\tau_2}{2p} = r_1 - \tau_2' \left(\frac{1}{2} - \frac{1}{p} + \frac{1}{2p} - \frac{1}{2\tau_2} \right) \leq r_1 - \tau_2' \left(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2} \right) < 0,$$

then from inequality (28) we have

$$\sum_{l=n_1}^{n_2-1} 2^{-l(r_1-\frac{1}{2}+\frac{\tau_2}{2p})} \tilde{S}_l^2 \leq C \sum_{l=n_1}^{n_2-1} 2^{-l(r_1-\tau_2'(\frac{1}{2}-\frac{1}{p}+\frac{1}{2p}-\frac{1}{2\tau_2}))} \leq C 2^{-n_2(r_1-\tau_2'(\frac{1}{2}-\frac{1}{p}+\frac{1}{2p}-\frac{1}{2\tau_2}))}. \tag{29}$$

Since the function $f \in W_{2,\tau_1}^{\bar{r}}$ and $r_1 - \frac{1}{2} < 0$, we can prove similarly that

$$\sum_{l=n_1}^{n_2-1} 2^{l(\frac{1}{2}-r_1)} \tilde{S}_l \leq C 2^{n_2(\frac{1}{2}-r_1)}. \tag{30}$$

Now it follows from inequalities (27), (29), and (30) that

$$\begin{aligned} \|F_3\|_A &\leq C \left\{ (2^n n^{\nu-1})^{\frac{\tau_2'}{4}} 2^{-n_2(r_1-\tau_2'(\frac{1}{2}-\frac{1}{p}+\frac{1}{2p}-\frac{1}{2\tau_2}))} + 2^{n_2(\frac{1}{2}-r_1)} \right\} \leq \\ &\leq C (2^n n^{\nu-1})^{-\frac{p}{2}(r_1-\frac{1}{2})} \end{aligned}$$

for a function $f \in W_{2,\tau_1}^{\bar{r}}$ when $2 < p < \infty$, $1 < \tau_1 \leq 2$ and $1 < \tau_2 < \infty$, $r_1 < \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{2p} - \frac{1}{2\tau_2})$.

Therefore, according to Lemma 1, for the function F_3 , by a constructive method, there is an M -term trigonometric polynomial $G_M(F_3)$ such that

$$\|F_3 - G_M(F_3)\|_{p,\tau_2} \leq C M^{-\frac{1}{2}} (2^n n^{\nu-1})^{-\frac{p}{2}(r_1-\frac{1}{2})} \leq C M^{-\frac{p}{2}(r_1+\frac{1}{p}-\frac{1}{2})} \tag{31}$$

for a function $f \in W_{2,\tau_1}^{\bar{r}}$ for $2 < p < \infty$, $1 < \tau_1 \leq 2$, $1 < \tau_2 < \infty$, $r_1 < \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{p\tau_1} - \frac{1}{2\tau_2})$.

Let us estimate $\|F_2\|_{p,\tau_2}$. To do this, note that if $\bar{s} \notin G(l)$, then

$$\|\delta_{\bar{s}}(f)\|_2 \leq \tilde{m}_l^{-\frac{1}{2}} 2^{-lr_1} \tilde{S}_l \tag{32}$$

and

$$\begin{aligned} \|F_2\|_{p,\tau_2} &\leq C \left(\sum_{l=n_1}^{n_2-1} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1, \bar{s} \notin G(l)} 2^{\langle \bar{s}, \bar{l} \rangle (\frac{1}{2}-\frac{1}{p})\tau_2} \|\delta_{\bar{s}}(f)\|_2^{\tau_2} \right)^{1/\tau_2} = \\ &= C \left(\sum_{l=n_1}^{n_2-1} \sum_{l \leq \langle \bar{s}, \bar{\gamma} \rangle < l+1, \bar{s} \notin G(l)} 2^{\langle \bar{s}, \bar{l} \rangle (\frac{1}{2}-\frac{1}{p})\tau_2} \|\delta_{\bar{s}}(f)\|_2^{\tau_2-2} \|\delta_{\bar{s}}(f)\|_2^2 \right)^{1/\tau_2}. \end{aligned}$$

Further, if $\tau_2 - 2 \geq 0$, then using inequality (32) and repeating the arguments of the proof (18), we obtain

$$\|F_2\|_{p,\tau_2} \leq C (2^n n^{\nu-1})^{-\frac{p}{2}(r_1+\frac{1}{p}-\frac{1}{2})} \leq C M^{-\frac{p}{2}(r_1+\frac{1}{p}-\frac{1}{2})} \tag{33}$$

for a function $f \in W_{2,\tau_1}^{\bar{r}}$ when $q = 2 < p < \infty$, $1 < \tau_1 \leq 2 \leq \tau_2 < \infty$, $r_1 < \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{2p} - \frac{1}{2\tau_2})$.

Now inequalities (25), (31), (33) imply that

$$e_M(f)_{p,\tau_2} \leq \|f - (S_{Q_{n,\bar{\gamma}}}(f) + G_M^p(F_1) + G_M^p(F_3))\|_{p,\tau_2} \leq C M^{-\frac{p}{2}(r_1+\frac{1}{p}-\frac{1}{q})} (\log M)^{\frac{1}{2}-\frac{1}{\tau_1}}$$

for a function $f \in W_{2,\tau_1}^{\bar{r}}$ when $2 < p < \infty$, $1 < \tau_1 \leq 2 \leq \tau_2 < \infty$, $r_1 < \tau_2'(\frac{1}{2} - \frac{1}{p} + \frac{1}{2p} - \frac{1}{2\tau_2})$. The proof is complete.

Remark 1. In case when $\tau_1 = 2$, Theorem 1 complements Theorem 4 in [14].

Acknowledgments

This research was funded by the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan (Grant No. AP22683029).

Author Contributions

G. Akishev and A.Kh. Myrzagaliyeva collected and analyzed data. Both authors participated in the revision of the manuscript and approved the final submission.

All authors contributed equally to this work.

Conflict of Interest

The authors declare no relevant financial or non-financial competing interests.

References

- 1 Stein E.M. Introduction to Fourier analysis on Euclidean spaces / E.M. Stein, G. Weiss. — Princeton: Princeton Univ. Press, 1971. — 312 p.
- 2 Temlyakov V.N. Multivariate approximation / V.N. Temlyakov. — Cambridge University Press, 2018. <https://doi.org/10.1017/9781108689687>
- 3 Темляков В.Н. О приближении периодических функций многих переменных / В.Н. Темляков // ДАН СССР. — 1984. — 279. — № 2. — С. 301–305.
- 4 Темляков В.Н. Приближение функций с ограниченной смешанной производной / В.Н. Темляков // Тр. Мат. ин-та им. В.А. Стеклова. — 1986. — 178. — С. 1–112.
- 5 Стечкин С.Б. Об абсолютной сходимости ортогональных рядов / С.Б. Стечкин // ДАН СССР. — 1955. — 102. — № 1. — С. 37–40.
- 6 Исмагилов Р.С. Поперечники множеств в линейных нормированных пространствах и приближение функций тригонометрическими многочленами / Р.С. Исмагилов // Успехи мат. наук. — 1974. — 29. — № 3. — С. 161–178.
- 7 Майоров В.Е. Тригонометрические поперечники соболевских классов W_p^r в пространстве L_q / В.Е. Майоров // Мат. заметки. — 1986. — 40. — № 2. — С. 161–173.
- 8 Makovoz Y. On trigonometric n -widths and their generalization / Y. Makovoz // J. Approx. Theory. — 1984. — 41. — No. 4. — P. 361–366.
- 9 Белинский Э.С. Приближение периодических функций с «плавающей» системой экспонент и тригонометрические поперечники / Э.С. Белинский // Исследования по теории функций многих вещественных переменных: сб. ст. — 1984. — С. 10–24.
- 10 Белинский Э.С. Приближение «плавающей» системой экспонент на классах периодических функций с ограниченной смешанной производной / Э.С. Белинский // Исследования по теории функций многих вещественных переменных: сб. ст. — 1988. — С. 16–33.
- 11 DeVore R.A. Nonlinear approximation / R.A. DeVore // Acta Numerica. — 1998. — 7. — P. 51–150. <https://doi.org/10.1017/S0962492900002816>
- 12 Dinh Dũng. Hyperbolic Cross Approximation. Advanced Courses in Mathematics. CRM Barcelona / Dinh Dũng, V.N. Temlyakov, T. Ullrich. — Birkhäuser/Springer, 2018. — 222 p.
- 13 Akishev G. Estimations of the best M -term approximations of functions in the Lorentz space with constructive methods / G. Akishev // Bulletin of the Karaganda University. Mathematics series. — 2017. — No. 3(87). — P. 13–26. <https://doi.org/10.31489/2017m3/13-26>
- 14 Akishev G. On estimates of M -term approximations on classes of functions with bounded mixed derivative in Lorentz spaces / G. Akishev, A. Myrzagaliyeva // Journal Math. Sci. — 2023. — P. 1–16. <https://doi.org/10.1007/s10958-022-06146-7>
- 15 Акишев Г. Оценки M -членных приближений функций класса $W_{q,\tau}^{a,b,\bar{r}}$ в пространстве Лоренца / Г. Акишев // Современные проблемы математического анализа и теории функций:

- Матер. Междунар. науч. конф., посвящ. 70-летию акад. НАН Таджикистана М.Ш. Шабозова. — 2022. — С. 22–25.
- 16 Temlyakov V.N. Constructive sparse trigonometric approximation and other problems for functions with mixed smoothness / V.N. Temlyakov // Sb. Math. — 2015. — 206. — P. 1628–1656. <https://doi.org/10.1070/SM2015v206n11ABEH004507>
- 17 Temlyakov V.N. Constructive sparse trigonometric approximation for functions with small mixed smoothness / V.N. Temlyakov // Constr. Approx. — 2017. — 45. — P. 467–495. <https://doi.org/10.1007/s00365-016-9345-3>
- 18 Bazarkhanov D.B. Nonlinear tensor product approximation of functions / D.B. Bazarkhanov, V.N. Temlyakov // Preprint ArXiv: 1409.1403v1 [stat ML]. — 2014. — P. 1–23.
- 19 Bazarkhanov D.B. Nonlinear trigonometric approximation of multivariate function classes / D.B. Bazarkhanov // Proceedings Steklov Inst. Math. — 2016. — 293. — P. 2–36. <https://doi.org/10.1134/S0081543816040027>
- 20 Акишев Г. Оценки наилучших приближений функций класса логарифмической гладкости в пространстве Лоренца / Г. Акишев // Тр. Ин-та математики и механики УрО РАН. — 2018. — 23. — С. 3–21. <https://doi.org/10.21538/0134-4889-2017-23-3-3-21>
- 21 Акишев Г. Оценки наилучших приближений функций класса Никольского-Бесова в пространстве Лоренца тригонометрическими полиномами / Г. Акишев // Тр. Ин-та математики и механики УрО РАН. — 2020. — 26. — С. 5–27. <https://doi.org/10.21538/0134-4889-2020-26-2-5-27>

Лоренц кеңістігіндегі Соболев класының M -мүшелік жуықтауларын бағалау туралы

Г. Акишев¹, А.Х. Мырзағалиева²

¹*М.В. Ломоносов атындағы Мәскеу мемлекеттік университетінің Қазақстан филиалы, Астана, Қазақстан;*
²*Astana IT University, Астана, Қазақстан*

Жұмыста бірнеше айнымалы периодты функциялар кеңістіктері зерделенген, атап айтқанда Лоренц кеңістігі $L_{2,\tau}(T^m)$, шектеулі аралас бөлшек туындысы бар функциялар класы $W_{2,\tau}^r$, $1 \leq \tau < \infty$ және $f \in L_{p,\tau}(T^m)$ функциясының тригонометриялық көпмүшеліктермен ең жақсы M -мүшелік жуықтауларының реті зерттелген. Мақала кіріспеден, негізгі бөлімнен және қорытындыдан тұрады. Кіріспеде негізгі нәтижелерді дәлелдеу үшін ұғымдар, анықтамалар және қажетті тұжырымдар қарастырылған. Сонымен қатар, осы тақырып бойынша алдыңғы зерттеулер жайлы ақпаратты табуға болады. Негізгі бөлімде W_{2,τ_1}^r Соболев класы функцияларының $L_{p,\tau_2}(T^m)$ кеңістігінің нормасы бойынша p, τ_1, τ_2 параметрлері арасындағы қатынастар үшін ең жақсы M -мүшелік жуықтауларының нақты реттік бағалаулары анықталған.

Кілт сөздер: Лоренц кеңістігі, Соболев класы, аралас туынды, тригонометриялық көпмүшеліктер, M -мүшелік жуықтау.

Об оценках M -членных приближений класса Соболева в пространстве Лоренца

Г. Акишев¹, А.Х. Мырзагалиева²

¹Казахстанский филиал Московского государственного университета имени М.В. Ломоносова, Астана, Казахстан;

²Astana IT University, Астана, Казахстан

В работе изучены пространства периодических функций нескольких переменных, а именно пространство Лоренца $L_{2,\tau}(T^m)$, класс функций с ограниченной смешанной дробной производной $W_{2,\tau}^r$, $1 \leq \tau < \infty$, и порядок наилучшего M -членного приближения функции $f \in L_{p,\tau}(T^m)$ тригонометрическими полиномами. Статья состоит из введения, основной части и заключения. Во введении рассмотрены основные понятия, определения и необходимые утверждения для доказательства основных результатов. Также можно найти информацию о предыдущих результатах по этой теме. В основной части установлены точные по порядку оценки для наилучших M -членных приближений функций класса Соболева W_{2,τ_1}^r по норме пространства $L_{p,\tau_2}(T^m)$ для различных соотношений между параметрами p, τ_1, τ_2 .

Ключевые слова: пространство Лоренца, класс Соболева, смешанная производная, тригонометрический полином, M -членное приближение.

References

- 1 Stein, E.M., & Weiss, G. (1971). *Introduction to Fourier analysis on Euclidean spaces*. Princeton Univ. Press.
- 2 Temlyakov, V. (2018). *Multivariate approximation*. Cambridge University Press. <https://doi.org/10.1017/9781108689687>
- 3 Temlyakov, V.N. (1984). О приближении периодических функций нескольких переменных [On the approximation of periodic functions of several variables]. *Doklady Akademii nauk SSSR — Reports of the USSR Academy of sciences*, 279(2), 301–305 [in Russian].
- 4 Temlyakov, V.N. (1986). Приближение функции с ограниченной смешанной производной [Approximation of functions with bounded mixed derivative]. *Trudy Ordena Lenina i Ordena Oktiabrskoi Revoliutsii Matematicheskogo instituta imeni V.A. Steklova — Proc. of the Order of Lenin and the Order of the October Revolution of the V.A. Steklov Mathematical Institute*, 178, 3–113 [in Russian].
- 5 Stechkin, S.B. (1955). Об абсолютной сходимости ортогональных рядов [On the absolute convergence of orthogonal series]. *Doklady Akademii nauk SSSR — Reports of the USSR Academy of Sciences*, 102(1), 37–40 [in Russian].
- 6 Ismagilov, R.S. (1974). Поперечники множеств в линейных нормированных пространствах и приближение функции тригонометрическими многочленами [Diameters of sets in normed linear spaces and the approximation of functions by trigonometric polynomials]. *Uspekhi matematicheskikh nauk — Russian Math. Surveys*, 29, 161–178 [in Russian].
- 7 Maiorov, V.E. (1986). Тригонометрические поперечники соболевских классов W_p^r в пространстве L_q [Trigonometric diameters of the Sobolev classes W_p^r in the space L_q]. *Matematicheskie zametki — Math. Notes*, 40(2), 161–173 [in Russian].
- 8 Makovoz, Y. (1984). On trigonometric n -widths and their generalization. *J. Approx. Theory*, 41, 361–366.
- 9 Belinsky, E.S. (1984). Приближение периодических функций с «плавающей» системой экспонент и тригонометрические поперечники [Approximation of periodic functions by a “floating” system of

- exponents and trigonometric diameters]. *Issledovaniia po teorii funktsii mnogikh veshchestvennykh peremennykh: sbornik statei — Research on the theory of functions of many real variables: collection of articles*, 10–24 [in Russian].
- 10 Belinsky, E.S. (1988). Priblizhenie «plavaiushchei» sistemoi eksponent na klassakh periodicheskikh funktsii s ogranichennoi smeshannoi proizvodnoi [Approximation by a “floating” system of exponentials on the classes of smooth periodic functions with bounded mixed derivative]. *Issledovaniia po teorii funktsii mnogikh veshchestvennykh peremennykh: sbornik statei — Research on the theory of functions of many real variables: collection of articles*, 16–33 [in Russian].
 - 11 DeVore, R.A. (1998). Nonlinear approximation. *Acta Numerica*, 7, 51–150. <https://doi.org/10.1017/S0962492900002816>
 - 12 Dinh, Dũng, Temlyakov, V.N., & Ullrich, T. (2018). *Hyperbolic Cross Approximation. Advanced Courses in Mathematics*. Birkhäuser/Springer.
 - 13 Akishev, G. (2017). Estimations of the best M -term approximations of functions in the Lorentz space with constructive methods. *Bulletin of the Karaganda University. Mathematics series*, 3(87), 13–26. <https://doi.org/10.31489/2017m3/13-26>
 - 14 Akishev, G., & Myrzagaliyeva, A. (2023). On estimates of M -term approximations on classes of functions with bounded mixed derivative in Lorentz spaces. *Journal Math. Sci.*, 1–16. <https://doi.org/10.1007/s10958-022-06146-7>
 - 15 Akishev, G. (2022). Otsenki M -chlenykh priblizhenii funktsii klassa $W_{q,\tau}^{a,b,\bar{r}}$ v prostranstve Lorentsa [Estimates for M -term approximations of functions of the class $W_{q,\tau}^{a,b,\bar{r}}$ in the Lorentz space]. *Materialy Mezhdunarodnoi nauchnoi konferentsii, posviashchennoi 70-letiiu akademika NAN Tadjikistana M.Sh. Shabozova — Materials of the international scientific conference dedicated to the 70th anniversary of Academician of the National Academy of Sciences of Tajikistan M.Sh. Shabozov*, 22–25 [in Russian].
 - 16 Temlyakov, V.N. (2015). Constructive sparse trigonometric approximation and other problems for functions with mixed smoothness. *Sb. Math.*, 206, 1628–1656. <https://doi.org/10.1070/SM2015v206n11ABEH004507>
 - 17 Temlyakov, V.N. (2017). Constructive sparse trigonometric approximation for functions with small mixed smoothness. *Constr. Approx.*, 45, 467–495. <https://doi.org/10.1007/s00365-016-9345-3>
 - 18 Bazarkhanov, D.B., & Temlyakov, V.N. (2014). Nonlinear tensor product approximation of functions. *Preprint ArXiv: 1409.1403v1 [stat ML]*, 1–23.
 - 19 Bazarkhanov, D.B. (2016). Nonlinear trigonometric approximation of multivariate function classes. *Proceedings Steklov Inst. Math.*, 293, 2–36. <https://doi.org/10.1134/S0081543816040027>
 - 20 Akishev, G. (2018). Otsenki nailuchshikh priblizhenii funktsii klassa logarifmicheskoi gladkosti v prostranstve Lorentsa [Estimates of best approximations of functions of the logarithmic smoothness class in the Lorentz space]. *Trudy Instituta matematiki i mekhaniki UrO RAN — Proceedings of the Institute of Mathematics and Mechanics of the Ural Russian Academy of Sciences*, 23, 3–21 [in Russian]. <https://doi.org/10.21538/0134-4889-2017-23-3-3-21>
 - 21 Akishev, G. (2020). Otsenki nailuchshikh priblizhenii klassa funktsii Nikolskogo–Besova v prostranstve Lorentsa trigonometricheskimi polinomami [Estimates of the best approximations of functions of the Nikolsky–Besov class in the Lorentz space by trigonometric polynomials]. *Trudy Instituta matematiki i mekhaniki Uralskogo otdeleniia RAN — Proceedings of the Institute of Mathematics and Mechanics of the Ural Russian Academy of Sciences*, 26, 5–27. <https://doi.org/10.21538/0134-4889-2020-26-2-5-27> [in Russian].

*Author Information**

Gabdolla Akishev — Doctor of physical and mathematical sciences, Professor, Kazakhstan Branch of Lomonosov Moscow State University, 11 Kazhymukan Street, Astana, 010010, Kazakhstan; e-mail: akishev_g@mail.ru; <https://orcid.org/0000-0002-8336-6192>

Aigul Khamzиеvna Myrzagaliyeva (*corresponding author*) — PhD, Assistant-professor, Astana IT University, 55/11 Mangilik El Avenue, Astana, 010000, Kazakhstan; e-mail: aigul.myrzagaliyeva@astanait.edu.kz; <https://orcid.org/0000-0002-4996-9483>

Buketov University

*The author's name is presented in the order: First, Middle and Last Names.