

About unimprovability the embedding theorems for anisotropic Nikol'skii-Besov spaces with dominated mixed derivatives and mixed metric and anisotropic Lorentz spaces

Y. Toleugazy¹, K.Y. Kervenev^{2,*}

¹*M.V. Lomonosov Moscow State University, Kazakhstan Branch, Astana, Kazakhstan;*

²*Karaganda Buketov university, Karaganda, Kazakhstan*

(E-mail: toleugazy.yerzhan@gmail.com, kervenev@bk.ru)

The embedding theory of spaces of differentiable functions of many variables studies important connections and relationships between differential (smoothness) and metric properties of functions and has wide application in various branches of pure mathematics and its applications. Earlier, we obtained the embedding theorems of different metrics for Nikol'skii-Besov spaces with a dominant mixed smoothness and mixed metric, and anisotropic Lorentz spaces. In this work, we showed that the conditions for the parameters of spaces in the above theorems are unimprovable. To do this, we built the extreme functions included in the spaces from the left sides of the embeddings and not included in the "slightly narrowed" spaces from the spaces in the right parts of the embeddings.

Keywords: anisotropic Lorentz spaces, anisotropic Nikol'skii-Besov spaces, generalized mixed smoothness, mixed metric, embedding theorems.

2020 Mathematics Subject Classification: 46E35.

Introduction

One of the first results related to the theory of embedding of spaces of differentiable functions was a result of S.L. Sobolev [1]. This theory studies important relations of differential (smoothness) properties of functions in various metrics. Further development of this theory is associated with new classes of function spaces defined and studied in the works of S.M. Nikol'skii [2], O.V. Besov [3], P.I. Lizorkin [4], H. Triebel [5], J. Bergh and J. Löfström [6], and many others. The development of this research was determined both by its internal problems and by its applications in the theory of boundary value problems of mathematical physics and approximation theory (see, for example, [7–11]).

In the 1960s, in the works of S.M. Nikol'skii [1], A.D. Dzhabrailov [12] and T.I. Amanov [13] begins the study of spaces with a dominant mixed derivative. Further study of spaces with a dominant mixed derivative which is related with the theory of embedding and interpolation and the theory of approximations is associated with the works of A.P. Uninskij, V.N. Temlyakov, E.D. Nursultanov, D.B. Bazarkhanov, A.S. Romanyuk, G.A. Akishev, K.A. Bekmaganbetov, Ye. Toleugazy and others (see, for example, [14–20]).

In a serie of articles [21–23] we studied various properties of Nikol'skii-Besov spaces with a dominant mixed derivative and with a mixed metric. In these articles, we investigated the interpolation properties of these spaces, obtained limit embedding theorems for these spaces and anisotropic Lorentz spaces, and proved theorems on traces and continuations of functions.

*Corresponding author. *E-mail:* kervenev@bk.ru

This research of Y. Toleugazy was supported in part by the Committee of Science of the Ministry of Science and Higher Education of the Republic of Kazakhstan (project AP14869553).

Received: 23 December 2023; *Accepted:* 20 February 2024.

© 2024 The Authors. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

In the work of K.A. Bekmaganbetov, K.E. Kervenev, Ye. Toleugazy [22], embeddings for Nikol'skii-Besov spaces with a dominant mixed derivative and a mixed metric and anisotropic Lorentz spaces were studied. In this article we are showing that the conditions in the embedding theorems from the work [22] are unimprovable. We build the extreme functions included in the spaces from the left sides of the embeddings and not included in the "slightly narrowed" spaces from spaces in the right parts of the embeddings.

Preliminaries and auxiliary results

Let $f(\mathbf{x}) = f(x_1, \dots, x_n)$ be a measurable function defined on \mathbb{T}^n . Let multiindexes $\mathbf{1} \leq \mathbf{p} = (p_1, \dots, p_n) \leq \infty$. A Lebesgue space $L_{\mathbf{p}}(\mathbb{T}^n)$ with mixed metric is the set of functions for which the following quantity is finite

$$\|f\|_{L_{\mathbf{p}}(\mathbb{T}^n)} = \left(\int_{\mathbb{T}} \left(\dots \left(\int_{\mathbb{T}} |f(x_1, \dots, x_n)|^{p_1} dx_1 \right)^{p_2/p_1} \dots \right)^{p_n/p_{n-1}} dx_n \right)^{1/p_n}.$$

Here, the expression $\left(\int_{\mathbb{T}} |f(t)|^p dt \right)^{1/p}$ for $p=\infty$ is understood as $\text{esssup}_{t \in \mathbb{T}} |f(t)|$.

For multiple trigonometric series $f(\mathbf{x}) \sim \sum_{\mathbf{k} \in \mathbb{Z}^n} a_{\mathbf{k}} e^{i(\mathbf{k}, \mathbf{x})}$ we denote by

$$\Delta_{\mathbf{s}}(f, \mathbf{x}) = \sum_{\mathbf{k} \in \rho(\mathbf{s})} a_{\mathbf{k}}(f) e^{i(\mathbf{k}, \mathbf{x})},$$

where $\rho(\mathbf{s}) = \{\mathbf{k} = (k_1, \dots, k_n) \in \mathbb{Z}^n : 2^{s_i-1} \leq |k_i| < 2^{s_i}, i = 1, \dots, n\}$, $(\mathbf{k}, \mathbf{x}) = \sum_{j=1}^n k_j x_j$ is the inner product of vectors \mathbf{k} and \mathbf{x} .

Let $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{R}^n$ and $\mathbf{1} \leq \tau = (\tau_1, \dots, \tau_n) \leq \infty$. The anisotropic Nikol'skii-Besov space with generalized mixed derivatives and mixed metric $B_{\mathbf{p}}^{\alpha, \tau}(\mathbb{T}^n)$ is a set of the series $f \sim \sum_{\mathbf{k} \in \mathbb{Z}^n} a_{\mathbf{k}} e^{i(\mathbf{k}, \mathbf{x})}$

such that

$$\|f\|_{B_{\mathbf{p}}^{\alpha, \tau}(\mathbb{T}^n)} = \left\| \left\{ 2^{(\alpha, \mathbf{s})} \|\Delta_{\mathbf{s}}(f)\|_{L_{\mathbf{p}}(\mathbb{T}^n)} \right\} \right\|_{l_{\tau}} < \infty,$$

where $\|\cdot\|_{l_{\tau}}$ is the norm of a discrete Lebesgue space with mixed metric l_{τ} .

We will also need the anisotropic Lorentz spaces which introduced by E.D. Nursultanov in [24].

Let $f(\mathbf{x}) = f(x_1, \dots, x_n)$ be a measurable function defined on \mathbb{T}^n . We denote by $f^*(\mathbf{t}) = f^{*1, \dots, *n}(t_1, \dots, t_n)$ the function obtained from $f(\mathbf{x}) = f(x_1, \dots, x_n)$ by applying the non-increasing rearrangement successively with respect to each of the variables x_1, \dots, x_n (the other variables are assumed to be fixed).

Let multiindexes $\mathbf{q} = (q_1, \dots, q_n)$, $\theta = (\theta_1, \dots, \theta_n)$ satisfy the conditions: if $0 < q_j < \infty$, then $0 < \theta_j \leq \infty$, if $q_j = \infty$, then $\theta_j = \infty$ for every $j = 1, \dots, n$. An anisotropic Lorentz space $L_{\mathbf{q}, \theta}(\mathbb{T}^n)$ is the set of functions for which the following quantity is finite

$$\begin{aligned} \|f\|_{L_{\mathbf{q}, \theta}(\mathbb{T}^n)} &= \\ &= \left(\int_{\mathbb{T}} \dots \left(\int_{\mathbb{T}} \left(t_1^{1/q_1} \dots t_n^{1/q_n} f^{*1, \dots, *n}(t_1, \dots, t_n) \right)^{\theta_1} \frac{dt_1}{t_1} \right)^{\theta_2/\theta_1} \dots \frac{dt_n}{t_n} \right)^{1/\theta_n}. \end{aligned}$$

The following theorems were obtained in the work [22]:

Theorem A. Let $-\infty < \alpha_0 = (\alpha_1^0, \dots, \alpha_n^0) \leq \alpha_1 = (\alpha_1^1, \dots, \alpha_n^1) < \infty$, $\mathbf{1} \leq \tau = (\tau_1, \dots, \tau_n) \leq \infty$ and $\mathbf{1} < \mathbf{p}_0 = (p_1^0, \dots, p_n^0)$, $\mathbf{p}_1 = (p_1^1, \dots, p_n^1) < \infty$. Then the embedding

$$B_{\mathbf{p}_1}^{\alpha_1 \tau}(\mathbb{T}^n) \hookrightarrow B_{\mathbf{p}_0}^{\alpha_0 \tau}(\mathbb{T}^n)$$

holds for $\alpha_0 - \mathbf{1}/\mathbf{p}_0 = \alpha_1 - \mathbf{1}/\mathbf{p}_1$.

Theorem B. Let $\mathbf{1} < \mathbf{p} = (p_1, \dots, p_n) < \mathbf{q} = (q_1, \dots, q_n) < \infty$ and $\mathbf{1} \leq \tau = (\tau_1, \dots, \tau_n) \leq \infty$. Then the embedding

$$B_{\mathbf{p}}^{\alpha \tau}(\mathbb{T}^n) \hookrightarrow L_{\mathbf{q}\tau}(\mathbb{T}^n)$$

holds for $\alpha = \mathbf{1}/\mathbf{p} - \mathbf{1}/\mathbf{q}$.

Theorem C. Let $\mathbf{1} < \mathbf{q} = (q_1, \dots, q_n) < \mathbf{p} = (p_1, \dots, p_n) < \infty$ and $\mathbf{1} \leq \tau = (\tau_1, \dots, \tau_n) \leq \infty$. Then the embedding

$$L_{\mathbf{q}\tau}(\mathbb{T}^n) \hookrightarrow B_{\mathbf{p}}^{\alpha \tau}(\mathbb{T}^n)$$

holds for $\alpha = \mathbf{1}/\mathbf{p} - \mathbf{1}/\mathbf{q}$.

Main results

In this work, we show that the conditions for the parameters providing attachments are unimprovable. The proof of these facts in Theorems A–C we carry out by constructing extreme functions.

The following theorem shows that the condition under which the embedding from Theorem A is valid is unimprovable.

Theorem 1. Let $-\infty < \alpha_0 = (\alpha_1^0, \dots, \alpha_n^0) \leq \alpha_1 = (\alpha_1^1, \dots, \alpha_n^1) < \infty$, $\mathbf{1} \leq \tau = (\tau_1, \dots, \tau_n) \leq \infty$, $\mathbf{1} < \mathbf{p}_0 = (p_1^0, \dots, p_n^0)$, $\mathbf{p}_1 = (p_1^1, \dots, p_n^1) < \infty$ and $\alpha_0 - \mathbf{1}/\mathbf{p}_0 = \alpha_1 - \mathbf{1}/\mathbf{p}_1$, then for arbitrary $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n) > \mathbf{0}$ and $\delta = (\delta_1, \dots, \delta_n) > \mathbf{0}$ there is a function $f_\beta^{(1)} \in B_{\mathbf{p}_1}^{\alpha_1 \tau}(\mathbb{T}^n)$ such that $f_\beta^{(1)} \notin B_{\mathbf{p}_0}^{(\alpha_0 + \varepsilon)\tau}(\mathbb{T}^n) \cup B_{(\mathbf{p}_0 + \delta)}^{\alpha_0 \tau}(\mathbb{T}^n)$.

Proof. Taking into account the estimate for the norm of a one-dimensional Dirichlet kernel, we obtain the relation

$$\left\| \sum_{k=2^{s-1}}^{2^s-1} e^{i(k, \cdot)} \right\|_{L_p(\mathbb{T})} \sim 2^{(1/p', s)}, \quad 1 < p < +\infty.$$

From this relation in the multiple case we have

$$\|\sigma_s(\cdot)\|_{L_{\mathbf{p}}(\mathbb{T}^n)} = \left\| \sum_{\mathbf{k}=2^{s-1}}^{2^s-1} e^{i(\mathbf{k}, \cdot)} \right\|_{L_{\mathbf{p}}(\mathbb{T}^n)} \sim 2^{(\mathbf{1}/\mathbf{p}', s)}. \tag{1}$$

Consider the function $f_\beta^{(1)}(\mathbf{x}) = \sum_{\mathbf{s}=\mathbf{0}}^{\infty} 2^{-(\beta, \mathbf{s})} \sigma_{\mathbf{s}}(\mathbf{x})$, where

$$\alpha_1 + \frac{\mathbf{1}}{\mathbf{p}_1'} < \beta < \min \left(\alpha_0 + \varepsilon + \frac{\mathbf{1}}{\mathbf{p}_0'}, \alpha_0 + \frac{\mathbf{1}}{(\mathbf{p}_0 + \delta)'} \right).$$

According to estimate (1) we have

$$\|f_\beta^{(1)}\|_{B_{\mathbf{p}_1}^{\alpha_1 \tau}(\mathbb{T}^n)} = \left(\sum_{\mathbf{s}=\mathbf{0}}^{\infty} \left(2^{(\alpha_1, \mathbf{s})} \|\Delta_{\mathbf{s}}(f_\beta^{(1)})\|_{L_{\mathbf{p}_1}(\mathbb{T}^n)} \right)^\tau \right)^{1/\tau} =$$

$$\begin{aligned}
 &= \left(\sum_{\mathbf{s}=0}^{\infty} \left(2^{(\alpha_1, \mathbf{s})} 2^{-(\beta, \mathbf{s})} \|\sigma_{\mathbf{s}}\|_{L_{\mathbf{p}_1}(\mathbb{T}^n)} \right)^{\tau} \right)^{1/\tau} = \\
 &= \left(\sum_{\mathbf{s}=0}^{\infty} \left(2^{(\alpha_1 - \beta + \frac{1}{\mathbf{p}_1}, \mathbf{s})} \right)^{\tau} \right)^{1/\tau} < +\infty,
 \end{aligned}$$

as $\alpha_1 + \frac{1}{\mathbf{p}_1} - \beta < \mathbf{0}$.

This means that $f_{\beta}^{(1)} \in B_{\mathbf{p}_0}^{\alpha_0 \tau}(\mathbb{T}^n)$.

Similarly, we obtain that

$$\begin{aligned}
 \|f_{\beta}^{(1)}\|_{B_{\mathbf{p}_0}^{(\alpha_0 + \varepsilon)\tau}(\mathbb{T}^n)} &= \left(\sum_{\mathbf{s}=0}^{\infty} \left(2^{(\alpha_0 + \varepsilon, \mathbf{s})} \|\Delta_{\mathbf{s}}(f_{\beta}^{(1)})\|_{L_{\mathbf{p}_0}(\mathbb{T}^n)} \right)^{\tau} \right)^{1/\tau} \geq \\
 &\geq C_1 \left(\sum_{\mathbf{s}=0}^{\infty} \left(2^{(\alpha_0 + \varepsilon, \mathbf{s})} 2^{(\frac{1}{\mathbf{p}_0} - \beta, \mathbf{s})} \right)^{\tau} \right)^{1/\tau} = \\
 &= C_1 \left(\sum_{\mathbf{s}=0}^{\infty} \left(2^{(\alpha_0 + \varepsilon + \frac{1}{\mathbf{p}_0} - \beta, \mathbf{s})} \right)^{\tau} \right)^{1/\tau} = +\infty,
 \end{aligned}$$

as $\alpha_0 + \varepsilon + \frac{1}{\mathbf{p}_0} - \beta > \mathbf{0}$. Therefore $f_{\beta}^{(1)} \notin B_{\mathbf{p}_0}^{(\alpha_0 + \varepsilon)\tau}(\mathbb{T}^n)$.

Further, we will show that $f_{\beta}^{(1)} \notin B_{(\mathbf{p}_0 + \delta)}^{\alpha_0 \tau}(\mathbb{T}^n)$. We have

$$\begin{aligned}
 \|f_{\beta}^{(1)}\|_{B_{(\mathbf{p}_0 + \delta)}^{\alpha_0 \tau}(\mathbb{T}^n)} &= \left(\sum_{\mathbf{s}=0}^{\infty} \left(2^{(\alpha_0, \mathbf{s})} \|\Delta_{\mathbf{s}}(f_{\beta}^{(1)})\|_{L_{(\mathbf{p}_0 + \delta)}(\mathbb{T}^n)} \right)^{\tau} \right)^{1/\tau} \geq \\
 &\geq C_2 \left(\sum_{\mathbf{s}=0}^{\infty} \left(2^{(\alpha_0, \mathbf{s})} 2^{(\frac{1}{(\mathbf{p}_0 + \delta)} - \beta, \mathbf{s})} \right)^{\tau} \right)^{1/\tau} = \\
 &= C_2 \left(\sum_{\mathbf{s}=0}^{\infty} \left(2^{(\alpha_1 + \frac{1}{(\mathbf{p}_0 + \delta)} - \beta, \mathbf{s})} \right)^{\tau} \right)^{1/\tau} = +\infty,
 \end{aligned}$$

considering that $\alpha_0 + \frac{1}{(\mathbf{p}_0 + \delta)} - \beta > \mathbf{0}$. Therefore $f_{\beta}^{(1)} \notin B_{(\mathbf{p}_0 + \delta)}^{\alpha_0 \tau}(\mathbb{T}^n)$.

Thus, we have shown that $f_{\beta}^{(1)} \notin B_{\mathbf{p}_0}^{(\alpha_0 + \varepsilon)\tau}(\mathbb{T}^n) \cup B_{(\mathbf{p}_0 + \delta)}^{\alpha_0 \tau}(\mathbb{T}^n)$.

The proof is complete.

The following theorem shows that the condition under which the embedding of Theorem B is valid, is not improved.

Theorem 2. Let $\mathbf{1} < \mathbf{p} = (p_1, \dots, p_n) < \mathbf{q} = (q_1, \dots, q_n) < \infty$ and $\alpha = \mathbf{1}/\mathbf{p} - \mathbf{1}/\mathbf{q}$. Then for an arbitrary $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n) > \mathbf{0}$ there is a function $f_{\beta}^{(2)} \in B_{\mathbf{p}}^{\alpha \tau}(\mathbb{T}^n)$ such that $f_{\beta}^{(2)} \notin L_{\mathbf{q} + \varepsilon, \tau}(\mathbb{T}^n)$.

Proof. First, let's show that $f_{\beta}^{(2)} \in B_{\mathbf{p}}^{\alpha \tau}(\mathbb{T}^n)$. Consider the function $f_{\beta}^{(2)}(\mathbf{x}) = \sum_{\mathbf{s}=0}^{\infty} 2^{-(\beta, \mathbf{s})} \sigma_{\mathbf{s}}(\mathbf{x})$,

where $\alpha + \frac{1}{\mathbf{p}'} < \beta \leq \frac{1}{(\mathbf{q} + \varepsilon)'}$.

By analogy with Theorem 1, we have

$$\|f_\beta^{(2)}\|_{B_{\mathbf{p}^{\alpha\tau}}(\mathbb{T}^n)} \sim \left(\sum_{s=0}^{\infty} \left(2^{(\alpha-\beta+\frac{1}{\mathbf{p}'},s)}\right)^\tau\right)^{1/\tau} < \infty$$

as $\alpha + \frac{1}{\mathbf{p}'} - \beta < \mathbf{0}$. It means that $f_\beta^{(2)} \in B_{\mathbf{p}^{\alpha\tau}}(\mathbb{T}^n)$.

In order to show that $f_\beta^{(2)} \notin L_{\mathbf{q}+\varepsilon,\tau}(\mathbb{T}^n)$ we use Theorem C.

We have

$$\begin{aligned} \|f_\beta^{(2)}\|_{L_{\mathbf{q}+\varepsilon,\tau}(\mathbb{T}^n)} &\geq \|f_\beta^{(2)}\|_{B_{\frac{1}{\mathbf{p}}-\frac{1}{\mathbf{q}+\varepsilon},\tau}(\mathbb{T}^n)} = \\ &= \left(\sum_{s=0}^{\infty} \left(2^{\left(\frac{1}{\mathbf{p}}-\frac{1}{\mathbf{q}+\varepsilon},s\right)} \|\Delta_s(f_\beta^{(2)})\|_{L_{\mathbf{p}}(\mathbb{T}^n)}\right)^\tau\right)^{1/\tau} = \\ &= C_3 \left(\sum_{s=0}^{\infty} \left(2^{\left(\frac{1}{\mathbf{p}}-\frac{1}{\mathbf{q}+\varepsilon},s\right)} 2^{\left(\frac{1}{\mathbf{p}'}-\beta,s\right)}\right)^\tau\right)^{1/\tau} = \\ &= C_3 \left(\sum_{s=0}^{\infty} \left(2^{\left(\frac{1}{\mathbf{p}}+\frac{1}{\mathbf{p}'}-\frac{1}{\mathbf{q}+\varepsilon}-\beta,s\right)}\right)^\tau\right)^{1/\tau} = \\ &= C_3 \left(\sum_{s=0}^{\infty} \left(2^{\left(1-\frac{1}{\mathbf{q}+\varepsilon}-\beta,s\right)}\right)^\tau\right)^{1/\tau} = C_3 \left(\sum_{s=0}^{\infty} \left(2^{\left(\frac{1}{(\mathbf{q}+\varepsilon)'}-\beta,s\right)}\right)^\tau\right)^{1/\tau} = +\infty, \end{aligned}$$

as $\frac{1}{(\mathbf{q}+\varepsilon)'} - \beta > 0$. It means that $f_\beta^{(2)} \notin L_{\mathbf{q}+\varepsilon,\tau}(\mathbb{T}^n)$.

The proof is complete.

The following theorem shows that the condition, under which the embedding of Theorem C is valid, is not improved.

Theorem 3. Let $\mathbf{1} < \mathbf{q} = (q_1, \dots, q_n) < \mathbf{p} = (p_1, \dots, p_n) < \infty$ and $\alpha = \mathbf{1}/\mathbf{p} - \mathbf{1}/\mathbf{q}$. Then for an arbitrary $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n) > \mathbf{0}$ and $\delta = (\delta_1, \dots, \delta_n) > \mathbf{0}$ there is a function $f_\beta^{(3)} \in L_{\mathbf{q},\tau}(\mathbb{T}^n)$ such that $f_\beta^{(3)} \notin B_{\mathbf{p}^{\alpha+\varepsilon,\tau}}(\mathbb{T}^n) \cup B_{(\mathbf{p}+\delta)'}(\mathbb{T}^n)$.

Proof. Let's choose a function $f_\beta^{(3)}(\mathbf{x})$ the same as in Theorem 1 with β , satisfying the condition $\frac{\mathbf{1}}{\mathbf{q}'} < \beta \leq \min\left(\alpha + \varepsilon + \frac{\mathbf{1}}{\mathbf{p}'}, \alpha + \frac{\mathbf{1}}{(\mathbf{p} + \delta)'}\right)$.

In order to show that $f_\beta^{(3)} \in L_{\mathbf{q},\tau}(\mathbb{T}^n)$ let's use Theorem B. We have

$$\begin{aligned} \|f_\beta^{(3)}\|_{L_{\mathbf{q},\tau}(\mathbb{T}^n)} &\leq C_4 \|f_\beta^{(3)}\|_{B_{\left(\frac{1}{\mathbf{p}}-\frac{1}{\mathbf{q}}\right)\tau}(\mathbb{T}^n)} = \\ &= C_4 \left(\sum_{s=0}^{\infty} \left(2^{\left(\frac{1}{\mathbf{p}}-\frac{1}{\mathbf{q}},s\right)} \|\Delta_s(f_\beta^{(3)})\|_{L_{\mathbf{p}}(\mathbb{T}^n)}\right)^\tau\right)^{1/\tau} \leq \\ &\leq C_5 \left(\sum_{s=0}^{\infty} \left(2^{\left(\frac{1}{\mathbf{p}}-\frac{1}{\mathbf{q}},s\right)} 2^{\left(-\beta+\frac{1}{\mathbf{p}'},s\right)}\right)^\tau\right)^{1/\tau} = \\ &= C_5 \left(\sum_{s=0}^{\infty} \left(2^{\left(\frac{1}{\mathbf{p}}+\frac{1}{\mathbf{p}'}-\beta-\frac{1}{\mathbf{q}},s\right)}\right)^\tau\right)^{1/\tau} = \end{aligned}$$

$$= C_5 \left(\sum_{s=0}^{\infty} \left(2^{(1-\frac{1}{q}-\beta, s)} \right)^\tau \right)^{1/\tau} = C_5 \left(\sum_{s=0}^{\infty} \left(2^{(\frac{1}{q}-\beta, s)} \right)^\tau \right)^{1/\tau} < +\infty,$$

as $\frac{1}{q} - \beta < \mathbf{0}$, i.e. $f_\beta^{(3)} \in L_{\mathbf{q}, \tau}(\mathbb{T}^n)$.

Let's show that $f_\beta \notin B_{\mathbf{p}}^{\alpha \mathbf{q}}(\mathbb{T}^n)$.

Let us estimate the norm of this function from below

$$\begin{aligned} \|f_\beta^{(3)}\|_{B_{\mathbf{p}}^{(\alpha+\varepsilon)\tau}(\mathbb{T}^n)} &= \left(\sum_{s=0}^{\infty} \left(2^{(\alpha+\varepsilon, s)} \|\Delta_s(f_\beta^{(3)})\|_{L_{\mathbf{p}}(\mathbb{T}^n)} \right)^\tau \right)^{1/\tau} \geq \\ &\geq C_6 \left(\sum_{s=0}^{\infty} \left(2^{(\alpha+\varepsilon, s)} 2^{(\frac{1}{p'}-\beta, s)} \right)^\tau \right)^{1/\tau} = \\ &= C_6 \left(\sum_{s=0}^{\infty} \left(2^{(\alpha+\varepsilon+\frac{1}{p'}-\beta, s)} \right)^\tau \right)^{1/\tau} = +\infty, \end{aligned}$$

as $\alpha + \varepsilon - \beta + \frac{1}{p'} > \mathbf{0}$, i.e. $f_\beta^{(3)} \notin B_{\mathbf{p}}^{(\alpha+\varepsilon)\tau}(\mathbb{T}^n)$.

Further, we will show that $f_\beta^{(3)} \notin B_{(\mathbf{p}_0+\delta)}^{\alpha_0 \tau}(\mathbb{T}^n)$.

Considering that $\alpha_0 + \frac{1}{(\mathbf{p}_0+\delta)'} - \beta > \mathbf{0}$, we have

$$\begin{aligned} \|f_\beta^{(3)}\|_{B_{(\mathbf{p}_0+\delta)}^{\alpha_0 \tau}(\mathbb{T}^n)} &= \left(\sum_{s=0}^{\infty} \left(2^{(\alpha_0, s)} \|\Delta_s(f_\beta^{(3)})\|_{L_{(\mathbf{p}_0+\delta)}(\mathbb{T}^n)} \right)^\tau \right)^{1/\tau} \geq \\ &\geq C_7 \left(\sum_{s=0}^{\infty} \left(2^{(\alpha_0, s)} 2^{(\frac{1}{(\mathbf{p}_0+\delta)'}-\beta, s)} \right)^\tau \right)^{1/\tau} = \\ &= C_7 \left(\sum_{s=0}^{\infty} \left(2^{(\alpha_1+\frac{1}{(\mathbf{p}_0+\delta)'}-\beta, s)} \right)^\tau \right)^{1/\tau} = +\infty, \end{aligned}$$

as $f_\beta^{(3)} \notin B_{(\mathbf{p}_0+\delta)}^{\alpha_0 \tau}(\mathbb{T}^n)$.

Thus, we have shown that $f_\beta^{(3)} \notin B_{\mathbf{p}_0}^{(\alpha_0+\varepsilon)\tau}(\mathbb{T}^n) \cup B_{(\mathbf{p}_0+\delta)}^{\alpha_0 \tau}(\mathbb{T}^n)$.

The proof is complete.

Author Contributions

All authors contributed equally to this work.

Conflict of Interest

The authors declare no conflict of interest.

References

- 1 Соболев С. Об одной теореме функционального анализа / С. Соболев // Мат. сб. — 1938. — Т. 46, № 4. — С. 471–497.

- 2 Nikol'skii S.M. Approximation of functions of several variables and imbedding theorems / S.M. Nikol'skii. — Heidelberg: Springer-Verlag, 1975. — 418 p.
- 3 Besov O.V. Integral representations of functions and imbedding theorems / O.V. Besov, V.P. Il'in, S.M. Nikol'skii. — New York – Toronto – London: J. Wiley and Sons, 1979. — 311 p.
- 4 Лизоркин П.И. Пространства $L_p^r(\Omega)$. Теоремы продолжения и вложения / П.И. Лизоркин // ДАН СССР. — 1962. — Т. 145, № 3. — С. 527–530.
- 5 Triebel H. Theory of interpolation, function spaces, differential operators / H. Triebel. — Berlin: VEB Deutscher Verlag der Wissenschaften, 1978. — 528 p.
- 6 Bergh J. Interpolation spaces. An introduction / J. Bergh, J. Löfström. — Berlin – Heidelberg – New York: Springer-Verlag, 1976. — 207 p.
- 7 Bekmaganbetov K.A. Homogenization of random attractors for reaction-diffusion systems / K.A. Bekmaganbetov, G.A. Chechkin, V.V. Chepyzhov // CR Mecanique. — 2016. — Vol. 344, No. 11–12. — P. 753–758. <https://doi.org/10.1016/j.crme.2016.10.015>
- 8 Bekmaganbetov K.A. Homogenization of trajectory attractors of 3D Navier-Stokes system with randomly oscillating force / K.A. Bekmaganbetov, G.A. Chechkin, V.V. Chepyzhov, A.Yu. Goritsky // Discrete Contin. Dyn. Syst. — 2017. — Vol. 37, No 5. — P. 2375–2393. <https://doi.org/10.3934/dcds.2017103>
- 9 Bekmaganbetov K.A. Weak convergence of attractors of reaction–diffusion systems with randomly oscillating coefficients / K.A. Bekmaganbetov, G.A. Chechkin, V.V. Chepyzhov // Applicable Analysis. — 2019. — Vol. 98, No. 1–2. — P. 256–271. <https://doi.org/10.1080/00036811.2017.1400538>
- 10 Nursultanov E. Norm convolution inequalities in Lebesgue spaces / E. Nursultanov, S. Tikhonov, N. Tleukhanova // Revista Matematica Iberoamericana: Universidad Autonoma de Madrid. — 2018. — Vol. 34, No. 2. — P. 811–838.
- 11 Акишев Г.А. Неравенство разных метрик в обобщенном пространстве Лоренца / Г.А. Акишев // Тр. ИММ УрО РАН. — 2018. — Т. 24, № 4. — С. 5–18. <https://doi.org/10.21538/0134-4889-2018-24-4-5-18>
- 12 Джабраилов А.Д. О некоторых функциональных пространствах. Прямые и обратные теоремы вложения / А.Д. Джабраилов // ДАН СССР. — 1964. — Т. 159, № 2. — С. 254–257.
- 13 Аманов Т.И. Теоремы представления и вложения для функциональных пространств $S_{p,\theta}^{(r)}B(\mathbb{R}^n)$ и $S_{p,\theta}^{(r)}B(0 \leq x_j \leq 2\pi; j = 1, \dots, n)$ / Т.И. Аманов // Тр. МИАН СССР. — 1965. — Т. 77. — С. 5–34.
- 14 Унинский А.П. Теоремы вложения для класса функций со смешанной нормой / А.П. Унинский // ДАН СССР. — 1966. — Т. 166, № 4. — С. 806–808.
- 15 Temlyakov V.N. Approximation of functions with small mixed smoothness in the uniform norm / V.N. Temlyakov, T. Ullrich // J. Approx. Theory. — 2022. — Vol. 277, article No. 105718. — 23 p. <https://doi.org/10.1016/j.jat.2022.105718>
- 16 Базарханов Д.Б. Нелинейные тригонометрические приближения классов функций многих переменных / Д.Б. Базарханов // Тр. МИАН. — 2016. — Т. 293. — С. 8–42. <https://doi.org/10.1134/S0371968516020023>
- 17 Romanyuk A.S. Kolmogorov and trigonometric widths of the Besov classes $B_{p,\theta}^r$ of multivariate periodic functions // Sbornik: Math. — 2006. — Vol. 197, No. 1. — P. 69–93.
- 18 Bekmaganbetov K.A. Interpolation of Besov $B_{pt}^{\sigma q}$ and Lizorkin-Triebel $F_{pt}^{\sigma q}$ spaces / K.A. Bekmaganbetov, E.D. Nursultanov // Analysis Mathematica. — 2009. — Vol. 35, No. 3. — P. 169–188. <https://doi.org/10.1007/s10476-009-0301-3>
- 19 Bekmaganbetov K.A. Embedding theorems for anisotropic Besov spaces $B_{\text{pr}}^{\alpha q}([0, 2\pi]^n)$ / K.A. Bekmaganbetov, E.D. Nursultanov // Izvestiia: Mathematics. — 2009. — Vol. 73, No. 4. —

- P. 655–668. <https://doi.org/10.4213/im2741>
- 20 Bekmaganbetov K.A. Order of the orthoprojection widths of the anisotropic Nikol'skii-Besov classes in the anisotropic Lorentz space / K.A. Bekmaganbetov, Ye. Toleugazy // Eurasian Math. J. — 2016. — Vol. 7, No. 3. — P. 8–16.
- 21 Bekmaganbetov K.A. Interpolation theorem for Nikol'skii-Besov type spaces with mixed metric / K.A. Bekmaganbetov, K.Ye. Kervenev, Ye. Toleugazy // Bulletin of the Karaganda university. Mathematics series. — 2020. — No. 4(100). — P. 33–42. <https://doi.org/10.31489/2020M4/33-42>
- 22 Bekmaganbetov K.A. The embedding theorems for Nikol'skii-Besov spaces with generalized mixed smoothness / K.A. Bekmaganbetov, K.Ye. Kervenev, Ye. Toleugazy // Bulletin of the Karaganda university. Mathematics series. — 2021. — No. 4(104). — P. 28–34. <https://doi.org/10.31489/2021M4/28-34>
- 23 Bekmaganbetov K.A. The theorems about traces and extensions for functions from Nikol'skii-Besov spaces with generalized mixed smoothness / K.A. Bekmaganbetov, K.Ye. Kervenev, Ye. Toleugazy // Bulletin of the Karaganda university. Mathematics series. — 2022. — No. 4(108). — P. 42–50. <https://doi.org/10.31489/2022M4/42-50>
- 24 Нурсултанов Е.Д. Интерполяционные теоремы для анизотропных функциональных пространств и их приложения / Е.Д. Нурсултанов // ДАН СССР — 2004. — Т. 394, № 1. — С. 22–25.

Үстем аралас туындысы және аралас метрикасы бар анизотропты Никольский-Бесов кеңістіктері және анизотропты Лоренц кеңістіктері үшін ену теоремаларының жетілдірілмейтіндігі туралы

Е. Төлеуғазы¹, Қ.Е. Кервенев²

¹М.В. Ломоносов атындағы Мәскеу мемлекеттік университетінің Қазақстан филиалы, Астана, Қазақстан;

²Академик Е.А. Бөкетов атындағы Қарағанды университеті, Қарағанды, Қазақстан
(E-mail: toleugazy.yerzhan@gmail.com, kervenev@bk.ru)

Дифференциалданатын функциялар кеңістіктерінің енгізу теориясы әртүрлі метрикалардағы функциялардың дифференциалдық (тегістіліктік) қасиеттерінің маңызды байланыстары мен қатынастарын зерттейді. Математикалық физиканың шектік есептер теориясында, жуықтау теориясында және математиканың басқа да салаларында кеңінен қолданысқа ие. Мақалада үстем аралас тегістілігі және аралас метрикасы бар Никольский-Бесовтың кеңістіктері үшін және Лоренцтің анизотропты кеңістіктері үшін енгізу теоремалары берілген. Ұсынылған жұмыста жоғарыда көрсетілген теоремалардағы параметрлердің жетілдірілмейтіндігі көрсетілді. Осыны көрсетуге біз сол жақтағы енулердегі кеңістіктер үшін шекті функцияларды құрамыз және олар оң жақтағы енулерде “сәл ғана жіңішкертілген” кеңістіктерде жатпайтындығы көрсетілген.

Клт сөздер: Лоренцтің анизотропты кеңістіктері, Никольский-Бесов типтес кеңістіктер, үстем аралас туынды, аралас метрика, ену теоремалары.

О неулучшаемости теорем вложения для анизотропных пространств Никольского-Бесова с доминирующей смешанной производной и смешанной метрикой и анизотропных пространств Лоренца

Е. Толеугазы¹, К.Е. Кервенеv²

¹Казахстанский филиал Московского государственного университета имени М.В. Ломоносова, Астана, Казахстан;

²Карагандинский университет имени академика Е.А. Букетова, Караганда, Казахстан
(E-mail: toleugazy.yerzhan@gmail.com, kervenev@bk.ru)

Теория вложения пространств дифференцируемых функций многих переменных изучает важные связи и соотношения между дифференциальными (гладкостными) и метрическими свойствами функций и имеет широкое применение в различных разделах чистой математики и ее приложениях. Ранее нами были получены предельные теоремы вложения разных метрик для пространств Никольского-Бесова с доминирующей смешанной гладкостью и со смешанной метрикой и для анизотропных пространств Лоренца. В данной работе мы показали, что условия на параметры пространств в отмеченных выше теоремах являются неулучшаемыми. Для этого мы построили крайние функции, входящие в пространства из левых сторон вложений и не входящие в «немного зауженные» пространства, чем пространства, стоящие в правых частях вложений.

Ключевые слова: анизотропные пространства Лоренца, пространства Никольского-Бесова, доминирующая смешанная производная, смешанная метрика, теоремы вложения.

References

- 1 Sobolev, S. (1938). Ob odnoi teoreme funktsionalnogo analiza [On a theorem of functional analysis]. *Matematicheskii sbornik — Math. Coll.*, 46(4), 471–497 [in Russian].
- 2 Nikol'skii, S.M. (1975). *Approximation of functions of several variables and imbedding theorems*. Heidelberg: Springer-Verlag.
- 3 Besov, O.V., Il'in, V.P., & Nikol'skii, S.M. (1979). *Integral representations of functions and imbedding theorems*. New York – Toronto – London: J. Wiley and Sons.
- 4 Lizorkin, P.I. (1962). Prostranstva $L_p^r(\Omega)$. Teoremy prodolzheniia i vlozheniia [$L_p^r(\Omega)$ spaces. Extension and imbedding theorems]. *Doklady Akademii nauk SSSR — Soviet Math. Dokl.*, 145(3), 527–530 [in Russian].
- 5 Triebel, H. (1978). *Theory of interpolation, function spaces, differential operators*. Berlin: VEB Deutscher Verlag der Wissenschaften.
- 6 Bergh, J., & Löfström, J. (1976). *Interpolation spaces. An introduction*. Berlin – Heidelberg – New York: Springer-Verlag.
- 7 Bekmaganbetov, K.A., Chechkin, G.A., & Chepyzhov, V.V. (2016). Homogenization of random attractors for reaction-diffusion systems. *CR Mecanique*, 344(11-12), 753–758. <https://doi.org/10.1016/j.crme.2016.10.015>
- 8 Bekmaganbetov, K.A., Chechkin, G.A., Chepyzhov, V.V., & Goritsky, A.Yu. (2017). Homogenization of trajectory attractors of 3D Navier-Stokes system with randomly oscillating force. *Discrete Contin. Dyn. Syst.*, 37(5), 2375–2393. <https://doi.org/10.3934/dcds.2017103>
- 9 Bekmaganbetov, K.A., Chechkin, G.A., & Chepyzhov, V.V. (2019). Weak convergence of attractors of reaction-diffusion systems with randomly oscillating coefficients. *Applicable Analysis*, 98(1–2), 256–271. <https://doi.org/10.1080/00036811.2017.1400538>

- 10 Nursultanov, E., Tikhonov, S., & Tleukhanova, N. (2018). Norm convolution inequalities in Lebesgue spaces. *Revista Matemática Iberoamericana: Universidad Autónoma de Madrid (Spain)*, 34(2), 811–838.
- 11 Akishev, G.A. (2018). Neravenstvo raznykh metrik v obobshchennom prostranstve Lorentsa [Inequality of different metrics in the generalized Lorentz space]. *Trudy Instituta matematiki i mekhaniki Uralskogo otdeleniia Rossiiskoi akademii nauk — Proceedings of the Institute of Mathematics and Mechanics of the Ural Branch of the Russian Academy of Sciences*, 24(4), 5–18 [in Russian]. 10.21538/0134-4889-2018-24-4-5-18
- 12 Dzhabrailov, A.D. (1964). O nekotorykh funktsionalnykh prostranstvakh. Priamye i obratnye teoremy vlozheniia [About some functional spaces. Direct and inverse theories of implementation]. *Doklady Akademii nauk SSSR — Soviet Math. Dokl.*, 159(2), 254–257 [in Russian].
- 13 Amanov, T.I. (1965). Teoremy predstavleniia i vlozheniia dlia funktsionalnykh prostranstv $S_{p,\theta}^{(r)}B(\mathbb{R}^n)$ i $S_{p,\theta}^{(r)}B(0 \leq x_j \leq 2\pi; j = 1, \dots, n)$ [Representation and embedding theorems for function spaces $S_{p,\theta}^{(r)}B(\mathbb{R}^n)$ and $S_{p,\theta}^{(r)}B(0 \leq x_j \leq 2\pi; j = 1, \dots, n)$]. *Trudy MIAN SSSR — Proceedings of the V.A. Steklov Mathematical Institute*, 77, 5–34 [in Russian].
- 14 Uninskii, A.P. (1966). Teoremy vlozheniia dlia klassa funktsii so smeshannoi normoi [Embedding theorems for a class of functions with a mixed norm]. *Doklady Akademii nauk SSSR — Soviet Math. Dokl.*, 166(4), 806–808 [in Russian].
- 15 Temlyakov, V.N., & Ullrich, T. (2022). Approximation of functions with small mixed smoothness in the uniform norm. *Journal of Approximation Theory*, 277(105718), 23 p. <https://doi.org/10.1016/j.jat.2022.105718>
- 16 Bazarkhanov, D.B. (2016). Nelineinye trigonometricheskie priblizheniia klassov funktsii mnogikh peremennykh [Nonlinear trigonometric approximations of classes of functions of many variables]. *Trudy MIAN — Proceedings of the V.A. Steklov Mathematical Institute*, 293, 8–42 [in Russian]. <https://doi.org/10.1134/S0371968516020023>
- 17 Romanyuk, A.S. (2006). Kolmogorov and trigonometric widths of the Besov classes $B_{p,\theta}^r$ of multivariate periodic functions. *Sbornik: Math.* 197(1), 69–93.
- 18 Bekmaganbetov, K.A., & Nursultanov, E.D. (2009). Interpolation of Besov $B_{pt}^{\sigma q}$ and Lizorkin-Triebel $F_{pt}^{\sigma q}$ spaces. *Analysis Mathematica*. 35(3), 169–188. <https://doi.org/10.1007/s10476-009-0301-3>
- 19 Bekmaganbetov, K.A., & Nursultanov, E.D. (2009). Embedding theorems for anisotropic Besov spaces $B_{pr}^{\alpha q}([0, 2\pi)^n)$. *Izvestiia: Mathematics*, 73(4), 655–668. <https://doi.org/10.4213/im2741>
- 20 Bekmaganbetov, K.A., & Toleugazy, Ye. (2016). Order of the orthoprojection widths of the anisotropic Nikol'skii-Besov classes in the anisotropic Lorentz space. *Eurasian Math. J.*, 7(3), 8–16.
- 21 Bekmaganbetov, K.A., Kervenev, K.Ye., & Toleugazy, Ye. (2020). Interpolation theorem for Nikol'skii-Besov type spaces with mixed metric. *Bulletin of the Karaganda university. Mathematics series*, 4(100), 32–41. <https://doi.org/10.31489/2020M4/33-42>
- 22 Bekmaganbetov, K.A., Kervenev, K.Ye., & Toleugazy, Ye. (2021). The embedding theorems for Nikol'skii-Besov spaces with generalized mixed smoothness. *Bulletin of the Karaganda university. Mathematics series*, 4(104), 28–34. <https://doi.org/10.31489/2021M4/28-34>
- 23 Bekmaganbetov, K.A., Kervenev, K.Ye., & Toleugazy, Ye. (2022). The theorems about traces and extensions for functions from Nikol'skii-Besov spaces with generalized mixed smoothness. *Bulletin of the Karaganda university. Mathematics series*, 4(108), 42–50. <https://doi.org/10.31489/2022M4/42-50>

- 24 Nursultanov, E.D. (2004). Interpoliatsionnye teoremy dlia anizotropnykh funktsionalnykh prost-
ranstv i ikh prilozheniia [Interpolation theories for anisotropic function spaces and their applica-
tions]. *Doklady Akademii nauk SSSR — Soviet Math. Dokl.*, 394(1), 22–25 [in Russian].

*Author Information**

Yerzhan Toleugazy — PhD, Senior lecturer, M.V. Lomonosov Moscow State University, Kazakhstan
Branch, 11 Kazhymukan street, Astana, 100008, Kazakhstan; e-mail: *toleugazy-y@yandex.ru*

Kabylgazy Yerezhovich Kervenev (*corresponding author*) — Master, Senior lecturer, Karaganda
Buketov University, 28 Universitetskaya street, Karaganda, 100028, Kazakhstan; e-mail: *kervenev@bk.ru*

Buketov University

*The author's name is presented in the order: First, Middle and Last Names.